



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07K 14/605, A61K 38/26</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/43708 (43) International Publication Date: 2 September 1999 (02.09.99)</p>
<p>(21) International Application Number: PCT/DK99/00086 (22) International Filing Date: 25 February 1999 (25.02.99) (30) Priority Data: 0274/98 27 February 1998 (27.02.98) DK 60/084,357 5 May 1998 (05.05.98) US (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): KNUDSEN, Liselotte, Bjerre [DK/DK]; Valby Langgade 49A, 1. tv., DK-2500 Valby (DK). HUUSFELDT, Per, Olaf [DK/DK]; Applebys Plads 27,5. mf., DK-1411 Copenhagen K (DK). NIELSEN, Per, Franklin [DK/DK]; Dalsø Park 59, DK-3500 Værløse (DK). MADSEN, Kjeld [DK/DK]; Nyvestergårdsvej 3, DK-3500 Værløse (DK). (74) Common Representative: NOVO NORDISK A/S; Novo Allé, DK-2880 Bagsvaerd (DK).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: GLP-1 DERIVATIVES OF GLP-1 AND EXENDIN WITH PROTRACTED PROFILE OF ACTION</p>		
<p>(57) Abstract</p> <p>The present invention relates to derivatives exendin and of GLP-1(7-C), wherein C is 35 or 36, which derivatives have just one lipophilic substituent which is attached to the C-terminal amino acid residue.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

GLP-1 DERIVATIVES OF GLP-1 AND EXENDIN WITH PROTRACTED PROFILE OF ACTION

FIELD OF THE INVENTION

5 The present invention relates to novel derivatives of human glucagon-like peptide-1 (GLP-1) and fragments thereof and analogues of such fragments which have a protracted profile of action and to methods of making and using them. The invention furthermore relates to novel derivatives of exendin and the uses of such derivatives.

10

BACKGROUND OF THE INVENTION

Peptides are widely used in medical practice, and since they can be produced by recombinant DNA technology it can be expected that their importance will increase also in the years to come. When native peptides or analogues thereof are used in therapy it is generally found that they have a high clearance. A high clearance of a therapeutic agent is inconvenient in cases where it is desired to maintain a high blood level thereof over a prolonged period of time since repeated administrations will then be necessary. Examples of peptides which have a high clearance are: ACTH, corticotropin-releasing factor, angiotensin, calcitonin, insulin, glucagon, glucagon-like peptide-1, glucagon-like peptide-2, insulin-like growth factor-1, insulin-like growth factor-2, gastric inhibitory peptide, growth hormone-releasing factor, pituitary adenylate cyclase activating peptide, secretin, enterogastrin, somatostatin, somatotropin, somatomedin, parathyroid hormone, thrombopoietin, erythropoietin, hypothalamic releasing factors, prolactin, thyroid stimulating hormones, endorphins, enkephalins, vasopressin, oxytocin, opioids and analogues thereof, superoxide dismutase, interferon, asparaginase, arginase, arginine deaminase, adenosine deaminase and ribonuclease. In some cases it is possible to influence the release profile of peptides by applying suitable pharmaceutical compositions, but this approach has various shortcomings and is not generally applicable.

30 The hormones regulating insulin secretion belong to the so-called enteroinsular axis, designating a group of hormones, released from the gastrointestinal mucosa in response to the presence and absorption of nutrients in the gut, which promote an early and potentiated release of insulin. The enhancing effect on insulin secretion, the so-called incretin effect, is probably essential for a normal glucose tolerance. Many of the gastrointestinal hormones, including gastrin and secretin (cholecystokinin is not insulinotropic in man), are insulinotro-

35

pic, but the only physiologically important ones, those that are responsible for the incretin effect, are the glucose-dependent insulinotropic polypeptide, GIP, and glucagon-like peptide-1 (GLP-1). Because of its insulinotropic effect, GIP, isolated in 1973 (1) immediately attracted considerable interest among diabetologists. However, numerous investigations carried out during the following years clearly indicated that a defective secretion of GIP was not involved in the pathogenesis of insulin dependent diabetes mellitus (IDDM) or non insulin-dependent diabetes mellitus (NIDDM) (2). Furthermore, as an insulinotropic hormone, GIP was found to be almost ineffective in NIDDM (2). The other incretin hormone, GLP-1 is the most potent insulinotropic substance known (3). Unlike GIP, it is surprisingly effective in stimulating insulin secretion in NIDDM patients. In addition, and in contrast to the other insulinotropic hormones (perhaps with the exception of secretin) it also potently inhibits glucagon secretion. Because of these actions it has pronounced blood glucose lowering effects particularly in patients with NIDDM.

GLP-1, a product of the proglucagon (4), is one of the youngest members of the secretin-VIP family of peptides, but is already established as an important gut hormone with regulatory function in glucose metabolism and gastrointestinal secretion and metabolism (5). The glucagon gene is processed differently in the pancreas and in the intestine. In the pancreas (9), the processing leads to the formation and parallel secretion of 1) glucagon itself, occupying positions 33-61 of proglucagon (PG); 2) an N-terminal peptide of 30 amino acids (PG (1-30)) often called glicentin-related pancreatic peptide, GRPP (10, 11); 3) a hexapeptide corresponding to PG (64-69); 4) and, finally, the so-called major proglucagon fragment (PG (72-158)), in which the two glucagon-like sequences are buried (9). Glucagon seems to be the only biologically active product. In contrast, in the intestinal mucosa, it is glucagon that is buried in a larger molecule, while the two glucagon-like peptides are formed separately (8). The following products are formed and secreted in parallel: 1) glicentin, corresponding to PG (1-69), with the glucagon sequence occupying residues Nos. 33-61 (12); 2) GLP-1(7-36)amide (PG (78-107))amide (13), not as originally believed PG (72-107)amide or 108, which is inactive). Small amounts of C-terminally glycine-extended but equally bioactive GLP-1(7-37), (PG (78-108)) are also formed (14); 3) intervening peptide-2 (PG (111-122)amide) (15); and 4) GLP-2 (PG (126-158)) (15, 16). A fraction of glicentin is cleaved further into GRPP (PG (1-30)) and oxyntomodulin (PG (33-69)) (17, 18). Of these peptides, GLP-1, has the most conspicuous biological activities.

Being secreted in parallel with glicentin/enteroglucagon, it follows that the many studies of enteroglucagon secretion (6, 7) to some extent also apply to GLP-1 secretion, but GLP-1 is metabolised more quickly with a plasma half-life in humans of 2 min (19). Carbohydrate or fat-rich meals stimulate secretion (20), presumably as a result of direct interaction of yet

unabsorbed nutrients with the microvilli of the open-type L-cells of the gut mucosa. Endocrine or neural mechanisms promoting GLP-1 secretion may exist but have not yet been demonstrated in humans.

5 The incretin function of GLP-1(29-31) has been clearly illustrated in experiments with the GLP-1 receptor antagonist, exendin 9-39, which dramatically reduces the incretin effect elicited by oral glucose in rats (21, 22). The hormone interacts directly with the β -cells via the GLP-1 receptor (23) which belongs to the glucagon/VIP/calcitonin family of G-protein-coupled 7-transmembrane spanning receptors. The importance of the GLP-1 receptor in regulating insulin secretion was illustrated in recent experiments in which a targeted disruption of the GLP-1 receptor gene was carried out in mice. Animals homozygous for the disruption had greatly deteriorated glucose tolerance and fasting hyperglycaemia, and even heterozygous animals were glucose intolerant (24). The signal transduction mechanism (25) primarily involves activation of adenylate cyclase, but elevations of intracellular Ca^{2+} are also essential (25, 26). The action of the hormone is best described as a potentiation of glucose stimulated insulin release (25), but the mechanism that couples glucose and GLP-1 stimulation is not known. It may involve a calcium-induced calcium release (26, 27). As already mentioned, the insulinotropic action of GLP-1 is preserved in diabetic β -cells. The relation of the latter to its ability to convey "glucose competence" to isolated insulin-secreting cells (26, 28), which respond poorly to glucose or GLP-1 alone, but fully to a combination of the two, is also not known. Equally importantly, however, the hormone also potently inhibits glucagon secretion (29). The mechanism is not known, but seems to be paracrine, via neighbouring insulin or somatostatin cells (25). Also the glucagonostatic action is glucose-dependent, so that the inhibitory effect decreases as blood glucose decreases. Because of this dual effect, if the plasma GLP-1 concentrations increase either by increased secretion or by exogenous infusion the molar ratio of insulin to glucagon in the blood that reaches the liver via the portal circulation is greatly increased, whereby hepatic glucose production decreases (30). As a result blood glucose concentrations decrease. Because of the glucose dependency of the insulinotropic and glucagonostatic actions, the glucose lowering effect is self-limiting, and the hormone, therefore, does not cause hypoglycaemia regardless of dose (31). The effects are preserved in patients with diabetes mellitus (32), in whom infusions of slightly supra-physiological doses of GLP-1 may completely normalise blood glucose values in spite of poor metabolic control and secondary failure to sulphonylurea (33). The importance of the glucagonostatic effect is illustrated by the finding that GLP-1 also lowers blood glucose in type-1 diabetic patients without residual β -cell secretory capacity (34).

In addition to its effects on the pancreatic islets, GLP-1 has powerful actions on the gastrointestinal tract. Infused in physiological amounts, GLP-1 potently inhibits pentagastrin-

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.