


# Principles of Optics

ELECTROMAGNETIC THEORY OF PROPAGATION  
INTERFERENCE AND DIFFRACTION OF LIGHT

Sixth Edition

MAX BORN & EMIL WOLF



Pergamon Press

# Principles of Optics

*Electromagnetic Theory of Propagation,  
Interference and Diffraction of Light*

by

MAX BORN

M.A., Dr.Phil., F.R.S.

Nobel Laureate

*Formerly Professor at the Universities of Göttingen and Edinburgh*

and

EMIL WOLF

Ph.D., D.Sc.

*Professor of Physics, University of Rochester, N.Y.*

*with contributions by*

A. B. BHATIA, P. C. CLEMMOW, D. GABOR, A. R. STOKES,

A. M. TAYLOR, P. A. WAYMAN and W. L. WILCOCK

SIXTH EDITION



PERGAMON PRESS

OXFORD · NEW YORK · TORONTO · SYDNEY · PARIS · FRANKFURT

U.K.	Pergamon Press Ltd., Headington Hill Hall, Oxford OX3 0BW, England
U.S.A.	Pergamon Press Inc., Maxwell House, Fairview Park, Elmsford, New York 10523, U.S.A.
CANADA	Pergamon of Canada, Suite 104, 150 Consumers Road, Willowdale, Ontario M2J 1P9, Canada
AUSTRALIA	Pergamon Press (Aust.) Pty. Ltd., P.O. Box 544, Potts Point, N.S.W. 2011, Australia
FRANCE	Pergamon Press SARL, 24 rue des Ecoles, 75240 Paris, Cedex 05, France
FEDERAL REPUBLIC OF GERMANY	Pergamon Press GmbH, 6242 Kronberg-Taunus, Hammerweg 6, Federal Republic of Germany

Copyright © 1980 Max Born and Emil Wolf  
*All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means: electronic, electrostatic, magnetic tape, mechanical, photocopying, recording or otherwise, without permission in writing from the copyright holders*

First edition 1959  
 Second (revised) edition 1964  
 Third (revised) edition 1965  
 Fourth (revised) edition 1970  
 Fifth (revised) edition 1975  
 Reprinted 1975, 1977  
 Sixth edition 1980

Library of Congress Cataloging in Publication Data

Born, Max  
 Principles of optics. - 6th ed.  
 I. Optics - Collected works  
 I. Title II. Wolf, Emil  
 535 QC351 80-41470

ISBN 0-08-026482-4 hardcover  
 ISBN 0-08-026481-6 flexicover

Printed in Great Britain by A. Wheaton & Co. Ltd., Exeter

The id  
of publ  
twenty  
research  
years,  
field. I  
a subst  
In plan  
develop  
book v  
restrict  
The opt  
full cor  
old bo  
subject  
relativ  
book n  
was th  
restrict  
phenom  
of mat  
mecha  
The fac  
some i  
classica  
We  
plete p  
such a  
MAXW  
In C  
effect o  
formall  
questio  
presenc  
may be  
tion of  
conside  
(Chapte  
treated  
by Prof  
A co  
follows  
additio

\* MA

Let  $c$  be the distance between the focal points  $F'_0$  and  $F_1$ . Since the image space of the first transformation coincides with the object space of the second,

$$Z_1 = Z'_0 - c, \quad Y_1 = Y'_0. \quad (23)$$

Elimination of the coordinates of the intermediate space from (22) by means of (23) gives

$$\left. \begin{aligned} Y'_1 &= \frac{Z'_1 Y_1}{f'_1} = \frac{Z'_1 Y'_0}{f'_1} = \frac{Z'_1 f_0 Y_0}{f'_1 Z_0} = \frac{f_0 f_1 Y_0}{f_0 f'_0 - c Z_0}, \\ Z'_1 &= \frac{f_1 f'_1}{Z_1} = \frac{f_1 f'_1}{Z'_0 - c} = \frac{f_1 f'_1}{f_0 f'_0 - c} = \frac{f_1 f'_1 Z_0}{f_0 f'_0 - c Z_0} \end{aligned} \right\} \quad (24)$$

Let

$$\left. \begin{aligned} Y &= Y_0, & Z &= Z_0 - \frac{f_0 f'_0}{c}, \\ Y' &= Y'_1, & Z' &= Z'_1 + \frac{f_1 f'_1}{c} \end{aligned} \right\} \quad (25)$$

Equations (25) express a change of coordinates, the origins of the two systems being shifted by distances  $f'_0 f_0 / c$  and  $-f_1 f'_1 / c$  respectively in the  $Z$ -direction. In terms of these variables, the equations of the combined transformation become

$$\frac{Y'}{Y} = \frac{f}{Z} = \frac{Z'}{f'}, \quad (26)$$

where

$$f = -\frac{f_0 f_1}{c}, \quad f' = \frac{f'_0 f'_1}{c}. \quad (27)$$

The distance between the origins of the new and the old systems of coordinates, i.e. the distances  $\delta = F_0 F$  and  $\delta' = F'_1 F'$  of the foci of the equivalent transformation from the foci of the individual transformations are seen from (25) to be

$$\delta = \frac{f_0 f'_0}{c}, \quad \delta' = -\frac{f_1 f'_1}{c}. \quad (28)$$

If  $c = 0$ , then  $f = f' = \infty$  so that the equivalent collineation is telescopic. The equations (24) then reduce to

$$\left. \begin{aligned} Y'_1 &= \frac{f_1}{f'_0} Y_0, \\ Z'_1 &= \frac{f_1 f'_1}{f_0 f'_0} Z_0; \end{aligned} \right\} \quad (29)$$

the constants  $\alpha$  and  $\beta$  in (18) of the equivalent transformation, are therefore

$$\alpha = \frac{f_1}{f'_0}, \quad \beta = \frac{f_1 f'_1}{f_0 f'_0}. \quad (30)$$

The angular magnification is now

$$\frac{\tan \gamma'}{\tan \gamma} = \frac{\alpha}{\beta} = \frac{f_0}{f'_1}. \quad (31)$$

If one or both of the transformations are telescopic, the above considerations must be somewhat modified.

We shall now tions. In this lie in the inn powers of off. be neglected.

#### 4.4.1 Refra

Consider a pe two homoger rays in both origin will be symmetry.

Let  $P_0(x_0, y_0)$  and  $P_1(x_1, y_1)$  respectively. § 4.1 (40), ar the two rays

where, acco

r being the

Let us ex to lie in th  $P_1$  will ther rays, so the Now for

and substit

$y_1 =$

\* As befo used. The v the Teachin

## 4.4 GAUSSIAN OPTICS

We shall now study the elementary properties of lenses, mirrors, and their combinations. In this elementary theory only those points and rays will be considered which lie in the immediate neighbourhood of the axis; terms involving squares and higher powers of off-axis distances, or of the angles which the rays make with the axis, will be neglected. The resulting theory is known as Gaussian optics.\*

## 4.4.1 Refracting surface of revolution

Consider a pencil of rays incident on a refracting surface of revolution which separates two homogeneous media of refractive indices  $n_0$  and  $n_1$ . To begin with, points and rays in both media will be referred to the same Cartesian reference system, whose origin will be taken at the pole  $O$  of the surface, with the  $z$ -direction along the axis of symmetry.

Let  $P_0(x_0, y_0, z_0)$  and  $P_1(x_1, y_1, z_1)$  be points on the incident and on the refracted ray respectively. Neglecting terms of degree higher than first, it follows from § 4.1 (29), § 4.1 (40), and § 4.1 (44) that the coordinates of these points and the components of the two rays are connected by the relations

$$\left. \begin{aligned} x_0 - \frac{p_0}{n_0} z_0 &= \frac{\partial T^{(2)}}{\partial p_0} = 2\mathcal{A}p_0 + \mathcal{C}p_1, \\ x_1 - \frac{p_1}{n_1} z_1 &= -\frac{\partial T^{(2)}}{\partial p_1} = -2\mathcal{B}p_1 - \mathcal{C}p_0, \end{aligned} \right\} \quad (1a)$$

$$\left. \begin{aligned} y_0 - \frac{q_0}{n_0} z_0 &= \frac{\partial T^{(2)}}{\partial q_0} = 2\mathcal{A}q_0 + \mathcal{C}q_1, \\ y_1 - \frac{q_1}{n_1} z_1 &= -\frac{\partial T^{(2)}}{\partial q_1} = -2\mathcal{B}q_1 - \mathcal{C}q_0, \end{aligned} \right\} \quad (1b)$$

where, according to § 4.1 (45),

$$\mathcal{A} = \mathcal{B} = \frac{1}{2} \frac{r}{n_1 - n_0}, \quad \mathcal{C} = -\frac{r}{n_1 - n_0}, \quad (2)$$

$r$  being the paraxial radius of curvature of the surface.

Let us examine under what conditions all the rays from  $P_0$  (which may be assumed to lie in the plane  $x = 0$ ) will, after refraction, pass through  $P_1$ . The coordinates of  $P_1$  will then depend only on the coordinates of  $P_0$  and not on the components of the rays, so that when  $q_1$  is eliminated from (1b),  $q_0$  must also disappear.

Now from the first equation (1b)

$$q_1 = \frac{1}{\mathcal{C}} \left( y_0 - q_0 \left( 2\mathcal{A} + \frac{1}{n_0} z_0 \right) \right), \quad (3)$$

and substituting this into the second equation, we obtain

$$y_1 = - \left( 2\mathcal{B} - \frac{1}{n_1} z_1 \right) \frac{1}{\mathcal{C}} y_0 + \left[ \frac{1}{\mathcal{C}} \left( 2\mathcal{B} - \frac{1}{n_1} z_1 \right) \left( 2\mathcal{A} + \frac{1}{n_0} z_0 \right) - \mathcal{C} \right] q_0. \quad (4)$$

\* As before, the usual sign convention of analytical geometry (Cartesian sign convention) is used. The various sign conventions employed in practice are very fully discussed in a Report on the Teaching of Geometrical Optics published by the Physical Society (London) in 1934.

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.