

[54] 2-(SECONDARY AMINOALKOXYMETHYL) DIHYDROPYRIDINE DERIVATIVES AS ANTI-ISCHAEMIC AND ANTIHYPERTENSIVE AGENTS

[75] Inventors: Simon F. Campbell, Deal; Peter E. Cross, Canterbury; John K. Stubbs, Deal, all of England

[73] Assignee: Pfizer Inc., New York, N.Y.

[21] Appl. No.: 576,982

[22] Filed: Feb. 3, 1984

**Related U.S. Application Data**

[63] Continuation-in-part of Ser. No. 463,081, Feb. 2, 1983, abandoned.

**Foreign Application Priority Data**

Mar. 11, 1982 [GB] United Kingdom ..... 8207180

[51] Int. Cl.<sup>4</sup> ..... C07D 211/90; A61K 31/455

[52] U.S. Cl. .... 514/356; 546/321

[58] Field of Search ..... 544/333; 546/321, 283, 546/274, 280, 257, 271, 167, 284, 270; 424/251, 258, 266; 514/356

**References Cited**

**U.S. PATENT DOCUMENTS**

4,430,333 2/1984 Campbell et al. .... 546/321

**FOREIGN PATENT DOCUMENTS**

318101 7/1981 European Pat. Off. .... 546/321

**OTHER PUBLICATIONS**

Schramm, M., "Novel Dihydropyridines with Positive Inotropic Action", Nature, vol. 303 (Jun. 9, 1983) pp. 535-537.

Bossert, F. et al, "4-Aryldihydropyridines", Angew. Chem. Int. Ed. Engl. 20, pp. 762-769 (1981).


Primary Examiner—Henry R. Jiles

Assistant Examiner—Dale A. Bjorkman

Attorney, Agent, or Firm—Charles J. Knuth; Albert E. Frost; James M. McManus

[57] **ABSTRACT**

A dihydropyridine compound of the formula



or a pharmaceutically acceptable acid addition salt thereof,

wherein

Y is  $-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-CH_2CH(CH_3)-$  or  $-CH_2C(CH_3)_2-$ ;

R is aryl or heteroaryl;

R<sup>1</sup> and R<sup>2</sup> are each independently C<sub>1</sub>-C<sub>4</sub> alkyl or 2-methoxyethyl; and

R<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, 2-(C<sub>1</sub>-C<sub>4</sub> alkoxy)ethyl, cyclopropylmethyl, benzyl, or  $-(CH_2)_mCOR^4$  where m is 1, 2 or 3 and


R<sup>4</sup> is hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy or  $-NR^5R^6$  where R<sup>5</sup> and R<sup>6</sup> are each independently hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl

can be employed for treating or preventing a heart condition or hypertension.

17 Claims, No Drawings



3



[R, R<sup>1</sup>, R<sup>2</sup> and Y are as defined for formula (I)].


One preferred amino-protecting group is benzyl. It is typically removed by hydrogenation, using e.g. H<sub>2</sub>/Pd on charcoal under acidic conditions in a suitable organic solvent, e.g. methanol. The acidic conditions are preferably obtained by using compound (II) in the form of an organic acid addition salt, e.g. as an oxalate or acetate salt.

A typical procedure involving the removal of a benzyl group is as follows. Compound (II) as an oxalate salt in methanol is added to a suspension of 10% pre-hydrogenated palladium on charcoal in methanol, and the mixture is then stirred under hydrogen at 50 p.s.i. for up to about 18 hours, e.g. overnight, and at room temperature. If necessary, heating at up to about 60° C. can be provided. The product can then be isolated and purified by conventional procedures.

When both Q and R<sup>3</sup> are benzyl, hydrogenation under the above conditions normally only removes one of the benzyl groups. Further hydrogenation of the resulting monobenzyl product under the above conditions with fresh catalyst can then be used to remove the remaining benzyl group.

Many of the starting materials of the formula (II) in which Q is benzyl are described and claimed in our European patent application publication No. 0060674. Typical methods to the N-benzyl starting materials of the formula (II) are as follows:

(a) The benzyl-protected intermediates (II) can be prepared by the Hantzsch synthesis, as follows:




4


In a typical procedure, the ketoester (IV) and aldehyde are heated under reflux in a suitable organic solvent, e.g. a C<sub>1</sub>-C<sub>4</sub> alkanol solvent such as ethanol, for about 15 minutes, and then the aminocrotonate (III) is added. Alternatively the aminocrotonate (III), ketoester (IV) and aldehyde can be heated together in the solvent. Preferably a small amount of a lower alkanolic acid such as acetic acid is added to neutralise the solution. The resulting solution can then be heated at 60°-130° C., preferably under reflux, until the reaction is essentially complete, typically in 24 hours or less. The product of the formula (II) can then be isolated and purified by conventional procedures.

The ketoesters (IV) are either known compounds or can be prepared by methods analogous to those of the prior art, such as the method illustrated in the Preparations hereinafter, which are essentially the method of Troostwijk and Kellogg, *J.C.S. Chem. Comm.*, 1977, page 932. Similarly the amino-crotonates (III) are either known compounds or can be prepared by conventional procedures. Also the aldehydes are either known or can be prepared by known methods.

(b) The benzyl-containing intermediates (II) can also be prepared by the following process:




The crotonate (VI) is typically prepared in situ by reaction of the corresponding acetoacetate (IV):



with ammonium acetate, e.g. by refluxing in a suitable organic solvent, e.g. a C<sub>1</sub>-C<sub>4</sub> alkanol such as ethanol, for, say, up to an hour. The crotonate (VI) is then reacted with compound (V), typically by heating in the solvent for up to about 5 hours at 60° C.-130° C., e.g. under reflux. The product (II) can then be isolated and purified by conventional procedures.

The starting materials (V) are either known compounds or may be prepared by methods analogous to those of the prior art, see e.g. *Can. J. Chem.*, 1967, 45, 1001.


The compounds of the formula (I) in which R<sup>3</sup> is H can be prepared from the corresponding phthalimido derivatives according to conventional procedures, e.g.:



The preferred primary amine is methylamine. The preferred alkali metal hydroxide is potassium hydroxide.

The reaction using methylamine is typically carried out in ethanol at room temperature, with heating if necessary. The reaction using hydrazine hydrate is typically carried out in ethanol at the reflux temperature or below. The reaction using potassium hydroxide is typically carried out at room temperature (although with heating if necessary) in tetrahydrofuran, following by the addition of the acid and heating at the reflux temperature or below. In all cases the product can be isolated conventionally.

The phthalimido starting materials can again be obtained conventionally, e.g.:




This is again the Hantzsch reaction.


Compounds of the formula (I) in which R<sup>3</sup> is H can also be purified to very high levels by reacting them with phthalic anhydride to form the phthalimido derivatives which can then be converted back to the compounds in which R<sup>3</sup> is H by the methods previously described.

To prepare compounds in which R<sup>3</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, -COOCH<sub>2</sub>CCl<sub>3</sub> can be used as the amino-protecting group. This can be removed in a conventional manner using zinc and either formic or acetic acid. The N-

protected starting materials necessary for this process can be prepared as follows:




or




Typically the reaction with 2,2,2-trichloroethyl chloroformate is carried by heating the reactants at up to reflux temperature in e.g. toluene. Many of the dialkyl-amino and N-alkyl-N-benzylamino starting materials needed to prepare these N-protected intermediates are described and claimed in our corresponding European patent application publication No. 0060674, and others can be prepared analogously.

The compounds of the formula (I) where R<sup>3</sup>=H can also be obtained from the corresponding azido compounds, the azido group being convertible to -NH<sub>2</sub> by reduction, e.g. with triphenylphosphine, or zinc and hydrochloric acid, or H<sub>2</sub>/Pd, under conventional conditions.




In a typical procedure using zinc dust, the reaction is carried out in methanol/aqueous hydrochloric acid. Heating is possible but is not generally necessary. Similarly hydrogenation can be carried out in e.g. methanol or ethanol in the presence of a catalyst such as Pd/CaCO<sub>3</sub> at room temperature.

Again the azido starting materials can be prepared by the Hantzsch synthesis under conditions similar to those previously described:




The azido-containing acetoacetates can also be obtained by conventional procedures:



Similarly the azido starting materials can also be prepared analogously to route (b) above for preparing the N-benzyl starting materials.

Some of the compounds of the invention can be prepared from other compounds of the invention by conventional techniques, e.g.:



The ability of the compounds to inhibit the movement of calcium into the cell is shown by their effectiveness in reducing the response of isolated heart tissue to an increase in calcium ion concentration in vitro. The test is performed by mounting spirally cut strips of rat aorta with one end fixed and the other attached to a force transducer. The tissue is immersed in a bath of physiological saline solution containing potassium ions at a concentration of 45 millimolar and no calcium. Calcium chloride is added to the bath with a pipette to give a final calcium ion concentration of 2 millimolar. The change in tension caused by the resulting contraction of the tissue is noted. The bath is drained and replaced with fresh saline solution and, after 45 minutes, the test is repeated with the particular compound under

test present in the saline solution. The concentration of compound required to reduce the response by 50% is recorded.

The antihypertensive activity of the compounds is also evaluated after oral administration by measuring the fall in blood pressure in spontaneously hypertensive rats or renally hypertensive dogs.

For administration to man in the curative or prophylactic treatment of cardiac conditions and hypertension, oral dosages of the compounds will be in the range of from 2-50 mg daily for an average adult patient (70 kg). Thus for a typical adult patient, individual tablets or capsules are likely to contain from 1 to 10 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier. Dosages for intravenous administration would be within the range 1 to 10 mg per single dose as required.

In a further aspect the invention provides a pharmaceutical composition comprising a compound of the formula (I), or a pharmaceutically acceptable acid addition salt thereof, together with a pharmaceutically acceptable diluent or carrier.


The invention also provides a compound of the formula (I), or a pharmaceutically acceptable acid addition salt thereof, for use in treating ischaemic heart disease, especially angina, or hypertension, in a human being.

The following Examples illustrate the invention: all temperatures are in °C.:

#### EXAMPLE 1

##### Preparation of

4-(2-chlorophenyl)-2-[2-(methylamino)ethoxymethyl]-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine, oxalate salt



A solution of 2-[2-(N-benzyl-N-methylamino)ethoxymethyl]-4-[2-chlorophenyl]-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine, oxalate salt (4.3 g) in methanol (220 ml) was added to a suspension of 10% (by weight) palladium on charcoal (0.4 g) pre-hydrogenated in methanol (50 ml). Stirring under hydrogen at 50 p.s.i. and room temperature overnight resulted in complete removal of the benzyl group. After removal of the catalyst by filtration, the methanol was removed by evaporation and the residue crystal-



# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.