



For Advanced
High School
Chemistry

Brown LeMay Bursten

CHEMISTRY

THE CENTRAL SCIENCE

Revised Eighth Edition



CD-ROM Inside

DOCKET
A L A R M

Find authenticated court documents without watermarks at docketalarm.com.

Chemistry

The Central Science

Eighth Revised Edition

Theodore L. Brown

University of Illinois at Urbana-Champaign

H. Eugene LeMay, Jr.

University of Nevada, Reno

Bruce E. Bursten

The Ohio State University

With contributions by Julia R. Burdge, University of Akron

PRENTICE HALL
Upper Saddle River, New Jersey 07458

Editor: John Challice
Development Editor/Editor in Chief, Development: Carol Trueheart
Associate Editor: Mary Hornby
Editorial Assistants: Amanda K. Griffith, Gillian Buonanno
Media Editor: Paul Draper
Editorial/Production Supervision: Bob Walters
Art Director: Joseph Sengotta
Assistant Art Director: John Christiana
Page layout: Richard Foster, Karen Noferi, Karen Stephens, Amy Peltier, Jeff Henn,
Joanne Del Ben, Donna Marie Paukovits
Art Studios: Academy Artworks/Michael Goodman/BioGrafx/Wellington
Editor in Chief: Paul F. Corey
Director of Marketing: John Tweeddale
Assistant Vice President ESM Production and Manufacturing: David W. Riccardi
Executive Managing Editor: Kathleen Schiaparelli
Art Manager: Gus Vibal
Art Editor: Karen Branson
Assistant Art Editor: Adam Velthaus
Senior Marketing Manager: Steve Sartori
Marketing Assistant: Dorothy Marrero
Director, Creative Services: Paul Belfanti
Associate Creative Director: Amy Rosen
Interior Design: Judith A. Matz-Coniglio
Manufacturing Manager: Trudy Piscioti
Photo Editor: Melinda Reo
Photo Researcher: Yvonne Gerin
Cover Illustration: © Kenneth Eward/BioGrafx, 1999
Copy Editor: Fay Ahuja

© 2002, 2000, 1997, 1994, 1991, 1988, 1985, 1981, 1977 by Prentice-Hall, Inc.
Upper Saddle River, NJ 07458

All rights reserved. No part of this book may be
reproduced, in any form or by any means,
without permission in writing from the publisher.
Printed in the United States of America
10 9 8 7 6 5 4 3 2 1

ISBN 0-13-061142-5

Prentice-Hall International (UK) Limited, *London*
Prentice-Hall of Australia Pty. Limited, *Sydney*
Prentice-Hall Canada Inc., *Toronto*
Prentice-Hall Hispanoamericana, S.A., *Mexico*
Prentice-Hall of India Private Limited, *New Delhi*
Prentice-Hall of Japan, Inc., *Tokyo*
Prentice-Hall (Singapore) Pte. Ltd., *Singapore*
Editora Prentice-Hall do Brasil, Ltda., *Rio de Janeiro*

Weak Acids

K_{a3}

4.0×10^{-7}

4.2×10^{-11}

Apply as "large" if it proton. Thus, it:

ization)

$< 10^{-2}$.

1 constants for acids from the first factor of 10^2 as polyprotic acid

37 M. The carbonic acid.

constants, K_{a1} and K_{a2} . I can be determined a monoprotic equilibrium

$[\text{CO}_3^{2-}(aq)]$

0
+x M
x M

$1.0 \times 10^{-5} M$ information that is

Solving for x , we have $x^2 = (0.0037)(4.3 \times 10^{-7}) = 1.6 \times 10^{-9}$
 $x = [\text{H}^+] = [\text{HCO}_3^-] = \sqrt{1.6 \times 10^{-9}} = 4.0 \times 10^{-5} M$

The small value of x indicates that our simplifying assumption was justified. The pH is therefore

$$\text{pH} = -\log [\text{H}^+] = -\log (4.0 \times 10^{-5}) = 4.40$$

If we were asked to solve for $[\text{CO}_3^{2-}]$, we would need to use K_{a2} . Let's illustrate that calculation. Using the values of $[\text{HCO}_3^-]$ and $[\text{H}^+]$ calculated above, and setting $[\text{CO}_3^{2-}] = y$, we have the following initial and equilibrium concentration values:



Initial	$4.0 \times 10^{-5} M$	$4.0 \times 10^{-5} M$	0
Change	$-y M$	$+y M$	$+y M$
Equilibrium	$(4.0 \times 10^{-5} - y) M$	$(4.0 \times 10^{-5} + y) M$	$y M$

Assuming that y is small compared to 4.0×10^{-5} , we have

$$K_{a2} = \frac{[\text{H}^+][\text{CO}_3^{2-}]}{[\text{HCO}_3^-]} = \frac{(4.0 \times 10^{-5})(y)}{4.0 \times 10^{-5}} = 5.6 \times 10^{-11}$$

$$y = 5.6 \times 10^{-11} M = [\text{CO}_3^{2-}]$$

The value calculated for y is indeed very small in comparison to 4.0×10^{-5} , showing that our assumption was justified. It also shows that the ionization of HCO_3^- is negligible in comparison to that of H_2CO_3 as far as production of H^+ is concerned. However, it is the *only* source of CO_3^{2-} , which has a very low concentration in the solution.

Our calculations thus tell us that in a solution of carbon dioxide in water most of the CO_2 is in the form of CO_2 or H_2CO_3 , a small fraction ionizes to form H^+ and HCO_3^- , and an even smaller fraction ionizes to give CO_3^{2-} .

PRACTICE EXERCISE


Calculate the pH and concentration of oxalate ion, $[\text{C}_2\text{O}_4^{2-}]$, in a 0.020 M solution of oxalic acid, $\text{H}_2\text{C}_2\text{O}_4$ (see Table 16.3). **Answers:** pH = 1.80; $[\text{C}_2\text{O}_4^{2-}] = 6.4 \times 10^{-5} M$

16.7 Weak Bases

Many substances behave as weak bases in water. Such substances react with water, removing protons from H_2O , thereby forming the conjugate acid of the base and OH^- ions:



The most commonly encountered weak base is ammonia:



The equilibrium-constant expression for this reaction can be written as

$$K = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3][\text{H}_2\text{O}]} \quad [16.32]$$

Because the concentration of water is essentially constant, the $[\text{H}_2\text{O}]$ term is incorporated into the equilibrium constant, giving

$$K_b = K[\text{H}_2\text{O}] = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} \quad [16.33]$$

The constant K_b is called the **base-dissociation constant**, by analogy with the acid-dissociation constant, K_a , for weak acids. The constant K_b always refers to the equi-

TABLE 16.4 Some Weak Bases and Their Aqueous Solution Equilibria


Base	Lewis Structure	Conjugate Acid	Equilibrium Reaction	K_b
Ammonia (NH_3)		NH_4^+	$\text{NH}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$	1.8×10^{-5}
Pyridine ($\text{C}_5\text{H}_5\text{N}$)		$\text{C}_5\text{H}_5\text{NH}^+$	$\text{C}_5\text{H}_5\text{N} + \text{H}_2\text{O} \rightleftharpoons \text{C}_5\text{H}_5\text{NH}^+ + \text{OH}^-$	1.7×10^{-9}
Hydroxylamine (H_2NOH)		H_3NOH^+	$\text{H}_2\text{NOH} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{NOH}^+ + \text{OH}^-$	1.1×10^{-8}
Methylamine (NH_2CH_3)		NH_3CH_3^+	$\text{NH}_2\text{CH}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NH}_3\text{CH}_3^+ + \text{OH}^-$	4.4×10^{-4}
Hydrosulfide ion (HS^-)		H_2S	$\text{HS}^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{S} + \text{OH}^-$	1.8×10^{-7}
Carbonate ion (CO_3^{2-})		HCO_3^-	$\text{CO}_3^{2-} + \text{H}_2\text{O} \rightleftharpoons \text{HCO}_3^- + \text{OH}^-$	1.8×10^{-4}
Hypochlorite ion (ClO^-)		HClO	$\text{ClO}^- + \text{H}_2\text{O} \rightleftharpoons \text{HClO} + \text{OH}^-$	3.3×10^{-7}

librium in which a base reacts with H_2O to form the conjugate acid and OH^- . Table 16.4 \blacktriangle lists the names, formulas, Lewis structures, equilibrium reactions, and values of K_b for several weak bases in water. Appendix D includes a more extensive list. Notice that these bases contain one or more lone pairs of electrons. A lone pair is necessary to form the bond with H^+ . Notice also that in the neutral molecules the lone pairs are on nitrogen atoms and that the other bases are anions derived from weak acids.

SAMPLE EXERCISE 16.14


Calculate the concentration of OH^- in a 0.15 M solution of NH_3 .

Solution We use essentially the same procedure here as used in solving problems involving the ionization of weak acids. The first step is to write the ionization reaction and the corresponding equilibrium-constant (K_b) expression:



$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = 1.8 \times 10^{-5}$$

We then tabulate the equilibrium concentrations involved in the equilibrium:



Initial	0.15 M	—	0	0
Change	$-x$ M	—	$+x$ M	$+x$ M
Equilibrium	$(0.15 - x)$ M	—	x M	x M

(Notice the equilibrium-constant expression)

Because K_b compared 0.15 M. The

Notice that 0.15 M. The

PRACTICE
Which of the following is pyridine?

Types of V

How can we be able to behave first category pair of electrons including all These substances amines. In with a bond NH_3 with CH_3NH_2), forming an

$\text{H}-\ddot{\text{N}}-\text{H}$
I

The chemical CH_3NH_3^+ .

The weak acids nitrite, NaClO ion is always the ClO^- ion. Consequently, the

ClO^-

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.