

Compilation and alignment of DNA polymerase sequences

Junetsu Ito and Dan K. Braithwaite

Department of Microbiology and Immunology, College of Medicine, University of Arizona Health Sciences Center, Tucson, AZ 85724, USA

Received May 29, 1991; Revised and Accepted July 9, 1991

INTRODUCTION

More than 40 different DNA polymerases, including some putative DNA polymerase sequences deduced from nucleotide sequence data, have recently been reported (1-39). The amino acid sequences of these DNA polymerases have been aligned and partial homologous regions identified by many investigators (2-4,9,10,12-25,27-36,42-51). Based on the segmental amino acid sequence similarities, DNA polymerases have been classified into two major groups; *E. coli* DNA polymerase I-Type and eukaryotic DNA polymerase α -Type (14,44,47,48,51), or family A DNA polymerases and family B DNA polymerases (4,9,50). As the number of DNA polymerase sequences increases, the classification of DNA polymerases becomes increasingly ambiguous. For example, DNA polymerase delta of yeast was shown to have amino acid sequence similarity to the α -Type DNA polymerases (17). It has become necessary to establish a unified classification of DNA polymerases. Here we propose to classify DNA polymerases into families A, B, and C (Figure 1: A, B, and C), according to the amino acid sequence homologies with *E. coli* DNA polymerases I, II, and III, respectively. As new and different prokaryotic and eukaryotic DNA polymerases are identified, the number of families can easily be expanded by using additional letters of the alphabet (i.e., D, E, etc.).

The bacterium *E. coli* (strain K12) contains three distinct DNA polymerases I, II, and III (52). *E. coli* DNA polymerase I, the first DNA polymerase discovered, is specified by the *polA* gene (52). *E. coli* DNA polymerase II, encoded by the *polB* gene, was recently sequenced and found to be identical to the *dna* gene, a DNA damage inducible gene whose expression is regulated by the SOS system in *E. coli* (8,53). Amino acid sequence alignment shows that *E. coli* DNA polymerase II has significant homology with family B (α -Type) DNA polymerases (8,53,54).

E. coli DNA polymerase III is a multisubunit enzyme encoded by various *dna* genes (55); the DNA polymerizing α -subunit encoded by the *polC* (*dnaE*) gene (56) and the 3' \rightarrow 5' exonuclease performing ϵ -subunit encoded by the *dnaQ* gene (57). The α -subunit of *E. coli* DNA polymerase III exhibits an extensive homology with the corresponding α -subunit of *Salmonella typhimurium* DNA polymerase III (35); and both show significant homology to *Bacillus subtilis* DNA polymerase III, a single-polypeptide encoded by the *polC* gene (36).

In summary, family A DNA polymerases are named for their homology to the product of the *polA* gene encoding *E. coli* DNA polymerase I; family B DNA polymerases are named for their

homology to the product of the *polB* gene encoding *E. coli* DNA polymerase II; and family C DNA polymerases are named for their homology to the product of the *polC* gene encoding *E. coli* DNA polymerase III.

The eukaryotic DNA polymerase β , the smallest known DNA polymerase, does not have homology with those of any of the DNA polymerase families described above. Instead, DNA polymerase β has homology with terminal transferases (37). This β group we will call family X (Figure 1D). The classification and original reference(s) for the amino acid sequences of each DNA polymerase are shown in Table 1.

All of the family A DNA polymerases, except for yeast mitochondrial DNA polymerase I, are prokaryotic and are very sensitive to dideoxynucleotide inhibitors, and therefore are useful enzymes for DNA sequencing by the chain-termination method (58). The family A DNA polymerases are resistant to aphidicolin. The family B DNA polymerases are quite extensive in number and variety. Most of the family B DNA polymerases, if not all, are sensitive to aphidicolin and relatively resistant to dideoxynucleotide inhibitors. Most of the family B DNA polymerases, except for pAI2 (33) and yeast DNA polymerase II (16), contain the highly conserved amino acid sequence motif YGDTD, which has been suggested to form part of the dNTP binding site. Amino acid substitutions in this conserved sequence resulted in defects in the DNA polymerase activity without affecting the 3' \rightarrow 5' exonuclease activity (59,60,61). The family C DNA polymerases are major bacterial replicative DNA polymerases which do not have appreciable homology with those of family A and B DNA polymerases. *B. subtilis* DNA polymerase III is a single polypeptide that is highly sensitive to hydroxyphenylazouracil (62). It is anticipated that the number of sequenced family C DNA polymerases will increase rapidly, since all of the aerobic bacteria may contain a member of this family of DNA polymerases.

SEQUENCE ALIGNMENT

The 37 complete DNA polymerase sequences and 3 complete terminal deoxynucleotidyltransferase (TDT) sequences are listed in 4 groups; the family A DNA polymerases, the family B DNA polymerases (including the alignment), and the family C DNA polymerases are very similar to the family C DNA polymerases not

Columbia Ex. 2089
 Illumina, Inc. v. The Trustees
 of Columbia University in the
 City of New York
 152020 0000 01065

Compilation and alignment of DNA polymerase sequences

Junetsu Ito and Dan K. Braithwaite

Department of Microbiology and Immunology, College of Medicine, University of Arizona Health Sciences Center, Tucson, AZ 85724, USA

Received May 29, 1991; Revised and Accepted July 9, 1991

INTRODUCTION

More than 40 different DNA polymerases, including some putative DNA polymerase sequences deduced from nucleotide sequence data, have recently been reported (1–39). The amino acid sequences of these DNA polymerases have been aligned and partial homologous regions identified by many investigators (2–4,9,10,12–25,27–36,42–51). Based on the segmental amino acid sequence similarities, DNA polymerases have been classified into two major groups; *E. coli* DNA polymerase I-Type and eukaryotic DNA polymerase α -Type (14,44,47,48,51), or family A DNA polymerases and family B DNA polymerases (4,9,50). As the number of DNA polymerase sequences increases, the classification of DNA polymerases becomes increasingly ambiguous. For example, DNA polymerase delta of yeast was shown to have amino acid sequence similarity to the α -Type DNA polymerases (17). It has become necessary to establish a unified classification of DNA polymerases. Here we propose to classify DNA polymerases into families A, B, and C (Figure 1: A, B, and C), according to the amino acid sequence homologies with *E. coli* DNA polymerases I, II, and III, respectively. As new and different prokaryotic and eukaryotic DNA polymerases are identified, the number of families can easily be expanded by using additional letters of the alphabet (i.e., D, E, etc.).

The bacterium *E. coli* (strain K12) contains three distinct DNA polymerases I, II, and III (52). *E. coli* DNA polymerase I, the first DNA polymerase discovered, is specified by the *polA* gene (52). *E. coli* DNA polymerase II, encoded by the *polB* gene, was recently sequenced and found to be identical to the *dna* gene, a DNA damage inducible gene whose expression is regulated by the SOS system in *E. coli* (8,53). Amino acid sequence alignment shows that *E. coli* DNA polymerase II has significant homology with family B (α -Type) DNA polymerases (8,53,54).

E. coli DNA polymerase III is a multisubunit enzyme encoded by various *dna* genes (55); the DNA polymerizing α -subunit encoded by the *polC* (*dnaE*) gene (56) and the 3' → 5' exonuclease performing ϵ -subunit encoded by the *dnaQ* gene (57). The α -subunit of *E. coli* DNA polymerase III exhibits an extensive homology with the corresponding α -subunit of *Salmonella typhimurium* DNA polymerase III (35); and both show significant homology to *Bacillus subtilis* DNA polymerase III, a single-polypeptide encoded by the *polC* gene (36).

In summary, family A DNA polymerases are named for their homology to the product of the *polA* gene encoding *E. coli* DNA polymerase I; family B DNA polymerases are named for their

homology to the product of the *polB* gene encoding *E. coli* DNA polymerase II; and family C DNA polymerases are named for their homology to the product of the *polC* gene encoding *E. coli* DNA polymerase III.

The eukaryotic DNA polymerase β , the smallest known DNA polymerase, does not have homology with those of any of the DNA polymerase families described above. Instead, DNA polymerase β has homology with terminal transferases (37). This β group we will call family X (Figure 1D). The classification and original reference(s) for the amino acid sequences of each DNA polymerase are shown in Table 1.

All of the family A DNA polymerases, except for yeast mitochondrial DNA polymerase I, are prokaryotic and are very sensitive to dideoxynucleotide inhibitors, and therefore are useful enzymes for DNA sequencing by the chain-termination method (58). The family A DNA polymerases are resistant to aphidicolin. The family B DNA polymerases are quite extensive in number and variety. Most of the family B DNA polymerases, if not all, are sensitive to aphidicolin and relatively resistant to dideoxynucleotide inhibitors. Most of the family B DNA polymerases, except for pAI2 (33) and yeast DNA polymerase II (16), contain the highly conserved amino acid sequence motif **YGDTD**, which has been suggested to form part of the dNTP binding site. Amino acid substitutions in this conserved sequence resulted in defects in the DNA polymerase activity without affecting the 3' → 5' exonuclease activity (59,60,61). The family C DNA polymerases are major bacterial replicative DNA polymerases which do not have appreciable homology with those of family A and B DNA polymerases. *B. subtilis* DNA polymerase III is a single polypeptide that is highly sensitive to hydroxyphenylazouracil (62). It is anticipated that the number of sequenced family C DNA polymerases will increase rapidly, since all of the aerobic bacteria may contain a member of this family of DNA polymerases.

SEQUENCE ALIGNMENT

The 37 complete DNA polymerase sequences and 3 complete terminal deoxynucleotidyltransferase (TDT) sequences are listed in 4 groups; the family A DNA polymerases, the family B DNA polymerases, the family C DNA polymerases, and family X DNA polymerases (including TDTs). In order to limit the space needed for the alignment, we omitted DNA polymerase sequences that are very similar to the prototype DNA polymerase. The DNA polymerases not shown include: herpes virus type-2 (63),

adenovirus type-5 (64), bacteriophage T3 (65), and bacteriophage PZA (66).

ACCURACY OF SEQUENCE DATA

Whenever a sequence ambiguity existed in a published sequence, we contacted the authors to obtain the updated sequence information. We found that a few published amino acid sequences differ at one or more positions from their GenBank/EMBL entry. Again, we have communicated with the primary author to confirm the correct sequences.

The multiple alignment of the amino acid sequences was obtained by a series of pairwise alignments combined and adjusted by eye into larger and larger subsets of similar sequences. The process of combining and adjusting by eye was aided by modified versions of the MOTIF program (67) and the ALIGN program (68). The GAP and BESTFIT programs, from UWGCG (University of Wisconsin Genetic Computer Group) (69), initially generated the pairwise alignments, adjusted for maximum alignment that allowed for a considerable number of gaps. We then compressed these alignments by eye to give a more contiguous alignment. The alignment of the sequences for optimal similarity is straightforward in the areas of relatively conserved structure, but is much more arbitrary in the more varied sequence areas. The alignment of the varied areas should therefore be regarded as less than optimal in view of the difficulties concerned with multiple alignments in these areas.

Finally, we invite further correction from readers, and welcome suggested revisions and alternative alignments.

ACKNOWLEDGEMENTS

This work was supported by grant GM28013 from the National Institutes of Health and by grant NP-704 from the American Cancer Society.

REFERENCES

- Joyce, C.M., Kelley, W.S. and Grindley N.D.F. (1982) *J. Biol. Chem.* **257**, 1958–1964.
- Lopez, P., Martinez, S., Diaz, A., Espinosa, M. and Lacks, S.A. (1989) *J. Biol. Chem.* **264**, 4255–4263.
- Lawyer, F.C., Stoffel, S., Saiki, R.K., Myambo, K., Drummond, R. and Gelfand, D.H. (1989) *J. Biol. Chem.* **264**, 6427–6437.
- Leavitt, M.C. and Ito, J. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**, 4465–4469.
- Dunn, J.J. and Studier, F.W. (1983) *J. Mol. Biol.* **166**, 477–535.
- Rådén, B. and Rutberg, L. (1984) *J. Virol.* **52**, 9–15.
- Foury, F. (1989) *J. Biol. Chem.* **264**, 20552–20560.
- Iwasaki, H., Ishino, Y., Toh, H., Nakata, A. and Shinagawa, H. (1991) *Mol. Gen. Genet.* **226**, 24–33.
- Jung, G., Leavitt, M.C., Hsieh, J.-C. and Ito, J. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 8287–8291.
- Savilahti, H. and Bamford D.H. (1987) *Gene* **57**, 121–130.
- Yoshikawa, H. and Ito, J. (1982) *Gene* **17**, 323–335.
- Matsumoto, K., Takano, H., Kim, C.I. and Hirokawa, H. (1989) *Gene* **84**, 247–255.
- Spicer, E.K., Rush, J., Fung, C., Reha-Krantz, L.J., Karam, J.D. and Konigsberg, W.H. (1988) *J. Biol. Chem.* **263**, 7478–7486.
- Wong, S.W., Wahl, A.F., Yuan, P.-M., Arai, N., Pearson, B.E., Arai, K.-i., Korn, D., Hunkapiller, M.W. and Wang, T.S.-F. (1988) *EMBO J.* **7**, 37–47.
- Pizzagalli, A., Valsasini, P., Plevani, P. and Lucchini, G. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 3772–3776.
- Morrison, A., Araki, H., Clark, A.B., Hamatake, R.K. and Sugino, A. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 1112–1116.

- Boulet, A., Simon, M., Faye, G., Bauer, G.A. and Burgers, P.M.J. (1989) *EMBO J.* **8**, 1849–1854.
- Morrison, A., Christensen, R.B., Alley, J., Beck, A.K., Bernstine, E.G., Lemontt, J.F. and Lawrence, C.W. (1989) *J. Bacteriol.* **171**, 5659–5667.
- Gibbs, J.S., Chiou, H.C., Hall, J.D., Mount, D.W., Retondo, M.J., Weller, S.K. and Coen, D.M. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **82**, 7969–7973.
- Kouzariades, T., Bankier, A.T., Satchwell, S.C., Weston, K., Tomlinson, P. and Barrell, B.G. (1987) *J. Virol.* **61**, 125–133.
- Baer, R., Bankier, A.T., Biggin, M.D., Deininger, P.L., Farrell, P.J., Gibson, T.J., Hatfull, G., Hudson, G.S., Satchwell, S.C., Séguin, C., Tuffnell, P.S. and Barrell, B.G. (1984) *Nature* **310**, 207–211.
- Davison, A.J. and Scott, J.E. (1986) *J. Gen. Virol.* **67**, 1759–1816.
- Binns, M.M., Stenzler, L., Tomley, F.M., Campbell, J. and Boursnell, M.E.G. (1987) *Nucl. Acids Res.* **15**, 6563–6573.
- Earl, P.L., Jones, E.V. and Moss, B. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 3659–3663.
- Tomalski, M.D., Wu, J. and Miller, L.K. (1988) *Virology* **167**, 591–600.
- Gingras, T.R., Sciak, D., Gelinis, R.E., Bing-Dong, J., Yen, C.E., Kelly, M.M., Bullock, P.A., Parsons, B.L., O'Neill, K.E. and Roberts, R.J. (1982) *J. Biol. Chem.* **257**, 13475–13491.
- Engler, J.A., Hoppe, M.S. and van Bree, M.P. (1983) *Gene* **21**, 145–159.
- Shu, L., Hong, J.S., Wei, Y.-f. and Engler, J.A. (1986) *Gene* **46**, 187–195.
- Paillard, M., Sederoff, R.R. and Levings, C.S. III (1985) *EMBO J.* **4**, 1125–1128.
- Stark, M.J.R., Mileham, A.J., Romanos, M.A. and Boyd, A. (1984) *Nucl. Acids Res.* **12**, 6011–6030.
- Tommasino, M., Ricci, S. and Galeotti, C.L. (1988) *Nucleic Acids Res.* **16**, 5863–5878.
- Oeser, B. and Tudzynski, P. (1989) *Mol. Gen. Genet.* **217**, 132–140.
- Kempken, F., Meinhardt, F. and Esser, K. (1989) *Mol. Gen. Genet.* **218**, 523–530.
- Tomasiewicz, H.G. and McHenry, C.S. (1987) *J. Bacteriol.* **169**, 5735–5744.
- Lancy, E.D., Lifscis, M.R., Munson, P. and Maurer, R. (1989) *J. Bacteriol.* **171**, 5581–5586.
- Hammond, R.A., Barnes, M.H., Mack, S.L., Mitchener, J.A. and Brown, N.C. (1991) *Gene* **98**, 29–36.
- Matsukage, A., Nishikawa, K., Ooi, T., Seto, Y. and Yamaguchi, M. (1987) *J. Biol. Chem.* **262**, 8960–8962.
- Abbotts, J., SenGupta, D.N., Zmudzka, B., Widen, S.G., Notario, V. and Wilson, S.H. (1988) *Biochemistry* **27**, 901–909.
- SenGupta, D.N., Zmudzka, B.Z., Kumar, P., Cobiانchi, F., Skowronski, J. and Wilson, S.H. (1986) *Biochem. Biophys. Res. Comm.* **136**, 341–347.
- Peterson, R.C., Cheung, L.C., Mattaliano, R.J., White, S.T., Chang, L.M.S. and Bollum F.J. (1985) *J. Biol. Chem.* **260**, 10495–19502.
- Koiwai, O., Yokota, T., Kageyama, T., Hirose, T., Yoshida, S. and Arai, K.-i. (1986) *Nucl. Acids Res.* **14**, 5777–5792.
- Argos, P., Tucker, A.D. and Phillipson, L. (1986) *Virology* **149**, 208–216.
- Larder, B.A., Kemp, S.D. and Darby, G. (1987) *EMBO J.* **6**, 160–175.
- Hall, J.D. (1988) *Trends Genet.* **4**, 42–46.
- Reha-Krantz, L.J. (1988) *J. Mol. Biol.* **202**, 711–724.
- Bernad, A., Zaballos, A., Salas, M. and Blanco, L. (1987) *EMBO J.* **6**, 4219–4225.
- Wang, T.S.-F., Wong, S.W. and Korn, D. (1989) *FASEB J.* **3**, 14–21.
- Bernad, A., Blanco, L., Lazaro, J.M., Martin, G. and Salas, M. (1989) *Cell* **59**, 219–228.
- Polesky, A.H., Steitz, T.A., Grindley, N.D.F. and Joyce, C.M. (1990) *J. Biol. Chem.* **265**, 14579–14591.
- Ito, J. and Braithwaite, D.K. (1990) *Nucl. Acids Res.* **18**, 6716.
- Blanco, L., Bernad, A. and Salas, M. (1991) *Nucl. Acids Res.* **19**, 955.
- Kornberg, A. (1974) DNA replication. W.H. Freeman and Co., San Francisco.
- Bonner, C.A., Hays, S., McEntee, K. and Goodman M.F. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 7663–7667.
- Chen, H., Lawrence, C.B., Bryan, S.K. and Moses, R.E. (1990) *Nucl. Acids Res.* **18**, 7185–7186.
- McHenry, C.S. (1985) *Mol. Cell. Biochem.* **66**, 71–85.
- Shepard, D., Oberfelder, R.W., Welch, M.W. and McHenry, C.S. (1984) *J. Bacteriol.* **158**, 455–459.
- Scheuermann, R., Tam, S., Burgers, P.M.J., Lu, C. and Echols, H. (1983) *Proc. Natl. Acad. Sci. U.S.A.* **80**, 7085–7089.
- Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. U.S.A.* **74**, 5463–5467.
- Dorsky, D.I. and Crumpacker, C.S. (1990) *J. Virol.* **64**, 1394–1397.
- Bernad, A., Lázaro, J.M., Salas, M. and Blanco, L. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 1112–1116.

61. Jung, G., Leavitt, M.C., Schultz, M. and Ito, J. (1990) *Biochem. Biophys. Res. Comm.* **170**, 1294–1300.
62. Neville, M.N. and Brown, N.C. (1972) *Nature New Biol.* **240**, 80–82.
63. Tsurumi, T., Maeno, K. and Nishiyama, Y. (1987) *Gene* **52**, 129–137.
64. Dekker, B.M.M. and Van Ormondt, H. (1984) *Gene* **27**, 115–120.
65. Beck, P.J., Gonzalez, S., Ward, C.L. and Molineux, I.J. (1989) *J. Mol. Biol.* **210**, 687–701.
66. Paces, V., Vlcek, C., Urbanek, P. and Hostomsky, Z. (1985) *Gene* **38**, 45–56.
67. Smith, H.O., Annau, T.M. and Chandrasegaran, S. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 826–830.
68. Doolittle, R.F. and Feng, D.-F. (1990) In Doolittle, R.F. (ed.), *Methods in Enzymol. – Molecular Evolution: Computer Analysis of Protein and Nucleic Acid Sequences*. Academic Press, New York, Vol. 183, pp. 659–669.
69. Devereux, J., Haerberli, P. and Smithies, O. (1984) *Nucl. Acids Res.* **12**, 387–395.

Classification of DNA polymerases

| A. Family A DNA polymerases | | References |
|---|---|-------------------|
| 1. Bacterial DNA polymerases | | |
| a) | <i>E. coli</i> DNA polymerase I | (1) |
| b) | <i>Streptococcus pneumoniae</i> DNA polymerase I | (2) |
| c) | <i>Thermus aquaticus</i> DNA polymerase I | (3) |
| 2. Bacteriophage DNA polymerases | | |
| a) | T5 DNA polymerase | (4) |
| b) | T7 DNA polymerase | (5) |
| c) | Spo2 DNA polymerase | (6) |
| 3. Mitochondrial DNA polymerase | | |
| | Yeast mitochondrial DNA polymerase (MIP1) | (7) |
| B. Family B DNA polymerases | | |
| 1. Bacterial DNA polymerases | | |
| | <i>E. coli</i> DNA polymerase II | (8) |
| 2. Bacteriophage DNA polymerases | | |
| a) | PRD1 DNA polymerase* | (9,10) |
| b) | φ29 DNA polymerase* | (11) |
| c) | M2 DNA polymerase* | (12) |
| d) | T4 DNA polymerase | (13) |
| 3. Eukaryotic DNA polymerases | | |
| a) | Human DNA polymerase alpha | (14) |
| b) | Yeast DNA polymerase I | (15) |
| c) | Yeast DNA polymerase II | (16) |
| d) | Yeast DNA polymerase III (delta) | (17) |
| e) | Yeast DNA polymerase Rev3 | (18) |
| 4. Viral DNA polymerases | | |
| a) | Herpes-1 DNA polymerase | (19) |
| b) | Human cytomegalovirus DNA polymerase | (20) |
| c) | Epstein-Barr virus DNA polymerase | (21) |
| d) | Varicella-Zoster virus DNA polymerase | (22) |
| e) | Fowlpox virus DNA polymerase | (23) |
| f) | Vaccinia virus DNA polymerase | (24) |
| g) | Autographa californica nuclear polyhedrosis virus (AcMNPV) DNA polymerase | (25) |
| h) | Adenovirus-2 DNA polymerase* | (26) |
| i) | Adenovirus-7 DNA polymerase* | (27) |
| j) | Adenovirus-12 DNA polymerase* | (28) |
| 5. Eukaryotic linear DNA plasmid encoded DNA polymerases | | |
| a) | S-1 maize mitochondrial DNA polymerase* | (29) |
| b) | <i>Kluyveromyces lactis</i> plasmid pGKL1 DNA polymerase* | (30) |
| c) | <i>Kluyveromyces lactis</i> plasmid pGKL2 DNA polymerase* | (31) |
| d) | <i>Claviceps purpurea</i> plasmid pCLK1 DNA polymerase* | (32) |
| e) | <i>Ascobolus immersus</i> plasmid pAI2 DNA polymerase* | (33) |
| C. Family C DNA polymerases | | |
| Bacterial replicative DNA polymerases | | |
| a) | <i>E. coli</i> DNA polymerase III α subunit | (34) |
| b) | <i>Salmonella typhimurium</i> DNA polymerase III α subunit | (35) |
| c) | <i>Bacillus subtilis</i> DNA polymerase III | (36) |
| D. Family X DNA polymerases | | |
| a) | Rat DNA polymerase β | (37) |
| b) | Human DNA polymerase β | (38,39) |
| c) | Human terminal deoxynucleotidyltransferase (TdT) | (40) |
| d) | Bovine terminal deoxynucleotidyltransferase (TdT) | (41) |
| e) | Mouse terminal deoxynucleotidyltransferase (TdT) | (41) |

Table 1. The main families and subclassifications of DNA polymerases. Those DNA polymerases marked with a star (*) are protein-primed DNA polymerases.




Figure 1A. Family A DNA polymerases—*E. coli* DNA pol. I (E.c. pol D(1)), *Streptococcus pneumoniae* DNA pol. I (S.p. pol D(2)), *Thermus aquaticus* DNA pol. I (Taq pol D(3)), T5 DNA pol. (4), T7 DNA pol. (5), Spo2 DNA pol. (6), and yeast mitochondrial DNA pol. (MIP1)(7).

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.