

## SYNTHESIS OF FLUORESCENT, PHOTOLABILE 3'-O-PROTECTED NUCLEOSIDE TRIPHOSPHATES FOR THE BASE ADDITION SEQUENCING SCHEME

Mike B. Welch and Kevin Burgess\*

Department of Chemistry, Texas A & M University, PO Box 300012,  
College Station, TX 77842-3012

**ABSTRACT:** The dansylated nucleoside triphosphates **1a** and **1b** were prepared as a prelude to investigating sequencing of DNA via a scheme that does not involve electrophoresis.

Throughput of sequence data is critical in The Human Genome Project and similar ventures. Gel electrophoresis is a "bottle-neck" in high throughput sequencing of DNA hence it is desirable to develop methods that avoid this technique. Several groups have been interested in an approach that we call, "The Base Addition Sequencing Scheme", BASS.<sup>1-6</sup> This involves modified nucleotides **I** that are: (i) incorporated by DNA replicating enzymes; (ii) spectroscopically distinct (at P\*) so that the parent base (A, T, G, and C) can be differentiated; and, (iii) labile at the 3'-position such that the 3'-hydroxyl terminus of a polynucleotide can be regenerated. This scheme may be slower and sequence less bases per experiment than existing methods but, unlike methods involving gels, there is potential for multiple experiments to be run in parallel. Such combinatorial sequencing schemes may lead to significantly improved throughput. A major challenge in development of BASS, however, is that

197

Copyright © 1999 by Marcel Dekker, Inc.

Columbia Ex. 2006  
Illumina, Inc. v. The Trustees  
of Columbia University  
in the City of New York  
IPR2020-01177

## **SYNTHESIS OF FLUORESCENT, PHOTOLABILE 3'-O-PROTECTED NUCLEOSIDE TRIPHOSPHATES FOR THE BASE ADDITION SEQUENCING SCHEME**

Mike B. Welch and Kevin Burgess\*

Department of Chemistry, Texas A & M University, PO Box 300012,  
College Station, TX 77842-3012

**ABSTRACT:** The dansylated nucleoside triphosphates **1a** and **1b** were prepared as a prelude to investigating sequencing of DNA via a scheme that does not involve electrophoresis.

Throughput of sequence data is critical in The Human Genome Project and similar ventures. Gel electrophoresis is a "bottle-neck" in high throughput sequencing of DNA hence it is desirable to develop methods that avoid this technique. Several groups have been interested in an approach that we call, "The Base Addition Sequencing Scheme", BASS.<sup>1-6</sup> This involves modified nucleotides **I** that are: (i) incorporated by DNA replicating enzymes; (ii) spectroscopically distinct (at P\*) so that the parent base (A, T, G, and C) can be differentiated; and, (iii) labile at the 3'-position such that the 3'-hydroxyl terminus of a polynucleotide can be regenerated. This scheme may be slower and sequence less bases per experiment than existing methods but, unlike methods involving gels, there is potential for multiple experiments to be run in parallel. Such combinatorial sequencing schemes may lead to significantly improved throughput. A major challenge in development of BASS, however, is that the 3'-blocking group must be





Diagram 1

removed in an aqueous medium using reagents/conditions that do not denature or modify double strand DNA. To address this issue, the focus of our efforts has been on photolabile protection. Earlier work from these labs proved that 3'-O-(2''-nitrobenzyl)adenosine triphosphate **II** could be incorporated by a DNA polymerase.<sup>1</sup> In an extension of these studies, we now describe syntheses of two systems **1a** and **1b** containing a 3'-O-(2-nitrobenzyl) group modified to carry a fluorescent label (Diagram 1).

The synthetic approach to target molecules **1a** and **1b** has parallels with chemistry developed for solid phase photolabile linkers.<sup>7,8</sup> Details of the synthetic route used are given in Scheme 1. The most difficult step in this procedure was coupling of a benzylic halide to the 3'-hydroxyl group of a nucleoside derivative. Development of conditions for this step required considerable experimentation.

Details of the synthesis shown in Scheme 1 are as follows. The allyloxycarbonyl (ALOC) protected bromide **2** was used to alkylate the vanillin derivative **3** to give the aldehyde **4**. Nitration then borohydride



**Scheme 1.** Reagents and conditions: i,  $\text{K}_2\text{CO}_3$ , KI, MeCN, reflux, 12 h (>95 %); ii,  $\text{HNO}_3$ ,  $\text{Ac}_2\text{O}$ ,  $0^\circ\text{C}$ ; iii,  $\text{NaBH}_4$ , EtOH,  $0^\circ\text{C}$ , 2 h then  $25^\circ\text{C}$ , 2 h (53 % for ii + iii); iv,  $\text{CBr}_4$ ,  $\text{PPh}_3$ , EtOAc,  $25^\circ\text{C}$ , 1 h; v, 5'-O-tert-butylidiphenylsilylnucleoside, NaI,  $\text{Bu}_4\text{NOH}$ ,  $\text{CH}_2\text{Cl}_2$ , 1 M NaOH,  $25^\circ\text{C}$ , 16 h (iv + v, 49 % for  $\text{B} = \text{A}^{\text{Bz}}$ , 88 % for  $\text{B} = \text{T}$ ); vi, cat.  $\text{Pd}(\text{PPh}_3)_4$ , THF,  $\text{HNEt}_3$ ,  $25^\circ\text{C}$ , 2.5 h; vii, dansyl-Cl,  $\text{NEt}_3$ , cat. 4-DMAP, PhMe/THF (1:1),  $25^\circ\text{C}$ , 1 h; viii,  $\text{Bu}_4\text{NF}$ , THF,  $25^\circ\text{C}$ , 11 min for T, and 10 min at  $0^\circ\text{C}$  then 1 h at  $25^\circ\text{C}$  for  $\text{A}^{\text{Bz}}$ ; (vi - viii, 69 % for  $\text{B} = \text{A}^{\text{Bz}}$ , 66 % for  $\text{B} = \text{T}$ ).

reduction of this gave the benzylic alcohol **5**, which emerged as a key intermediate in this work. Coupling of benzyl alcohol **5** to the 3'-hydroxyl group of nucleoside derivatives was difficult. After considerable experimentation, it was found that conversion to the benzylic bromide, then alkylation *under phase transfer conditions* worked whereas many obvious alternative approaches did not. Removal of the ALLOC group from the coupled products **6**, dansylation, then removal of the 5'-silyl protecting group gave the desired nucleosides **7**.

Triphosphorylation of nucleosides **7**, and of many other unnatural nucleosides prepared in these labs, is experimentally difficult and tends to give poor yields. Ultimately, we settled on the protocol developed by Eckstein and co-workers,<sup>9</sup> although none of the methods attempted were entirely satisfactory. Fortunately, only small amounts of the product are required for feasibility tests in bioassays. Debenzoylation of the adenosine derivative **7a** was performed after the triphosphorylation sequence ( $\text{NH}_4\text{OH}$ , 60 °C, 3 h). Both final products, **1a** and **1b**, were purified via chromatography, first on DEA cellulose, then by RP HPLC.

Preliminary tests<sup>1</sup> of compounds **1a** and **1b** as substrates for polymerases (Klenow, rTth DNA polymerase, and Vent (exo-) DNA polymerase) did not show evidence for incorporation. Further experiments are in progress, including modeling studies to attempt to dock **1a** and **1b** in the active sites of polymerases. The latter experiments should highlight bad interactions that must be avoided to advance this work further.

**Acknowledgments.** We thank Richard Gibbs of Baylor College of Medicine for valuable advice. Support for this work was provided by The Texas Technology Program, The Robert A. Welch Foundation, and The National Institutes of Health (HG/GM01745-01). KB thanks the NIH Research Career Development Award, and The Alfred P. Sloan Foundation for a fellowship.

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.