



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C07K 7/10, 7/34, A61K 37/02 A61K 37/28	A1	(11) International Publication Number: WO 87/ 06941 (43) International Publication Date: 19 November 1987 (19.11.87)
(21) International Application Number: PCT/US87/01005 (22) International Filing Date: 5 May 1987 (05.05.87) (31) Priority Application Number: 859,928 (32) Priority Date: 5 May 1986 (05.05.86) (33) Priority Country: US (71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; Fruit Street (Bar-3), Boston, MA 02114 (US). (72) Inventor: HABENER, Joel ; 217 Plymouth Road, Newton Highlands, MA 02161 (US). (74) Agents: GOLDSTEIN, Jorge, A. et al.; Saidman, Sterne, Kessler & Goldstein, 1225 Connecticut Avenue, N.W., Suite 300, Washington, DC 20036 (US).		(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: INSULINOTROPIC HORMONE (57) Abstract A fragment of glucagon-like peptide I (GLP-1) has been found to be an insulinotropic hormone. This insulinotropic hormone comprises amino acid residues 7-37 of GLP-1. The insulinotropic hormone is useful as a potential therapy for <i>Diabetes Mellitus</i> .		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-1-

INSULINOTROPIC HORMONE**BACKGROUND OF THE INVENTION****Field of the Invention**

This invention is directed to the discovery that certain peptide fragments of the prehormone, proglucagon, possess hormonal activities and can be used to stimulate the synthesis and secretion of the hormone, insulin. These peptide fragments are useful in therapy for the disease Diabetes mellitus.

Description of the Background Art

The endocrine secretions of the pancreatic islets are under complex control not only by blood-borne metabolites (glucose, amino acids, catecholemines, etc.), but also by local paracrine influences. The major pancreatic islet hormones (glucagon, insulin and somatostatin) interact amongst their specific cell types (A, B, and D cells, respectively) to modulate secretory responses mediated by the metabolites. Although insulin secretion is predominantly controlled by blood levels of glucose, glucagon and somatostatin stimulate and inhibit glucose-mediated insulin secretory responses, respectively. In addition to the pro-

-2-

posed interislet paracrine regulation of insulin secretion, there is evidence to support the existence of insulintropic factors in the intestine. This concept originates from the observations that glucose taken orally is a much more potent stimulant of insulin secretion than is a comparable amount of glucose given intravenously.

The human hormone, glucagon, is a 29-amino acid peptide hormone produced in the A-cells of the pancreas. The hormone belongs to a multi-gene family of structurally related peptides that include secretin, gastric inhibitory peptide, vasoactive intestinal peptide and glicentin. These peptides variously regulate carbohydrate metabolism, gastrointestinal mobility and secretory processing. The principal recognized actions of pancreatic glucagon, however, are to promote glycogenolysis and gluconeogenesis, resulting in an elevation of blood sugar levels. In this regard, the actions of glucagon are counterregulatory to those of insulin and may contribute to the hyperglycemia that accompanies Diabetes mellitus (Lund, P. K. et al., Proc. Natl. Acad. Sci., USA, 79: 345-349 (1982)).

Glucagon has been found to be capable of binding to specific receptors which lie on the surface of insulin producing cells. Glucagon, when bound to these receptors, stimulates the rapid synthesis of cAMP, by these cells. cAMP, in turn, has been found to stimulate insulin expression (Korman, L.Y. et al., Diabetes, 34:717-722 (1985)). Insulin acts to inhibit glucagon synthesis (Review of Medical Physiology, Ganong, W.F., 1979 Lange Publications, Los Altos, California (p. 273). Thus the expression of glucagon is carefully

-3-

regulated by insulin, and ultimately by the serum glucose level.

The glucagon gene is initially translated from a 630 base pair precursor to form the polypeptide, preproglucagon (Lund et al. (1982)). This polypeptide is subsequently processed to form proglucagon. Patzelt, C. et al., Nature, 282: 260-266 (1979), demonstrated that proglucagon was subsequently cleaved into glucagon and a second polypeptide. Subsequent work by Lund, P. K. et al., Lopez L. C. et al., and Bell, G. I. et al., (Nature) 302:716-718 (1983) demonstrated that the proglucagon molecule was cleaved immediately after lysine-arginine dipeptide residues. Studies of proglucagon produced by channel catfish (Ictalurus punctata) indicated that glucagon from this animal was also proteolytically cleaved after adjacent lysine-arginine and arginine-arginine dipeptide residues (Andrews, P. C. et al., J. Biol. Chem., 260: 3910-3914 (1985)). Lopez, L. C. et al., (Proc. Natl. Acad. Sci. USA 80:5485-5489 (1983)), and Bell, G. I. et al., discovered the mammalian proglucagon was cleaved at lysine-arginine or arginine-arginine dipeptides, and demonstrated that the proglucagon molecule contained three discrete and highly homologous peptide molecules which were designated glucagon, glucagon-like protein 1 (GLP-1) and glucagon-like protein 2 (GLP-2). Lopez et al. concluded that glucagon-like protein 1 was 37 amino acid residues long and that glucagon-like peptide 2 was 34 amino acid residues long. Analogous studies on the structure of rat preproglucagon revealed a similar pattern of proteolytic cleavage between adjacent lysine-arginine or arginine-arginine dipep-

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.