

REVIEW

Open Access



Endophthalmitis following intravitreal anti-vascular endothelial growth factor (VEGF) injection: a comprehensive review

Rohan Merani^{1,2,3,4*} and Alex P Hunyor^{1,2,3,5}

Abstract

The purpose of this review is to report and summarize previously reported studies and assess many of the individual steps of the intravitreal injection procedure's possible effect on the prevention of endophthalmitis. The pooled endophthalmitis rate from 20 large retrospective case series of anti-VEGF injections was 144/510,396 (0.028%; 1/3,544). Injections may be performed in an office-based location or in an operating room (OR) and low rates of endophthalmitis can be achieved in either location with careful attention to asepsis. Pre- or post-injection topical antibiotics have not been shown to be effective, and could select for more virulent microorganisms. Povidone-iodine prior to injection is accepted as the gold-standard antiseptic agent, but *aqueous* chlorhexidine may be an alternative. Antisepsis before and after gel or subconjunctival anesthetic is suggested. The preponderance of Streptococcal infections after intravitreal injection is discussed, including the possible role of aerosolization, which can be minimized by using face masks or maintaining silence. As with other invasive procedures in medicine, the use of sterile gloves, following adequate hand antisepsis, may be considered. Control of the eyelashes and lid margin is required to avoid contamination of the needle, but this can be achieved with or without a speculum. Techniques to minimize vitreous reflux have not been shown to reduce the risk of endophthalmitis. Same day bilateral injections should be performed as two separate procedures, preferably using drug from different lots, especially when using compounded drugs.

Keywords: Endophthalmitis, Intravitreal injection, Anti-VEGF, *Streptococcus*, Masks, Antisepsis, Povidone-Iodine, Chlorhexidine, Antibiotics, Speculum

Introduction

Intravitreal injection (IVI) is the most commonly performed ophthalmic procedure. In the USA, the number of injections performed has increased exponentially, from 4,215 injections in 2001 to 82,994 in 2004, to 812,413 in 2007, to 1.27 million in 2009 and to 2.5 million injections in 2011 [1, 2]. Similar increases have been observed in Canada and the United Kingdom [3, 4].

Infectious endophthalmitis (IE) secondary to IVI is a potentially devastating complication. It can be difficult to distinguish infectious endophthalmitis from "sterile" or non-infectious endophthalmitis. For the purpose of this review, IE refers to endophthalmitis that is clinically

suspected to be infectious, and treated as such with a vitreous tap and injection of antibiotics and/or vitrectomy surgery.

Bacteria are most likely inoculated into the vitreous cavity at the time of injection, or much less likely gain access later through the needle tract [5, 6]. The potential sources of bacteria include the patient's ocular or periorcular surfaces, aerosolized bacteria, or contamination of the needle, instruments, drug or drug vial [7].

Two meta-analyses including both retrospective series and clinical trials have calculated the pooled rate of endophthalmitis after anti-VEGF injections. McCannel found a rate of 52/105,536 injections (0.049%; 1 in 2030) [8] and more recently, Fileta et al. [9] calculated a rate of 197/350,535 (0.056%; 1 in 1,779). As patients typically receive ongoing intravitreal therapy, the per-patient risk

*Correspondence: rmerani@med.usyd.edu.au

¹ Retina Associates, Level 4, 8 Thomas St, Chatswood, NSW 2067, Australia
Full list of author information is available at the end of the article



© 2015 Merani and Hunyor. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/licenses/by/4.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

of endophthalmitis is significantly higher than the per-injection risk.

The rate of needle contamination after IVI has been reported as between 0.36 and 18%, which is significantly higher than the incidence of endophthalmitis after IVI [5, 7, 10]. The threshold inoculum size required to develop endophthalmitis is related to the type of bacteria and their virulence, intraocular immune mechanisms and anatomical characteristics of the vitreous [11, 12]. Animal studies have shown that a smaller number of bacterial colony-forming units are required to induce endophthalmitis when injected into the vitreous compared to when they are injected into the anterior chamber [13]. Endophthalmitis following intravitreal injection often presents earlier than after cataract surgery [14, 15].

The purpose of this review is to estimate the rate of endophthalmitis after intravitreal injection and to examine each step of the injection procedure that may influence the risk of endophthalmitis. To be able to prove that a particular measure reduces the risk of endophthalmitis would need huge numbers of patients in a randomized controlled trial, given that endophthalmitis is a relatively rare outcome. There is thus no Level 1 evidence for any preventative measure to reduce the incidence of endophthalmitis after intravitreal injection. As a result, this review largely summarizes retrospective papers, with their inherent biases.

Methods

A systematic literature search of the Medline database from 1996 to December 2014 was performed through Ovid, using search terms relevant to each section. Further literature was sourced from the reference lists of retrieved publications.

To estimate the per-injection rate of endophthalmitis after anti-VEGF injection, retrospective case series with at least 10,000 such injections were included. Studies that did not report a breakdown of the drugs used were excluded to avoid including triamcinolone and other injections in this calculation. Questionnaire-based and population-based studies were excluded given the incomplete data. Clinical trials were excluded as they may not reflect real-world practice, with more stringent requirements regarding injection technique often included in the protocols.

Results

Twenty retrospective case series meeting the inclusion criteria were identified. Details of the injection procedure and associated aseptic measures used in each series are

summarized in the Table 1. Where data were missing, the corresponding author for each study was contacted by email. Only two authors were not contactable.

We identified 144 cases of endophthalmitis from 510,396 anti-VEGF injections which equates to a pooled endophthalmitis rate of 0.028% or 1 in 3,544 injections [16–33].

Review

Location—office vs operating room (OR)

In the 2013 American Society of Retinal Specialists (ASRS) Preferences and Trends (PAT) Survey, over 98% of USA-based specialists reported performing injections in an office setting, compared with only 47% of international specialists [34]. In Germany and other parts of Europe, more injections are performed in the operating room (OR) [35, 36].

It has been [29] suggested that an advantage of the OR location is the superior air circulation systems. However, the ESCRS endophthalmitis study group was not able to find a relationship between the number of air changes per hour and the incidence of endophthalmitis after cataract surgery when they compared locations with minimal airflow, 20 air changes per hour and ultraclean air systems using laminar flow principles [37, 38].

Pooling the results of three OR-based injection series, the endophthalmitis rate was just 6/78,506 (0.0076% or 1 in 13,084) [19, 23, 25]. Common to these studies was the careful attention to asepsis with the use of sterile gloves, face masks, and drapes which were not used in most other office-based series (see Table 1). A notable exception is Shimada et al's series with no cases of endophthalmitis out of 15,144 injections where similar strict aseptic measures were followed in an office setting [27].

Abell et al. [29] reported an endophthalmitis rate of 4/3,376 (0.12%) for office-based injections compared with 0/8,873 (0%) for OR-based injections. In this non-randomized series, patients with private health insurance were treated in the OR while those without insurance were treated in the office. The difference in endophthalmitis rates may be a reflection of socioeconomic or other factors [39]. Tabendeh et al. [30] reported an endophthalmitis rate of 3/8,210 (0.037%) anti-VEGF injections in the office compared with 2/3,047 in the operating room (0.066%), in another non-randomised study that was not powered to be able to detect a difference. Compared with office-based injections, there was no apparent benefit to an OR environment in this small study.

Although there is no doubt that the OR has many advantages, there are logistical hurdles that make access to OR facilities difficult for many patients, and the OR location

Table 1 Endophthalmitis following intravitreal anti-VEGF injection—retrospective cases series with at least 10,000 injection

| Period and location | Authors | n = injections | Rate of clinically suspected IE | Pre-injection antibiotics | Post injection antibiotics | Mask | Drape | Conjunctival povidone-iodine concentration | Anaesthetic agents used | Sterile lid speculum | Gloves | Location |
|--|--|----------------|--|---------------------------|---|------|----------------|--|---------------------------|---|--------------------|----------|
| 1 Jan 2009 to 1 Oct 2012 Single-centre, USA | Storey et al. [16] (PIE study group) Ophthal 2013 | 117,171 | Overall 44/117,171 (0.038%); 1/2,663) By Agent Rani: 24/71,791 (0.033%; 1/2,991) Bev: 20/44,007 (0.045%; 1/2,200) Affib: 0/1,373 | Variable | Variable | No* | No | 5% | Drops Subconj (rarely) | Variable | Nil* | Office |
| 1 Jan 2005 to 31 Dec 2010 Single-centre (multi-site), USA | Moshfeghi et al. [17] Retina 2011 | 60,322 | Overall 12/60,322 (0.020%); 1/5,027) By Agent Rani: 5/18,607 (0.027%; 1/3,721) Bev: 7/39,700 (0.018%; 1/5,671) Peg: 0/2,015 | No | Variable | No* | No | 5% | Drops | Yes | Non-sterile | Office |
| 1 Jan 2007 to 31 Dec 2011 Single-centre, USA | Chaudhary et al. [18] Retina 2013 | 49,002 | Overall 17/49,002 (0.035%); 1 in 2,882) By Agent Rani: 2/20,297 (0.0099%; 1/10,149) Bev: 15/28,705 (0.052%; 1/1,914) | Yes | Yes | No | No | 5% | Drops Gel Subconj | Yes | Nil or non-sterile | Office |
| 2004 to 2012 Multicentre, Switzerland | Casparis et al. [19] Retina 2014 | 40,011 | Overall 3/40,011 (0.0075%); 1 in 13,337) By Agent Rani: 3/36,398 (0.0082%; 1/12,133) Bev: 0/3,518 Affib: 0/89 Peg: 0/6 | No | Variable (yes in one hospital, no in the other) | Yes | Yes (adhesive) | 5–10% | Drops | Yes | Sterile | OR |
| 1 Aug, 2006 to 31 Jul 2007 Multi-centre, USA | Klein et al. [20] Ophthal 2009 | 30,736 | Overall 15/30,736 (0.049%); 1/2,049) By Agent Rani: 10/22,579 (0.044%; 1/2,258) Bev: 5/8,039 (0.062%; 1/1,608) Peg: 0/128 | Variable | ? | ? | ? | 5–10% | ? | Variable: used in 14 of the 15 cases with endophthalmitis | ? | Office |

Table 1 continued

| Period and location | Authors | n = injections | Rate of clinically suspected IE | Pre-injection antibiotics | Post injection antibiotics | Mask | Drape | Conjunctival povidone-iodine concentration | Anaesthetic agents used | Sterile lid speculum | Gloves | Location |
|---|---|----------------|---|---------------------------|---|------|---------------------|--|-------------------------|--|--------------|----------|
| July 2000 to July 2010 Single-centre (multi-site), USA | Chen et al. [21] Retina 2011 | 29,995 | Overall 11/29,995 (0.037%); 1/2,727) By Agent Rani: 8/22,336 (0.036%); 1/2,792) Bev: 3/6,675 (0.045%); 1/2,225) Peg: 0/984 | Yes | Yes | No* | No* | 5–10% | Drops Gel Subconj | Yes | Non-sterile* | Office |
| 1 Jun 2005 to 7 Aug 2007 Multicentre, USA | Fintak et al. [22] Retina 2008 | 26,905 | Overall 6/26,905 (0.022%); 1/4,484) By Agent Rani: 3/14,320 (0.021%); 1/4,773) Bev: 3/12,585 (0.024%); 1/4,195) | No* | Variable (used in all the cases of endophthalmitis) | No* | No* | 5–10% | Drops Gel Subconj | Variable (used in all the cases of endophthalmitis) Fingers to spread the eyelids in a minority | Non-sterile* | Office |
| March 2007 to May 2013 Single-centre, Denmark | Brynskov et al. [23] Retina 2014 | 20,293 | Overall 0/20,293 By Agent Rani: 0/20,024 Aflib: 0/269 | No | Variable | Yes | Yes (adhesive) | 5% | Drops | Yes | Sterile | OR |
| 1 Aug 1997 to 31 Oct 2012 Single-centre, USA | Bhavsar and Sandler [24] Retina 2015 | 17,666 | Overall 1/17,666 (0.0057%); 1/17,666) By Agent Rani: 0/1,669 Bev: 1/15,479 (0.0065%); 1/15,479) Aflib: 0/148 Peg: 0/370 | No | No | No | No | 5% (before and after injection) | Drops | Yes | Non-sterile | Office |
| Jan 2005 to end July 2012 Single-centre, Germany | Nentwich et al. [25] Retina 2014 | 18,202 | Overall 3/18,202 (0.016%); 1/6,067) By Agent Rani: 1/10,097 (0.010%); 1/10,097) Bev: 2/7,865 (0.025%); 1/3,932) Peg: 0/240 | No | Yes | Yes | Yes (non-adhesive)* | 1% | Drops | Yes | Sterile | OR |

Table 1 continued

| Period and location | Authors | n = injections | Rate of clinically suspected IE | Pre-injection antibiotics | Post injection antibiotics | Mask | Drape | Conjunctival povidone-iodine concentration | Anaesthetic agents used | Sterile lid speculum | Gloves | Location |
|--|---------------------------------------|----------------|--|---------------------------|-----------------------------|---------------------------|---------------------------|--|-------------------------|----------------------|--|----------------------------------|
| Jan 2007 to May 2012 Multicentre, India | Mithal et al. [26] BJO 2013 | 15,925 | Overall 8/15,925 (0.050%;1/1,991) By Agent Rani: 1/705 (0.14%; 1/705) Bev: 7/15,035 (0.044%;1/2,275) Peg: 0/185 | No* | Yes* | Yes* | Variable | 2.5%* | Drops* | Yes* | Sterile* | Office* |
| July 2009 to July 2012 Single-centre, Japan | Shimada et al. [27] Graefes 2013 | 15,144 | Overall 0/15,144 By Agent Rani: 0/13,750 Bev: 0/846 Peg: 0/548 | Yes | Yes | Yes | Yes (adhesive) | 0.25% (before and after injection) | Drops + Subconj | Yes | Yes | Office |
| Jan 2005 to Aug 2010 Multi-centre, Canada | Cheung et al. [28] Ophthalmol 2012 | 14,960 | Overall 7/14,960 (0.047%); 1/2,137 By Agent Rani: 3/9,453 (0.032%); 1/3,151 Bev: 4/5,386 (0.074%); 1/1,347. Peg: 0/121 | Variable | Variable | No* | No | 10% | Drops Gel | Yes | Nil* | Office |
| Mar 2006 to Mar 2012 Multi-centre (single-surgeon), Australia | Abell et al. [29] BJO 2012 | 12,249 | Overall 4/12,249 (0.033%); 1/3,062 By Agent Rani: 3/10,574 (0.028%; 1/3,525)* Bev: 1/1,675 (0.060%); 1/1,675* | No | Variable (used until 2011)* | Yes | Yes (non-adhesive)* | 10% | Drops + Gel* | Yes | Sterile | Office = 4/3,376 OR = 0/8,873 |
| Jan 2009 to Dec 2011 Multi-centre, USA and Italy | Tabandeh et al. [30] Retina 2014 | 11,257 | Overall 5/11,257 By Agent Rani: 3/7,724 (0.11%); 1/908 Bev: 2/8,533 (0.023%); 1/4,267 | Yes in OR No in office | Yes | Yes in OR No in office | Yes in OR No in office | 5% | Drops Subconj | Yes | Sterile in OR Non-sterile in office | Office = 3/8,647 OR = 2/3,063 |

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.