

Vol 110, No 17, October 26, 2004

ISSN 0009-7322

http://circ.ahajournals.org

SUPPLEMENT TO

American Heart
Association®



Learn and LiveSM

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION

Abstracts
from

scientific
sessions
2004

Sessions: November 7–10
Exhibits: November 7–9
New Orleans, Louisiana
scientificsessions.org



basic science



clinical science



population science

Named and Invited Lectures 2004

2004 Russell Ross Memorial Lectureship in Vascular Biology • 2004 George Lyman Duff Memorial Lecture • 2004 Sol Sherry Distinguished Lecture in Thrombosis • 2004 Thomas W. Smith Memorial Lecture • 2004 George E. Brown Memorial Lecture • 2004 Dickinson W. Richards Memorial Lecture • 2004 Katharine A. Lembricht Award • 2004 William W. L. Glenn Lecture • 2004 William J. Rashkind Memorial Lecture • 2004 T. Duckett Jones Memorial Lecture • 2004 Charles T. Dotter Memorial Lecture • 2004 Laennec Society Lecture • 2004 Ancel Keys Lecture • 2004 Lewis K. Dahl Memorial Lecture • 2004 Robert I. Levy Endowed Lecture in Lipid Metabolism

New and Young Investigator Award/Prize Abstract Finalists

Lewis N. and Arnold M. Katz Basic Science Research Prize for Young Investigators • Melvin L. Marcus Young Investigator Awards in Cardiovascular Science • Cournand and Comroe Young Investigators Prizes in Cardiopulmonary and Critical Care • Outstanding Research Award in Pediatric Cardiology • Melvin Judkins Young Investigator Award in Cardiovascular Radiology • Vivian Thomas Young Investigator Award • Martha N. Hill New Investigator Awards • Elizabeth Barrett-Connor Research Award in Epidemiology and Prevention for Investigators in Training • Samuel A. Levine Young Clinical Investigator Awards • Laennec Society Young Clinician Award • NPAM New Investigator Award

Abstracts From the Scientific Sessions 2004

Diovan® (valsartan) Tablets

Rx only

BRIEF SUMMARY: Please see package insert for full prescribing information.

USE IN PREGNANCY: When used in pregnancy during the second and third trimesters, drugs that act directly on the renin-angiotensin system can cause injury and even death to the developing fetus. When pregnancy is detected, Diovan should be discontinued as soon as possible.
See **WARNINGS: Fetal/Neonatal Morbidity and Mortality.**

INDICATIONS AND USAGE: Hypertension: Diovan® (valsartan) is indicated for the treatment of hypertension. It may be used alone or in combination with other antihypertensive agents.

Heart Failure: Diovan is indicated for the treatment of heart failure (NYHA class II-IV) in patients who are intolerant of angiotensin converting enzyme inhibitors. In a controlled clinical trial, Diovan significantly reduced hospitalizations for heart failure. There is no evidence that Diovan provides added benefits when it is used with an adequate dose of an ACE inhibitor. (See **CLINICAL PHARMACOLOGY, Pharmacodynamics and Clinical Effects, Heart Failure in the full prescribing information for details.**)

CONTRAINDICATIONS: Diovan® (valsartan) is contraindicated in patients who are hypersensitive to any component of this product.

WARNINGS: Fetal/Neonatal Morbidity and Mortality: Drugs that act directly on the renin-angiotensin system can cause fetal and neonatal morbidity and death when administered to pregnant women. Several dozen cases have been reported in the world literature in patients who were taking angiotensin-converting enzyme inhibitors. When pregnancy is detected, Diovan® (valsartan) should be discontinued as soon as possible.

The use of drugs that act directly on the renin-angiotensin system during the second and third trimesters of pregnancy has been associated with fetal and neonatal injury, including hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnios has also been reported, presumably resulting from decreased fetal renal function; oligohydramnios in this setting has been associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development. Prematurity, intrauterine growth retardation, and patent ductus arteriosus have also been reported, although it is not clear whether these occurrences were due to exposure to the drug.

These adverse effects do not appear to have resulted from intrauterine drug exposure that has been limited to the first trimester. Mothers whose embryos and fetuses are exposed to an angiotensin II receptor antagonist only during the first trimester should be so informed. Nonetheless, when patients become pregnant, physicians should advise the patient to discontinue the use of valsartan as soon as possible.

Rarely (probably less often than once in every thousand pregnancies), no alternative to a drug acting on the renin-angiotensin system will be found. In these rare cases, the mothers should be apprised of the potential hazards to their fetuses, and serial ultrasound examinations should be performed to assess the intra-amniotic environment.

If oligohydramnios is observed, valsartan should be discontinued unless it is considered life-saving for the mother. Contraction stress testing (CST), a nonstress test (NST), or biophysical profiling (BPP) may be appropriate, depending upon the week of pregnancy. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury.

Infants with histories of *in utero* exposure to an angiotensin II receptor antagonist should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dialysis may be required as means of reversing hypotension and/or substituting for disordered renal function.

No teratogenic effects were observed when valsartan was administered to pregnant mice and rats at oral doses up to 600 mg/kg/day and to pregnant rabbits at oral doses up to 10 mg/kg/day. However, significant decreases in fetal weight, pup birth weight, pup survival rate, and slight delays in developmental milestones were observed in studies in which parental rats were treated with valsartan at oral, maternally toxic (reduction in body weight gain and food consumption) doses of 600 mg/kg/day during organogenesis or late gestation and lactation. In rabbits, fetotoxicity (i.e., resorptions, litter loss, abortions, and low body weight) associated with maternal toxicity (mortality) was observed at doses of 5 and 10 mg/kg/day. The no observed adverse effect doses of 600, 200 and 2 mg/kg/day in mice, rats and rabbits represent 9, 6, and 0.1 times, respectively, the maximum recommended human dose on a mg/m² basis. (Calculations assume an oral dose of 320 mg/day and a 60-kg patient.)

Hypertension: Excessive hypertension was rarely seen (0.1%) in patients with uncomplicated hypertension treated with Diovan alone. In patients with an activated renin-angiotensin system, such as volume- and/or salt-depleted patients receiving high doses of diuretics, symptomatic hypotension may occur. This condition should be corrected prior to administration of Diovan, or the treatment should start under close medical supervision.

If excessive hypotension occurs, the patient should be placed in the supine position and, if necessary, given an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further treatment, which usually can be continued without difficulty once the blood pressure has stabilized.

Hypertension in Heart Failure Patients: Caution should be observed when initiating therapy in patients with heart failure. Patients with heart failure given Diovan commonly have some reduction in blood pressure, but discontinuation of therapy because of continuing symptomatic hypotension usually is not necessary when dosing instructions are followed. In controlled trials, the incidence of hypotension in valsartan-treated patients was 5.5% compared to 1.8% in placebo-treated patients.

PRECAUTIONS: General: Impaired Hepatic Function: As the majority of valsartan is eliminated in the bile, patients with mild-to-moderate hepatic impairment, including patients with biliary obstructive disorders, showed lower valsartan clearance (higher AUCs). Care should be exercised in administering Diovan® (valsartan) to these patients.

Impaired Renal Function - Hypertension: In studies of ACE inhibitors in hypertensive patients with unilateral or bilateral renal artery stenosis, increases in serum creatinine or blood urea nitrogen have been reported. In a 4-day trial of valsartan in 12 hypertensive patients with unilateral renal artery stenosis, no significant increases in serum creatinine or blood urea nitrogen were observed. There has been no long-term use of Diovan in patients with unilateral or bilateral renal artery stenosis, but an effect similar to that seen with ACE inhibitors should be anticipated.

Impaired Renal Function - Heart Failure: As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients with severe heart failure whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, treatment with angiotensin-converting enzyme inhibitors and angiotensin receptor antagonists has been associated with oliguria and/or progressive azotemia and (rarely) with acute renal failure and/or death. Similar outcomes have been reported with Diovan.

Some patients with heart failure have developed increases in blood urea nitrogen, serum creatinine, and potassium. These effects are usually minor and transient, and they are more likely to occur in patients with pre-existing renal impairment. Dosage reduction and/or discontinuation of the diuretic and/or Diovan may be required. In the Valsartan Heart Failure Trial, in which 93% of patients were on concomitant ACE inhibitors, treatment was discontinued for elevations in creatinine or potassium (total of 1.0% on valsartan vs. 0.2% on placebo). Evaluation of patients with heart failure should always include assessment of renal function.

Concomitant Therapy in Patients with Heart Failure: In patients with heart failure, concomitant use of Diovan, an ACE inhibitor, and a beta blocker is not recommended. In the Valsartan Heart Failure Trial, this triple combination was associated with an increased risk of heart failure outcome (see **CLINICAL PHARMACOLOGY, Pharmacodynamics and Clinical Effects, Heart Failure in the full prescribing information**).

Information for Patients: Pregnancy: Female patients of childbearing age should be told about the consequences of second- and third-trimester exposure to drugs that act on the renin-angiotensin system, and they should also be told that these consequences do not appear to have resulted from intrauterine drug exposure that has been limited to the first trimester. These patients should be asked to report pregnancies to their physicians as soon as possible.

Drug Interactions: No clinically significant pharmacokinetic interactions were observed when valsartan was coadministered with amlodipine, atenolol, cimetidine, digoxin, furosemide, glyburide, hydrochlorothiazide, or indomethacin. The valsartan-atenolol combination was more antihypertensive than either component, but it did not lower the heart rate more than atenolol alone.

Coadministration of valsartan and warfarin did not change the pharmacokinetics of valsartan or the time-course of the anticoagulant properties of warfarin.

CYP 450 Interactions: The enzyme(s) responsible for valsartan metabolism have not been identified but do not seem to be CYP 450 isozymes. The inhibitory or induction potential of valsartan on CYP 450 is also unknown.

As with other drugs that block angiotensin II or its effects, concomitant use of potassium sparing diuretics (e.g., spironolactone, triamterene, amiloride), potassium supplements, or salt substitutes containing potassium may lead to increases in serum potassium and in heart failure patients to increases in serum creatinine.

Carcinogenesis, Mutagenesis, Impairment of Fertility: There was no evidence of carcinogenicity when valsartan was administered in the diet to mice and rats for up to 2 years at doses up to 160 and 200 mg/kg/day, respectively. These doses in mice and rats are about 2.6 and 6 times, respectively, the maximum recommended human dose on a mg/m² basis. (Calculations assume an oral dose of 320 mg/day and a 60-kg patient.)

Mutagenicity assays did not reveal any valsartan-related effects at either the gene or chromosome level. These assays included bacterial mutagenicity tests with *Salmonella* (Ames) and *E. coli*: a gene mutation test with Chinese hamster V79 cells; a cytogenetic test with Chinese hamster ovary cells; and a rat micronucleus test.

Valsartan had no adverse effects on the reproductive performance of male or female rats at oral doses up to 200 mg/kg/day. This dose is 6 times the maximum recommended human dose on a mg/m² basis. (Calculations assume an oral dose of 320 mg/day and a 60-kg patient.)

Pregnancy: Pregnancy Categories C (first trimester) and D (second and third trimesters): See **WARNINGS, Fetal/Neonatal Morbidity and Mortality.**

Nursing Mothers: It is not known whether valsartan is excreted in human milk, but valsartan was excreted in the milk of lactating rats. Because of the potential for adverse effects on the nursing infant, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use: Safety and effectiveness in pediatric patients have not been established.

Geriatric Use: In the controlled clinical trials of valsartan, 1,214 (36.2%) of hypertensive patients treated with valsartan were ≥ 65 years and 285 (7.9%) were ≥ 75 years. No overall difference in the efficacy or safety of valsartan was observed in this patient population, but greater sensitivity of some older individuals cannot be ruled out. Of the 2,511 patients with heart failure randomized to valsartan in the Valsartan Heart Failure Trial, 45% (1,141) were 65 years of age or older. There were no notable differences in efficacy or safety between older and younger patients.

ADVERSE REACTIONS: Hypertension: Diovan® (valsartan) has been evaluated for safety in more than 4,000 patients, including over 400 treated for over 6 months, and more than 160 for over 1 year. Adverse experiences have generally been mild and transient in nature and have only infrequently required discontinuation of therapy. The overall incidence of adverse experiences with Diovan was similar to placebo.

The overall frequency of adverse experiences was neither dose-related nor related to gender, age, race, or regimen. Discontinuation of therapy due to side effects was required in 2.3% of valsartan patients and 2.0% of placebo patients. The most common reasons for discontinuation of therapy with Diovan were headache and dizziness.

The adverse experiences that occurred in placebo-controlled clinical trials in at least 1% of patients treated with Diovan and at a higher incidence in valsartan (n=2,316) than placebo (n=888) patients included viral infection (3% vs. 2%), fatigue (2% vs. 1%), and abdominal pain (2% vs. 1%).

Headache, dizziness, upper respiratory infection, cough, diarrhea, rhinitis, sinusitis, nausea, pharyngitis, edema, and arthralgia occurred at a more than 1% rate but at about the same incidence in placebo and valsartan patients.

In trials in which valsartan was compared to an ACE inhibitor with or without placebo, the incidence of dry cough was significantly greater in the ACE-inhibitor group (7.9% than in the groups who received valsartan (2.6%) or placebo (1.5%). In a 129-patient trial limited to patients who had had dry cough when they had previously received ACE inhibitors, the incidences of cough in patients who received valsartan, HCTZ, or lisinopril were 20%, 19%, and 69% respectively (p < 0.001).

Dose-related orthostatic effects were seen in less than 1% of patients. An increase in the incidence of dizziness was observed in patients treated with Diovan 320 mg (8%) compared to 10 to 160 mg (2% to 4%).

Diovan has been used concomitantly with hydrochlorothiazide without evidence of clinically important adverse interactions.

Other adverse experiences that occurred in controlled clinical trials of patients treated with Diovan (> 0.2% of valsartan patients) are listed below. It cannot be determined whether these events were causally related to Diovan.

Body as a Whole: Allergic reaction and asthenia; **Cardiovascular:** Palpitations; **Dermatologic:** Pruritus and rash; **Digestive:** Constipation, dry mouth, dyspepsia, and flatulence; **Musculoskeletal:** Back pain, muscle cramps, and myalgia; **Neurologic and Psychiatric:** Anxiety, insomnia, paresthesia, and somnolence; **Respiratory:** Dyspnea; **Special Senses:** Vertigo; **Urogenital:** Impotence

Other reported events seen less frequently in clinical trials included chest pain, syncope, anorexia, vomiting, and angioedema.

Heart Failure: The adverse experience profile of Diovan in heart failure patients was consistent with the pharmacology of the drug and the health status of the patients. In the Valsartan Heart Failure Trial, comparing valsartan in total daily doses up to 320 mg (n=2,506) to placebo (n=2,494), 10% of valsartan patients discontinued for adverse events vs. 7% of placebo patients.

The table shows adverse events in double-blind short-term heart failure trials, including the first 4 months of the Valsartan Heart Failure Trial, with an incidence of at least 2% that were more frequent in valsartan-treated patients than in placebo-treated patients. All patients received standard drug therapy for heart failure, frequently as multiple medications, which could include diuretics, digitalis, beta-blockers, or ACE inhibitors.

	Valsartan (n=3,282)		Placebo (n=2,740)	
Dizziness	17%	9%	3%	2%
Hypotension	7%	2%	2%	1%
Diarrhea	5%	4%	2%	1%
Arthralgia	3%	2%	2%	1%
Fatigue	3%	2%	2%	1%

Other adverse events with an incidence greater than 1% and greater than placebo included headache NOS, nausea, renal impairment NOS, syncope, blurred vision, upper abdominal pain and vertigo. (NOS = not otherwise specified).

From the long term data in the Valsartan Heart Failure Trial, there did not appear to be any significant adverse events not previously identified.

Post-Marketing Experience: The following additional adverse reactions have been reported in post-marketing experience:

Hypersensitivity: There are rare reports of angioedema. **Digestive:** Elevated liver enzymes and very rare reports of hepatitis. **Renal:** Impaired renal function. **Clinical Laboratory Tests:** Hyperkalemia; **Dermatologic:** Alopecia.

Clinical Laboratory Test Findings: In controlled clinical trials, clinically important changes in standard laboratory parameters were rarely associated with administration of Diovan.

Creatinine: Minor elevations in creatinine occurred in 0.8% of patients taking Diovan and 0.6% given placebo in controlled clinical trials of hypertensive patients. In heart failure trials, greater than 50% increases in creatinine were observed in 3.9% of Diovan-treated patients compared to 0.9% of placebo-treated patients.

Hemoglobin and Hematocrit: Greater than 20% decreases in hemoglobin and hematocrit were observed in 0.4% and 0.8%, respectively, of Diovan patients, compared with 0.1% and 0.1% in placebo-treated patients. One valsartan patient discontinued treatment for microcytic anemia.

Liver Function Tests: Occasional elevations (greater than 150%) of liver chemistries occurred in Diovan-treated patients. Three patients (< 0.1%) treated with valsartan discontinued treatment for elevated liver chemistries.

Neutropenia: Neutropenia was observed in 1.9% of patients treated with Diovan and 0.8% of patients treated with placebo.

Serum Potassium: In hypertensive patients, greater than 20% increases in serum potassium were observed in 4.4% of Diovan-treated patients compared to 2.9% of placebo-treated patients. In heart failure patients, greater than 20% increases in serum potassium were observed in 10.0% of Diovan-treated patients compared to 5.1% of placebo-treated patients.

Blood Urea Nitrogen (BUN): In heart failure trials, greater than 50% increases in BUN were observed in 16.6% of Diovan-treated patients compared to 6.3% of placebo-treated patients.

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F)

[see USP Controlled Room Temperature].

Protect from moisture.

Dispense in tight container (USP).

REV: NOVEMBER 2003

PRINTED IN U.S.A.

T2003-71

89004209

Distributed by:

Novartis Pharmaceuticals Corporation, East Hanover, New Jersey 07936

©Novartis

Reference: 1. Hermida RC, Calvo C, Ayala DE, et al. Administration time-dependent effects of valsartan on ambulatory blood pressure in hypertensive subjects. *Hypertension*. 2003;42:283-290.



Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936

©2004 Novartis

Printed in U.S.A.

6/04

C-DI0-1101

Printed on Recycled Paper

DIOVAN DELIVERS

USE IN PREGNANCY: When used in pregnancy during the second and third trimesters, drugs that act directly on the renin-angiotensin system can cause injury and even death to the developing fetus. When pregnancy is detected, DIOVAN should be discontinued as soon as possible. See **WARNINGS** in brief summary of complete Prescribing Information on next page.

DIOVAN is contraindicated in patients who are hypersensitive to any component of this product.

In hypertension, AEs more frequent with DIOVAN than placebo: viral infection (3% vs 2%), fatigue (2% vs 1%), abdominal pain (2% vs 1%); the most common AEs: headache and dizziness. An increase in dizziness was observed with 320 mg (8%) vs 10 mg to 160 mg (2% to 4%).

Because of the risk of hypotension, caution should be observed when initiating therapy in HF patients. Evaluation of HF patients should always include assessment of renal function. In patients with Heart Failure, concomitant use of DIOVAN, an ACE inhibitor, and a beta-blocker is not recommended. In the Valsartan Heart Failure Trial, this triple combination was associated with an unfavorable Heart Failure outcome.

In Val-HeFT, discontinuation due to adverse events was 10% for valsartan vs 7% for placebo; most common side effects were: dizziness (17% vs 9%), hypotension (7% vs 2%), and diarrhea (5% vs 4%).

www.DIOVAN.com

**Proven highly effective
BP reductions¹**

**Proven cardiovascular benefits:
Exclusive ARB indication for
HF patients intolerant of ACEIs**

**Proven excellent tolerability
in hypertension**

Definitely...DIOVAN

Diovan[®]
valsartan tablets

52 31LML 303
XL
1/05 31151-200

More than 10 million prescriptions!

Ride
the CREST™

Greater LDL-C reductions

Greater reductions in LDL-C

- In a comparison of dose-response curves, CRESTOR 10 mg to 40 mg **reduced LDL-C significantly more than atorvastatin** 10 mg to 80 mg (4.1%; $P < .001$)¹

In patients without CHD/CHD risk equivalents


94% of patients reached LDL-C goal at a low 10-mg dose*²

CRESTOR is indicated as an adjunct to diet to reduce elevated total-C, LDL-C, ApoB, non-HDL-C, and TG levels and to increase HDL-C in patients with primary hypercholesterolemia (heterozygous familial and nonfamilial) and mixed dyslipidemia.

CRESTOR is contraindicated in patients with active liver disease or with unexplained persistent elevations of serum transaminases, in women who are or may become pregnant, and in nursing mothers.

Rare cases of rhabdomyolysis with acute renal failure secondary to myoglobinuria have been reported with CRESTOR and with other drugs in this class. Patients should be advised to promptly report unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever. Therapy with CRESTOR should be discontinued if markedly elevated CK levels occur or myopathy is diagnosed or suspected.

*NCEP ATP III goal for patients without CHD or CHD risk equivalents:
optimal, <100 mg/dL; ≥2 risk factors, <130 mg/dL; 0-1 risk factor, <160 mg/dL.



Approved in over 64 countries worldwide!

LANE MEDICAL LIBRARY
STANFORD UNIV MED CENTER

NOV 11 2004

STANFORD, CA 94305-5123

with proven safety

as an adjunct to diet for many patients with dyslipidemia

Proven safety similar to other leading statins³

- Adverse reactions were **mild and transient**; incidence and discontinuation rates similar to placebo and other statins^{3,4}
- The most frequent adverse events were myalgia (3.3%), constipation (1.4%), asthenia (1.3%), abdominal pain (1.3%), and nausea (1.3%)^{3,5}

CRESTOR was evaluated in over 10,000 patients in preapproval clinical trials—more than any other statin prior to approval^{4,6-11}

Overall, more than 42,000 patients have taken CRESTOR in clinical trials, including ongoing trials in the GALAXY™ Program¹²

- The GALAXY Program is a large, comprehensive, long-term evolving global research initiative evaluating the efficacy and safety of CRESTOR and demonstrating the AstraZeneca confidence in, and commitment to, this important therapeutic option

It is recommended that liver function tests be performed before and at 12 weeks following both the initiation of therapy and any elevation of dose, and periodically (eg, semiannually) thereafter.

The effect of CRESTOR on cardiovascular morbidity and mortality has not been determined; long-term outcome studies are currently under way.

Please see brief summary of full Prescribing Information on reverse side of this advertisement.



CRESTOR[®]
rosuvastatin calcium

BRIEF SUMMARY: For full Prescribing Information, see package insert.

INDICATIONS AND USAGE CRESTOR is indicated: 1. as an adjunct to diet to reduce elevated total-C, LDL-C, ApoB, nonHDL-C, and TG levels and to increase HDL-C in patients with primary hypercholesterolemia (heterozygous familial and nonfamilial) and mixed dyslipidemia (Fredrickson Type IIa and IIb); 2. as an adjunct to diet for the treatment of patients with elevated serum TG (Fredrickson Type IV); 3. to reduce LDL-C, total-C, and ApoB in patients with homozygous familial hypercholesterolemia as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable. **CONTRAINDICATIONS** CRESTOR is contraindicated in patients with a known hypersensitivity to any component of this product. Rosuvastatin is contraindicated in patients with active liver disease or with unexplained persistent elevations of serum transaminases (see WARNINGS, Liver Enzymes). **Pregnancy and Lactation** Atherosclerosis is a chronic process and discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hypercholesterolemia. Cholesterol and other products of cholesterol biosynthesis are essential components for fetal development (including synthesis of steroids and cell membranes). Since HMG-CoA reductase inhibitors decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol, they may cause fetal harm when administered to pregnant women. Therefore, HMG-CoA reductase inhibitors are contraindicated during pregnancy and in nursing mothers. ROSUVASTATIN SHOULD BE ADMINISTERED TO WOMEN OF CHILDBEARING AGE ONLY WHEN SUCH PATIENTS ARE HIGHLY UNLIKELY TO CONCEIVE AND HAVE BEEN INFORMED OF THE POTENTIAL HAZARDS. If the patient becomes pregnant while taking this drug, therapy should be discontinued immediately and the patient apprised of the potential hazard to the fetus. **WARNINGS** **Liver Enzymes** HMG-CoA reductase inhibitors, like some other lipid-lowering therapies, have been associated with biochemical abnormalities of liver function. The incidence of persistent elevations (>3 times the upper limit of normal [ULN]) occurring on 2 or more consecutive occasions) in serum transaminases in fixed dose studies was 0.4, 0.0, and 0.1% in patients who received rosuvastatin 5, 10, 20, and 40 mg, respectively. In most cases, the elevations were transient and resolved or improved on continued therapy or after a brief interruption in therapy. There were two cases of jaundice, for which a relationship to rosuvastatin therapy could not be determined, which resolved after discontinuation of therapy. There were no cases of liver failure or irreversible liver disease in these trials. It is recommended that liver function tests be performed before and at 12 weeks following both the initiation of therapy and any elevation of dose, and periodically (e.g., semiannually) thereafter. Liver enzyme changes generally occur in the first 3 months of treatment with rosuvastatin. Patients who develop increased transaminase levels should be monitored until the abnormalities have resolved. Should an increase in ALT or AST of >3 times ULN persist, reduction of dose or withdrawal of rosuvastatin is recommended. Rosuvastatin should be used with caution in patients who consume substantial quantities of alcohol and/or have a history of liver disease (see CLINICAL PHARMACOLOGY, Special Populations, Hepatic Insufficiency). Active liver disease or unexplained persistent transaminase elevations are contraindications to the use of rosuvastatin (see CONTRAINDICATIONS). **Myopathy/Rhabdomyolysis** Rare cases of rhabdomyolysis with acute renal failure secondary to myoglobinuria have been reported with rosuvastatin and with other drugs in this class. Uncomplicated myalgia has been reported in rosuvastatin-treated patients (see ADVERSE REACTIONS). Creatine kinase (CK) elevations (>10 times upper limit of normal) occurred in 0.2% to 0.4% of patients taking rosuvastatin at doses of up to 40 mg in clinical studies. Treatment-related myopathy, defined as muscle aches or muscle weakness in conjunction with increases in CK values >10 times upper limit of normal, was reported in up to 0.1% of patients taking rosuvastatin doses of up to 40 mg in clinical studies. Rare cases of rhabdomyolysis were seen with higher than recommended doses (80 mg) of rosuvastatin in clinical trials. Factors that may predispose patients to myopathy with HMG-CoA reductase inhibitors include advanced age (>65 years), hypothyroidism, and renal insufficiency. The incidence of myopathy increased at doses of rosuvastatin above the recommended dosage range. Consequently, 1. Rosuvastatin should be prescribed with caution in patients with predisposing factors for myopathy, such as, renal impairment (see DOSAGE AND ADMINISTRATION), advanced age, and hypothyroidism. 2. Patients should be advised to promptly report unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever. Rosuvastatin therapy should be discontinued if markedly elevated CK levels occur or myopathy is diagnosed or suspected. 3. The risk of myopathy during treatment with rosuvastatin may be increased with concurrent administration of other lipid-lowering therapies or cyclosporine, (see CLINICAL PHARMACOLOGY, Drug Interactions, PRECAUTIONS, Drug Interactions, and DOSAGE AND ADMINISTRATION). The benefit of further alterations in lipid levels by the combined use of rosuvastatin with fibrates or niacin should be carefully weighed against the potential risks of this combination. Combination therapy with rosuvastatin and gemfibrozil should generally be avoided. (See DOSAGE AND ADMINISTRATION and PRECAUTIONS, Drug Interactions). 4. The risk of myopathy during treatment with rosuvastatin may be increased in circumstances which increase rosuvastatin drug levels (see CLINICAL PHARMACOLOGY, Special Populations, Race and Renal Insufficiency, and PRECAUTIONS, General). 5. Rosuvastatin therapy should also be temporarily withheld in any patient with an acute, serious condition suggestive of myopathy or predisposing to the development of renal failure secondary to rhabdomyolysis (e.g., sepsis, hypotension, major surgery, trauma, severe metabolic, endocrine, and electrolyte disorders, or uncontrolled seizures). **PRECAUTIONS** **General** Before instituting therapy with rosuvastatin, an attempt should be made to control hypercholesterolemia with appropriate diet and exercise, weight reduction in obese patients, and treatment of underlying medical problems (see INDICATIONS AND USAGE). Administration of rosuvastatin 20 mg to patients with severe renal impairment ($CL_{CR} < 30 \text{ mL/min/1.73 m}^2$) resulted in a 3-fold increase in plasma concentrations of rosuvastatin compared with healthy volunteers (see WARNINGS, Myopathy/Rhabdomyolysis and DOSAGE AND ADMINISTRATION). Pharmacokinetic studies show an approximate 2-fold elevation in median exposure in Japanese subjects residing in Japan and in Chinese subjects residing in Singapore compared with Caucasians residing in North America and Europe. The contribution of environmental and genetic factors to the difference observed has not been determined. However, these increases should be considered when making rosuvastatin dosing decisions for patients of Japanese and Chinese ancestry. (See WARNINGS, Myopathy/Rhabdomyolysis; CLINICAL PHARMACOLOGY, Special Populations, Race.) **Information for Patients** Patients should be advised to report promptly unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever. When taking rosuvastatin with an aluminum and magnesium hydroxide combination antacid, the antacid should be taken at least 2 hours after rosuvastatin administration (see CLINICAL PHARMACOLOGY, Drug Interactions). **Laboratory Tests** In the rosuvastatin clinical trial program, dipstick-positive proteinuria and microscopic hematuria were observed among rosuvastatin-treated patients, predominantly in patients dosed above the recommended dose range (i.e., 80 mg). However, this finding was more frequent in patients taking rosuvastatin 40 mg, when compared to lower doses of rosuvastatin or comparator statins, though it was generally transient and was not associated with worsening renal function. Although the clinical significance of this finding is unknown, a dose reduction should be considered for patients on rosuvastatin 40 mg therapy with unexplained persistent proteinuria during routine urinalysis testing. **Drug Interactions** **Cyclosporine:** When rosuvastatin 10 mg was coadministered with cyclosporine in cardiac transplant patients, rosuvastatin mean C_{max} and mean AUC were increased 11-fold and 7-fold, respectively, compared with healthy volunteers. These increases are considered to be clinically significant and require special consideration in the dosing of rosuvastatin to patients taking concomitant cyclosporine (see WARNINGS, Myopathy/Rhabdomyolysis, and DOSAGE AND ADMINISTRATION). **Warfarin:** Coadministration of rosuvastatin to patients on stable

warfarin therapy resulted in clinically significant rises in INR (>4, baseline 2-3). In patients taking coumarin anticoagulants and rosuvastatin concomitantly, INR should be determined before starting rosuvastatin and frequently enough during early therapy to ensure that no significant alteration of INR occurs. Once a stable INR time has been documented, INR can be monitored at the intervals usually recommended for patients on coumarin anticoagulants. If the dose of rosuvastatin is changed, the same procedure should be repeated. Rosuvastatin therapy has not been associated with bleeding or with changes in INR in patients not taking anticoagulants. **Gemfibrozil:** Coadministration of a single rosuvastatin dose to healthy volunteers on gemfibrozil (600 mg twice daily) resulted in a 2.2- and 1.9-fold, respectively, increase in mean C_{max} and mean AUC of rosuvastatin (see DOSAGE AND ADMINISTRATION). **Endocrine Function** Although clinical studies have shown that rosuvastatin alone does not reduce basal plasma cortisol concentration or impair adrenal reserve, caution should be exercised if any HMG-CoA reductase inhibitor or other agent used to lower cholesterol levels is administered concomitantly with drugs that may decrease the levels or activity of endogenous steroid hormones such as ketoconazole, spiroinolactone, and cimetidine. **CNS Toxicity** CNS vascular lesions, characterized by perivascular hemorrhages, edema, and mononuclear cell infiltration of perivascular spaces, have been observed in dogs treated with several other members of this drug class. A chemically similar drug in this class produced dose-dependent optic nerve degeneration (Wallerian degeneration of retinogeniculate fibers) in dogs, at a dose that produced plasma drug levels about 30 times higher than the mean drug level in humans taking the highest recommended dose. Edema, hemorrhage, and partial necrosis in the interstitium of the choroid plexus was observed in a female dog sacrificed moribund at day 24 at 90 mg/kg/day by oral gavage (systemic exposures 100 times the human exposure at 40 mg/kg/day based on AUC comparisons). Corneal opacity was seen in dogs treated for 52 weeks at 6 mg/kg/day by oral gavage (systemic exposures 20 times the human exposure at 40 mg/kg/day based on AUC comparisons). Cataracts were seen in dogs treated for 12 weeks by oral gavage at 30 mg/kg/day (systemic exposures 60 times the human exposure at 40 mg/kg/day based on AUC comparisons). Retinal dysplasia and retinal loss were seen in dogs treated for 4 weeks by oral gavage at 90 mg/kg/day (systemic exposures 100 times the human exposure at 40 mg/kg/day based on AUC). Doses $\leq 30 \text{ mg/kg/day}$ (systemic exposures ≤ 60 times the

Experiences Adverse experiences, regardless of causality assessment, reported in $\geq 2\%$ of patients in placebo-controlled clinical studies of rosuvastatin are shown in Table 1; discontinuations due to adverse events in these studies of up to 12 weeks duration occurred in 3% of patients on rosuvastatin and 5% on placebo.

Table 1. Adverse Events in Placebo-Controlled Studies

Adverse event	Rosuvastatin N=744	Placebo N=382
Pharyngitis	9.0	7.6
Headache	5.5	5.0
Diarrhea	3.4	2.9
Dyspepsia	3.4	3.1
Nausea	3.4	3.1
Myalgia	2.8	1.3
Asthenia	2.7	2.6
Back pain	2.6	2.4
Flu syndrome	2.3	1.8
Urinary tract infection	2.3	1.6
Rhinitis	2.2	2.1
Sinusitis	2.0	1.8

In addition, the following adverse events were reported, regardless of causality assessment, in $\geq 1\%$ of 10,275 patients treated with rosuvastatin in clinical studies. The events in *italics* occurred in $\geq 2\%$ of these patients. **Body as a Whole:** Abdominal pain, accidental injury, chest pain, infection, pain, pelvic pain, and neck pain. **Cardiovascular System:** Hypertension, angina pectoris, vasodilatation, and palpitation. **Digestive System:** Constipation, gastroenteritis, vomiting, flatulence, periodontal abscess, and gastritis. **Endocrine:** Diabetes mellitus. **Hemic and Lymphatic System:** Anemia and ecchymosis. **Metabolic and Nutritional Disorders:** Peripheral edema. **Musculoskeletal System:** Arthritis, arthralgia, and pathological fracture. **Nervous System:** Dizziness, insomnia, hypertension, paraesthesia, depression, anxiety, vertigo, and neuralgia. **Respiratory System:** Bronchitis, cough increased, dyspnea, pneumonia, and asthma. **Skin and Appendages:** Rash and pruritus. **Laboratory Abnormalities:** In the rosuvastatin clinical trial program, dipstick-positive proteinuria and microscopic hematuria were observed among rosuvastatin-treated patients, predominantly in patients dosed above the recommended dose range (i.e., 80 mg). However, this finding was more frequent in patients taking rosuvastatin 40 mg, when compared to lower doses of rosuvastatin or comparator statins, though it was generally transient and was not associated with worsening renal function. (See PRECAUTIONS, Laboratory Tests.) Other abnormal laboratory values reported were elevated creatine phosphokinase, transaminases, hyperglycemia, glutamyl transpeptidase, alkaline phosphatase, bilirubin, and thyroid function abnormalities. Other adverse events reported less frequently than 1% in the rosuvastatin clinical study program, regardless of causality assessment, included arrhythmia, hepatitis, hypersensitivity reactions (i.e., face edema, thrombocytopenia, leukopenia, vesiculobullous rash, urticaria, and angioedema), kidney failure, syncope, myasthenia, myositis, pancreatitis, photosensitivity reaction, myopathy, and rhabdomyolysis. **OVERDOSAGE** There is no specific treatment in the event of overdose. In the event of overdose, the patient should be treated symptomatically and supportive measures instituted as required. Hemodialysis does not significantly enhance clearance of rosuvastatin. **DOSAGE AND ADMINISTRATION** The patient should be placed on a standard cholesterol-lowering diet before receiving CRESTOR and should continue on this diet during treatment. CRESTOR can be administered as a single dose at any time of day, with or without food. **Hypercholesterolemia (Heterozygous Familial and Nonfamilial) and Mixed Dyslipidemia (Fredrickson Type IIa and IIb)** The dose range for CRESTOR is 5 to 40 mg once daily. The therapy with CRESTOR should be individualized according to goal of therapy and response. The usual recommended starting dose of CRESTOR is 10 mg once daily. Initiation of therapy with 5 mg once daily may be considered for patients requiring less aggressive LDL-C reductions or who have predisposing factors for myopathy (see WARNINGS, Myopathy/Rhabdomyolysis). For patients with marked hypercholesterolemia (LDL-C > 190 mg/dL) and aggressive lipid targets, a 20-mg starting dose may be considered. The 40-mg dose of CRESTOR should be reserved for those patients who have not achieved goal LDL-C at 20 mg (see WARNINGS, Myopathy/Rhabdomyolysis). After initiation and/or upon titration of CRESTOR, lipid levels should be analyzed within 2 to 4 weeks and dosage adjusted accordingly. **Homozygous Familial Hypercholesterolemia** The recommended starting dose of CRESTOR is 20 mg once daily in patients with homozygous FH. The maximum recommended daily dose is 40 mg. CRESTOR should be used in these patients as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable. Response to therapy should be estimated from pre-apheresis LDL-C levels. **Dosage in Patients Taking Cyclosporine** In patients taking cyclosporine, therapy should be limited to CRESTOR 5 mg once daily (see WARNINGS, Myopathy/Rhabdomyolysis, and PRECAUTIONS, Drug Interactions). **Concomitant Lipid-Lowering Therapy** The effect of CRESTOR on LDL-C and total-C may be enhanced when used in combination with a bile acid binding resin. If CRESTOR is used in combination with gemfibrozil, the dose of CRESTOR should be limited to 10 mg once daily (see WARNINGS, Myopathy/Rhabdomyolysis, and PRECAUTIONS, Drug Interactions). **Dosage in Patients With Renal Insufficiency** No modification of dosage is necessary for patients with mild to moderate renal insufficiency. For patients with severe renal impairment ($CL_{CR} < 30 \text{ mL/min/1.73 m}^2$) not on hemodialysis, dosing of CRESTOR should be started at 5 mg once daily and not to exceed 10 mg once daily (see PRECAUTIONS, General, and CLINICAL PHARMACOLOGY, Special Populations, Renal Insufficiency).

Rx only

References: 1. Data on file, DA-CRS-02. 2. Shepherd J, Hunninghake DB, Barter P, et al. Guidelines for lowering lipids to reduce coronary artery disease risk: a comparison of rosuvastatin with atorvastatin, pravastatin, and simvastatin for achieving lipid-lowering goals. *Am J Cardiol* 2003;91(suppl):11C-19C. 3. Shepherd J, Hunninghake DB, Stein EA, et al. The safety of rosuvastatin. *Am J Cardiol*. In press. 4. Prescribing Information for CRESTOR. AstraZeneca, Wilmington, DE. 5. Data on file, DA-CRS-01. 6. New Drug Application for atorvastatin (four-month safety update). 7. New Drug Application for simvastatin (simvastatin safety update II). 8. New Drug Application for pravastatin (medical officer's review and evaluation of NDA report). 9. New Drug Application for cerivastatin (conclusion with regard to the safety of cerivastatin). 10. New Drug Application for fluvastatin (analysis). 11. New Drug Application for lovastatin (integrated safety results and exposure). 12. Data on file, DA-CRS-11.

CRESTOR is a registered trademark of the AstraZeneca group of companies. "Ride the CREST" and the GALAXY Program are trademarks of the AstraZeneca group of companies.

©2004 AstraZeneca Pharmaceuticals LP. All rights reserved.

POC 630100
09/04 229369



human exposure at 40 mg/kg/day based on AUC comparisons) following treatment up to one year, did not reveal retinal findings. **Carcinogenesis, Mutagenesis, Impairment of Fertility** In a 104-week carcinogenicity study in rats at dose levels of 2, 20, 60, or 80 mg/kg/day by oral gavage, the incidence of uterine stromal polyps was significantly increased in females at 80 mg/kg/day at systemic exposure 20 times the human exposure at 40 mg/kg/day based on AUC. Increased incidence of polyps was not seen at lower doses. In a 107-week carcinogenicity study in mice given 10, 60, 200 mg/kg/day by oral gavage, an increased incidence of hepatocellular adenoma/carcinoma was observed at 200 mg/kg/day at systemic exposures 20 times human exposure at 40 mg/kg/day based on AUC. An increased incidence of hepatocellular tumors was not seen at lower doses. Rosuvastatin was not mutagenic or clastogenic with or without metabolic activation in the Ames test with *Salmonella typhimurium* and *Escherichia coli*, the mouse lymphoma assay, and the chromosomal aberration assay in Chinese hamster lung cells. Rosuvastatin was negative in the *in vivo* mouse micronucleus test. In rat fertility studies with oral gavage doses of 5, 15, 50 mg/kg/day, males were treated for 9 weeks prior to and throughout mating and females were treated 2 weeks prior to mating and throughout mating until gestation day 7. No adverse effect on fertility was observed at 50 mg/kg/day (systemic exposures up to 10 times human exposure at 40 mg/kg/day based on AUC comparisons). In testicles of dogs treated with rosuvastatin at 30 mg/kg/day for one month, spermatid giant cells were seen. Spermatid giant cells were observed in monkeys after 6-month treatment at 30 mg/kg/day in addition to vacuolation of seminiferous tubular epithelium. Exposures in the dog were 20 times and in the monkey 10 times human exposure at 40 mg/kg/day based on body surface area comparisons. Similar findings have been seen with other drugs in this class. **Pregnancy** **Pregnancy Category X** See CONTRAINDICATIONS. Rosuvastatin may cause fetal harm when administered to a pregnant woman. Rosuvastatin is contraindicated in women who are or may become pregnant. Safety in pregnant women has not been established. There are no adequate and well-controlled studies of rosuvastatin in pregnant women. Rosuvastatin crosses the placenta and is found in fetal tissue and amniotic fluid at 3% and 20%, respectively, of the maternal plasma concentration following a single 20 mg/kg oral gavage dose on gestation day 16 in rats. A higher fetal tissue distribution (25% maternal plasma concentration) was observed in rabbits after a single oral gavage dose of 1 mg/kg on gestation day 18. If this drug is administered to a woman with reproductive potential, the patient should be apprised of the potential hazard to a fetus. In female rats given oral gavage doses of 5, 15, 50 mg/kg/day rosuvastatin before mating and continuing through day 7 postcoitus results in decreased fetal body weight (female pups) and delayed ossification at the high dose (systemic exposures 10 times human exposure at 40 mg/kg/day based on AUC comparisons). In pregnant rats given oral gavage doses of 2, 20, 50 mg/kg/day from gestation day 7 through lactation day 21 (weaning), decreased pup survival occurred in groups given 50 mg/kg/day, systemic exposures ≥ 12 times human exposure at 40 mg/kg/day based on body surface area comparisons. In pregnant rabbits given oral gavage doses of 0.3, 1, 3 mg/kg/day from gestation day 6 to lactation day 18 (weaning), exposures equivalent to human exposure at 40 mg/kg/day based on body surface area comparisons, decreased fetal viability and maternal mortality was observed. Rosuvastatin was not teratogenic in rats at $\leq 25 \text{ mg/kg/day}$ or in rabbits $\leq 3 \text{ mg/kg/day}$ (systemic exposures equivalent to human exposure at 40 mg/kg/day based on AUC or body surface comparison, respectively). **Nursing Mothers** It is not known whether rosuvastatin is excreted in human milk. Studies in lactating rats have demonstrated that rosuvastatin is secreted into breast milk at levels 3 times higher than that obtained in the plasma following oral gavage dosing. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from rosuvastatin, a decision should be made whether to discontinue nursing or administration of rosuvastatin taking into account the importance of the drug to the lactating woman. **Pediatric Use** The safety and effectiveness in pediatric patients have not been established. Treatment experience with rosuvastatin in a pediatric population is limited to 8 patients with homozygous FH. None of these patients was below 8 years of age. **Geriatric Use** Of the 10,275 patients in clinical studies with rosuvastatin, 3,159 (31%) were 65 years and older, and 698 (6.8%) were 75 years and older. The overall frequency of adverse events and types of adverse events were similar in patients above and below 65 years of age. (See WARNINGS, Myopathy/Rhabdomyolysis.) The efficacy of rosuvastatin in the geriatric population (≥ 65 years of age) was comparable to the efficacy observed in the non-elderly. **ADVERSE REACTIONS** Rosuvastatin is generally well tolerated. Adverse reactions have usually been mild and transient. In clinical studies of 10,275 patients, 3.7% were discontinued due to adverse experiences attributable to rosuvastatin. The most frequent adverse events thought to be related to rosuvastatin were myalgia, constipation, asthenia, abdominal pain, and nausea. **Clinical Adverse**

Supplement to Circulation

Volume 110 Number 17 October 26, 2004

Abstracts From Scientific Sessions 2004
Morial Convention Center
New Orleans, Louisiana
November 7–10, 2004

Named and Invited Lectures

2004 Russell Ross Memorial Lectureship in Vascular Biology <i>Garret A. FitzGerald, MD</i>	III-A
2004 George Lyman Duff Memorial Lecture <i>Henry N. Ginsberg, MD</i>	III-A
2004 Sol Sherry Distinguished Lecture in Thrombosis <i>Denisa D. Wagner, PhD</i>	III-A
2004 Thomas W. Smith Memorial Lecture <i>Donald M. Bers, PhD</i>	III-B
2004 George E. Brown Memorial Lecture <i>David A. Kass, MD, FAHA</i>	III-B
2004 Dickinson W. Richards Memorial Lecture <i>Kenneth Weir, MD, FAHA</i>	III-C
2004 Katharine A. Lembright Award <i>Diane M. Becker, RN, MPH, ScD, FAHA</i>	III-C
2004 William W. L. Glenn Lecture <i>Timothy Gardner, MD</i>	III-C
2004 William J. Rashkind Memorial Lecture <i>Betsy Dresser, PhD</i>	III-D
2004 T. Duckett Jones Memorial Lecture <i>Jane W. Newburger, MD, MPH</i>	III-D
2004 Charles T. Dotter Memorial Lecture <i>Lewis Wexler, MD</i>	III-D
2004 Laennec Society Lecture <i>Nanette K. Wenger, MD, FAHA</i>	III-E
2004 Ancel Keys Lecture <i>Darwin R. Labarthe, MD, MPH, PhD</i>	III-E

Note: Abstract numbers are arranged to coincide with abstract titles in the *American Heart Association's Scientific Sessions 2002 Program*.

Subject and author indexes are keyed to abstract numbers. Authors of all Named and Invited Lectures and New and Young Investigator Award/Prize Abstracts are presented in the expanded Table of Contents within this Abstracts issue. Indexes are located at the back of this issue. All other supplements to this volume of *Circulation* will be indexed in the last December issue.

CIRCULATION (ISSN 0009-7322) is published weekly except combined the first two weeks in January and the last two weeks in December by Lippincott Williams & Wilkins at 16522 Hunters Green Parkway, Hagerstown, MD 21740. Business offices are located at 530 Walnut Street, Philadelphia, PA 19106-3621. Production offices are located at 351 West Camden Street, Baltimore, MD 21201-2436. Individuals may subscribe for their personal use at the following rates: \$206 for members of an American Heart Association scientific council and \$289 for nonmembers; international: \$353 for members of an American Heart Association scientific council and \$495 for nonmembers. Periodicals postage paid at Hagerstown, MD, and additional mailing offices. POSTMASTER: Send address changes to CIRCULATION, American Heart Association, Lippincott Williams & Wilkins, 16522 Hunters Green Parkway, Hagerstown, MD 21740.

2004 Lewis K. Dahl Memorial Lecture*Rhian M. Touyz MD, PhD* III-F**2004 Robert I. Levy Endowed Lecture in Lipid Metabolism***Ronald M. Krauss, MD, FAHA* III-F**New and Young Investigator Award/Prize Abstract Finalists****Lewis N. and Arnold M. Katz Basic Science Research Prize for Young Investigators***Karl-Ludwig Laugwitz, Alessandra Moretti, Jason Lam, Peter Gruber, Yinhong Chen, Sarah Woodard, Lizhu Lin, Chen-Leng Cai, Michael Reth, Sylvia Evans, Kenneth R. Chien* III-G*Patrick Most, Mirko Voelkers, Sven T. Pleger, Erhe Gao, Melanie Boerries, Andrew Remppis, Hugo A. Katus, Walter J. Koch* III-G*Sathyamangla V. Naga Prasad, Aruni Jayatilleke, Howard A. Rockman* III-H*Da-Zhi Wang, Dongsun Cao, Zhigao Wang, Shijie Li, Eric N. Olson* III-H*Min Xie, Sam C. Wang, Dou Zhang, Arun J.C. Prahash, Hidemasa Oh, Motoaki Sano, Xiaozhen Wang, Jennifer S. Pocius, George E. Taffet, Lloyd H. Michael, Tse-Hua Tan, Mark L. Entman, Michael D. Schneider* III-H**Melvin L. Marcus Young Investigator Awards in Cardiovascular Science***Roselle Abraham, Peter Lim, Charles A. Henrikson, Roland Emokpae, Ronald Li, Leslie Tung, Eduardo Marbán* III-I*Garvan C. Kane, Atta Behfar, D. Fearghas O'Coilain, Roy B. Dyer, Xiao-Ke Liu, Denice M. Hodgson, Santiago Reyes, Takashi Miki, Susumu Seino, Andre Terzic* III-I*Ryoji Koshida, Petra Rocic, Shuichi Saito, William M. Chilian* III-I*Gangjian Qin, Masaaki Ii, Andrea Wecker, Evelyn Bord, Hong Ma, David A. Goukassian, Yung-sup Yoon, Raj Kishore, Yan Zhu, Douglas W. Losordo* III-J*Yao Liang Tang, Yi Tang, Keping Qian, Y. Clare Zhang, Leping Shen, M. Ian Phillips* III-J*Subodh Verma, Chao-Hung Wang, William Stanford, Shu-Hong Li, Richard D. Weisel* III-J**Cournand and Comroe Young Investigators Prizes in Cardiopulmonary and Critical Care***Kimberly A. Boddicker, Yi Zhang, Loyd R. Davies, Richard E. Kerber* III-K*Sebastien Bonnet, M. Evangelos D. Michelakis, Sean McMurtry, Gwyneth Harry, Sandra Bonnet, Alois Haromy, Stephen L. Archer* III-K*Pietro Caironi, Fumito Ichinose, Rong Liu, Rosemary C. Jones, Kenneth D. Bloch, Warren M. Zapol* III-L*Junyan Gu, Deyanira Prastein, Charles White, Jeffrey Manchio, Richard N. Pierson, Bartley P. Griffith, Paul Gurbel, Udaya Tandry, Robert Poston* III-L*Christopher J. Mingone, Sachin A. Gupte, Noorjahan Ali, Richard A. Oeckler, Susan C. Olson, Michael S. Wolin* III-L**Outstanding Research Award in Pediatric Cardiology***Brian Feingold, Seda Selamet, Sang C. Park, Beth F. Printz, C. Becket Mahnke, Shing Lee, Charles S. Kleinman, Donna M. Timchak, William H. Neches, Welton M. Gersony* III-M*Marco Stramba-Badiale, Karine Goulene, Giuliano Bosi, Roberta Bini, Silvia G. Priori, Raffaella Bloise, Lia Crotti, Patrizia Salice, Vlasta Fesslova, Savina Mannarino, Giuseppe Latini, Roberto Giorgetti, Peter J. Schwartz* III-M*Wayne Tworetzky, Audrey C. Marshall, Laura B. Myers, Lisa J. Bergersen, Russell T. Jennings, Carol B. Benson, Louise E. Wilkins-Haug, James E. Lock* III-N**Melvin Judkins Young Investigator Award in Cardiovascular Radiology***Fabao Gao, Bensheng Qiu, Sourav Kar, Xiangcan Zhan, Lawrence V. Hofmann, Xiaoming Yang* III-N*C. Javidan-Nejad, D.C. Lesniak, G.S. Chrysan, J. Zheng, R.J. Gropler, T.K. Pilgram, P.K. Woodard* III-O*Saman Nazarian, David A. Bluemke, Albert C. Lardo, Menekhem M. Zviman, Katherine C. Wu, Stanley Watkins, Clerio Azevedo, Timm L. Dickfeld, Glenn R. Meininger, Ronald D. Berger, Hugh Calkins, Joao A. Lima, Henry R. Halperin* III-O*Monvadi B. Srichai, Chelif Junor, L. Leonardo Rodriguez, Arthur E. Stillman, Richard A. Grimm, Michael L. Lieber, Joan A. Weaver, Nicholas G. Smedira, Richard D. White* III-O**Vivian Thomas Young Investigator Award***Sydney L. Gaynor, Hersh S. Maniar, Jeffrey B. Bloch, Paul Steendijk, Marc R. Moon* III-P*Faraz Kerendi, Michael E. Halkos, Hajime Kin, Joel S. Corvera, Daniel J. Brat, Mary B. Wagner, Jakob Vinten-Johansen, Zhi-Qing Zhao, Joseph M. Forbess, Kirk R. Kanter, Paul M. Kirshbom* III-P*Paul V. Kochupura, Evren U. Azeloglu, Damon J. Kelly, Sergey V. Doronin, Stephen F. Badylak, Irvin B. Krukenkamp, Ira S. Cohen, Glenn R. Gaudette* III-P*Stephen Kolakowski Jr, Mark F. Berry, Todd Grand, Omar Fisher, M. Astrid Moise, Jeffrey Cohen, Vivian Hsu, Y. Joseph Woo* III-Q*Frank Langer, Filiberto Rodriguez, Saskia Ortiz, Allen Cheng, Mary K. Zasio, David Liang, George T. Daughters, Neil B. Ingels, D. Craig Miller* III-Q*Pierre Voisine, Jian Li, Cesario Bianchi, Tanveer A. Khan, Marc Ruel, Shu-Hua Xu, Jun Feng, Audrey Rosinberg, Tamer Malik, Yasunari Nakai, Frank W. Sellke* III-R

Martha N. Hill New Investigator Awards

Wen-Wen Li, Anita L. Stewart, Nancy Stotts, Erika S. Froelicher III-R
Philip Moons, Kristien Van Deyk, Kristel Marquet, Els Raes, Leentje De Bleser, Sabina De Geest, Werner Budts III-R
Lorraine S. Evangelista, Kathleen Dracup, Lynn Doering, Debra K. Moser, Jon Kobashigawa III-S

Elizabeth Barrett-Connor Research Award in Epidemiology and Prevention for Investigators in Training

Matthew Allison, Michael Criqui, Elena Ho, Julie Denenberg III-T
Khawaja Afzal Ammar, Steven J. Jacobsen, Richard J. Rodeheffer III-T
Apoor S. Gami, Daniel E. Howard, Eric J. Olson, Virend K. Somers III-T
Frederick A. Masoudi, Charles Baillie, Yongfei Wang, Edward P. Havranek, JoAnne Micale Foody, John F. Steiner, Harlan M. Krumholz III-U
Nona Sotoodehnia, David Siscovick, Matteo Vatta, Bruce Psaty, Russell Tracy, Jeffrey Towbin, Rozenn Lemaitre, Thomas Rea, Peter Durda, Joel Chang, Thomas Lumley, Lewis Kuller, Greg Burke, Susan Heckbert III-U
Tianying Wu, Susan E. Hankinson, Walter C. Willett, Edward Giovannucci III-V

Samuel A. Levine Young Clinical Investigator Awards

Michaela Scheuermann-Freestone, Damian Tyler, George K. Radda, Stefan Neubauer, Kieran Clarke III-W
Michelle M. Kittleson, Shui Q. Ye, Rafael A. Irizarry, John V. Conte, Giovanni Parmigiani, Leslie W. Miller, Yingjie Chen, Jennifer L. Hall, Joe G.N. Garcia, Joshua M. Hare III-W
Hidenobu Koga, Seigo Sugiyama, Kiyotaka Kugiyama, Keisuke Watanabe, Hironobu Fukushima, Tomoko Tanaka, Tomohiro Sakamoto, Michihiro Yoshimura, Hideaki Jinnouchi, Hisao Ogawa III-W
Francesca Nesta, Maire Leyne, Charles Simpson, Daisy Dai, Jane E. Marshall, Chaim Yosefy, Judy Hung, Susan A. Slaughaupt, Robert A. Levine III-X
Khim Leng Tong, Sanjiv Kaul, Xin-Qun Wang, Diana Rinkevich, Saul Kalvaitis, Todd Belcik, Wolfgang Lepper, William A. Foster, Kevin Wei III-X

Laennec Society Young Clinician Award

Marcus Y. Chen, William A. Baker, Bryan R. Haugen, Robert A. Quaipe III-Y
Andrew C. Lee, Elyse Foster, Yerem Yeghiazarians III-Y
Molly Ware, Eric Awtry, Michael Klein, Thomas Ryan III-Y

NPAM New Investigator Award

M. Robinson, M. Scheuermann-Freestone, P. Leeson, K. Clarke, S. Neubauer, F. Wiesmann III-Z

Abstracts From the Scientific Sessions 2004 III-1


Author Index III-837

Subject Index III-893

Abstracts From the Resuscitation Science Symposium III-1095

American Heart Association Embargo Policy for Scientific Sessions Abstracts

Abstracts, lectures, and other presentations included in the Scientific Sessions 2004 Supplement published in the Oct. 26 issue of *Circulation: Journal of the American Heart Association* are embargoed for release at the time of presentation at the American Heart Association's Scientific Sessions 2004 (Nov. 7-10) and information may not be released before then. Presentation times are listed in the Scientific Sessions 2004 Final Program.



Ride the **CREST**TM

Please Visit
Booth #2547


CRESTOR[®]
rosuvastatin calcium

CRESTOR is a registered trademark and "Ride the CREST" is a trademark of the AstraZeneca group of companies.

Please visit our Web site at www.crestor.com

AstraZeneca 

©2004 AstraZeneca Pharmaceuticals LP. All rights reserved. 217471 01/04

Supplement to
Circulation

Abstracts From Scientific Sessions 2004

Committee on Scientific Sessions Program

The American Heart Association is most grateful to members of the Committee on Scientific Sessions Program for their continued dedication and leadership throughout the year in planning the program.

Raymond J Gibbons, Chairman
Timothy Gardner, Vice Chairman

Council Representatives

Donna Arnett, Functional Genomics and Translational Biology Interdisciplinary Working Group
Kenneth Bloch, Cardiopulmonary Perioperative and Critical Care
Roberto Bolli, Member-at-Large
Lawrence Brass, Stroke Council
Blase A. Carabello, Clinical Cardiology
Arlene Chapman, Kidney in Cardiovascular Disease
Mark A. Creager, Atherosclerotic Peripheral Vascular Disease Interdisciplinary Working Group
Alan Daugherty, Arteriosclerosis, Thrombosis, and Vascular Biology
Christi Deaton, Cardiovascular Nursing
David D. Gutterman, Basic Cardiovascular Sciences
David M. Herrington, Epidemiology and Prevention
Mariell Jessup, Member-at-Large
Alan Kadish, Member-at-Large
John Kugler, Cardiovascular Disease in the Young
Bruce Lerman, American College of Cardiology, Liaison
George A. Mensah, Member-at-Large
Richard C. Pasternak,* Member-at-Large
Robert C. Robbins, Cardiovascular Surgery and Anesthesia
David Robertson, High Blood Pressure Research
Beatriz Rodriguez, Nutrition, Physical Activity, and Metabolism
Arthur Stillman, Cardiovascular Radiology and Intervention
Robert Talbert, Member-at-Large
Gordon F. Tomaselli, Basic Cardiovascular Sciences
William S Weintraub, Quality of Care and Outcomes Research Interdisciplinary Working Group
William Wijns, European Society of Cardiology, Liaison

American Heart Association

Alice K. Jacobs, President
Augustus O. Grant, Immediate Past President
Rose Marie Robertson, Chief Science Officer

*Former member

Supplements to *Circulation* Published in 2004

Supplements to *Circulation* are published occasionally and for archival purposes may be bound with the regular issue. Supplement page numbers are preceded by a Roman numeral and a hyphen. Supplements are indexed with the regular issue with the exception of the Abstracts issue, which is indexed within that issue.

Diagnosis of Venous Thromboembolism Supplement

Gregory L. Moneta, MD, *Guest Editor*
Volume 109, Issue 12. March 30, 2004. Pages I-1-I-27.

Vascular Effects of Statins

Peter Libby, MD, *Guest Editor*
Volume 109, Issue 21. June 1, 2004. Pages II-1-II-48.

Atherosclerosis: Evolving Vascular Biology and Clinical Implications

Antonio M. Gotto, Jr, MD, DPhil, *Guest Editor*
Volume 109, Issue 23. June 15, 2004. Pages III-1-III-57.

Markers of Malign Across the Cardiovascular Continuum: Interpretation and Application

Victor J. Dzau, *Guest Editor*
Volume 109, Issue 25. June 29, 2004. Pages IV-1-IV-58.

Treatment of Venous Thromboembolism

Jeffrey I. Weitz, MD, *Guest Editor*
Volume 110, Issue 9. August 31, 2004. Pages I-1-I-34.

Cardiovascular Surgery Supplement 2004

Robert C. Robbins, MD, *Guest Editor*
Volume 110, Issue 14. September 14, 2004. Pages II-1-II-286.

Abstracts From Scientific Sessions 2004

Volume 110, Issue 17. October 26, 2004. Pages III-1-III-1102.

For rate and order information, contact

Lippincott Williams & Wilkins
Subscription Fulfillment Dept
351 West Camden Street
Baltimore, MD 21201-2436 USA
Telephone (410) 361-8080
Fax (410) 361-8048

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION

Editor-in-Chief

Joseph Loscalzo

Deputy Editor
Joseph A. Vita

Senior Associate Editor
Elliott M. Antman

Associate Editors

Gary J. Balady Emelia J. Benjamin Wilson S. Colucci Ravin Davidoff Jane E. Freedman John F. Keaney, Jr
Jane A. Leopold Jane W. Newburger Douglas B. Sawyer Richard J. Shemin William G. Stevenson James E. Udelson Kenneth Walsh

Africa

Lionel Opie

Asia

Shigetake Sasayama

Australia/New Zealand

Harvey D. White

Central/South America

Edgardo Escobar

Europe

Thomas F. Lüscher

Staff

Managing Editor, *Karen Barry*

Assistant Managing Editors, *Dale R. Golden* and *Doreen E. Kriegl*

Consultants

Immediate Past Editor, *James T. Willerson*

Statistical Consultant, *Martin G. Larson*

Editorial Board

Philip Ades
Charles Antzelevitch
Paul W. Armstrong
Robert J. Bache
Donald S. Baim
William H. Barry
Robyn J. Barst
Craig T. Basson
Anton E. Becker
George E. Beller
Ivor J. Benjamin
Bradford C. Berk
Michel E. Bertrand
Christoph Bode
Roberto Bolli
Robert O. Bonow
Jeffrey S. Borer
Penelope A. Boydén
Eugene Braunwald
Michael R. Bristow
Joan Heller Brown
Nancy J. Brown
Martina Brueckner
Robert M. Califf
Francois Cambien
Richard O. Cannon III
Oscar Carretero
Alan Chait
Bernard R. Chaitman
Peng-Sheng Chen
William M. Chilian
Alexander Clowes
David J. Cohen
Lawrence H. Cohn
Antonio Colombo
John P. Cooke
Delos M. Cosgrove
Filippo Crea
Mark A. Creager
Henry J. Dargie
John E. Deanfield
Linda L. Demer
David Dichek
John P. DiMarco
Stephanie Dimmeler
Gerald W. Dorn II
Helmut Drexler
Victor Dzau
Elazer R. Edelman
Eric J. Eichhorn
Stephen G. Ellis
Stephen E. Epstein

Michael D. Ezekowitz
Erling Falk
David P. Faxon
Douglas T. Fearon
James J. Ferguson III
Loren J. Field
Toren Finkel
Michael Fishbein
Glen I. Fishman
Desmond J. Fitzgerald
Garret A. FitzGerald
Alan M. Fogelman
Aaron R. Folsom
Thomas L. Force
Gregory L. Freeman
Victor Froelicher
Curt D. Furberg
Valentin Fuster
William H. Gaasch
Marie D. Gerhard-Herman
Bernard J. Gersh
Gary Gibbons
Raymond Gibbons
C. Michael Gibson
Samuel S. Gidding
Richard F. Gillum
Charles J. Glueck
Pascal J. Goldschmidt
Paolo Golino
Stephen S. Gottlieb
Augustus O. Grant
Paul A. Grayburn
Scott M. Grundy
Steven M. Haffner
Michel Haïssaguerre
Christian W. Hamm
Goran K. Hansson
Gerd Hasenfuss
Otto M. Hess
Charles B. Higgins
Joseph A. Hill
L. David Hillis
Karen Hirschi
Helen H. Hobbs
Judith S. Hochman
David R. Holmes, Jr
Jeffrey D. Hosenpud
O. Wayne Isom
David A. Kass
Arnold M. Katz
Sanjiv Kaul
Chuji Kawai

David M. Kaye
Mark T. Keating
Daniel P. Kelly
Dean J. Keretiaikes
Morton J. Kern
Spencer B. King III
Neal S. Kleinman
George J. Klein
Robert A. Kloner
Walter J. Koch
Marvin A. Konstam
Raymond T. Kwong
Kathryn G. Lamping
Leslie Leinwand
Bruce B. Lerman
E. Douglas Lewandowski
Martin M. LeWinter
James K. Liao
Peter Libby
Stephen B. Liggett
A. Michael Lincoff
William C. Little
James E. Lock
Douglas W. Losordo
Benedict R. Lucchesi
Lynn Mahony
Alberto Malliani
Donna M. Mancini
Teri Manolio
Eduardo Marbán
Andrew R. Marks
Barry J. Maron
John F. Martin
Attilio Maseri
Jay W. Mason
John E. Mayer, Jr
Hugh A. McAllister, Jr
Jeanie B. McMillin
Elizabeth M. McNally
Jawahar L. Mehta
D. Douglas Miller
Jeffrey D. Molkentin
Fred Morady
Ryuichi Morishita
David A. Morrow
Elizabeth Murphy
Charles E. Murry
Anthony J. Muslin
Elizabeth G. Nabel
Ryozo Nagai
Claudio Napoli
Toshio Ogihara

Ira Ockene
Eric N. Olson
William W. O'Neill
Suzanne Oparil
Catherine M. Otto
Milton Packer
Igor F. Palacios
Julio C. Palmaz
Julio A. Panza
Carlo Patrono
Cam Patterson
Dudley J. Pennell
Carl J. Pepine
Joseph K. Perloff
Mark A. Pfeffer
Mariann R. Piano
Michael H. Picard
Peipei Ping
Bertram Pitt
Gerald M. Pohost
Silvia G. Priors
Eric N. Prystowsky
Karin Pryzklenk
Reed Pyeritz
Marlene Rabinovitch
Daniel J. Rader
Shahbudin H. Rahimtoola
Nathaniel Reichek
Paul M. Ridker
Jeffrey Robbins
Robert C. Robbins
Robert Roberts
William C. Roberts
Rose Marie Robertson
Michael R. Rosen
Jean-Lucien Rouleau
Marschall S. Runge
Frank M. Sacks
Jeffrey E. Saffitz
David J. Sahn
Andrew I. Schafer
Hartzell V. Schaff
Jutta Schaper
Wolfgang Schaper
Melvin Scheinman
Heinrich R. Schelbert
Ann Marie Schmidt
Arnold Schwartz
Ketty Schwartz
Peter J. Schwartz
Robert J. Schwartz

Robert S. Schwartz
Stephen M. Schwartz
Christine Seidman
Jonathan Seidman
Prediman K. Shah
Maarten L. Simoons
Daniel I. Simon
Paul C. Simpson
Burton E. Sobel
Michael J. Sole
Virend K. Somers
Sidney C. Smith, Jr
Francis G. Spinale
David H. Spodick
Lynne Warner Stevenson
A. Jamil Tajik
Akira Takeshita
Anne L. Taylor
Pierre Theroux
James D. Thomas
Gianni Tognoni
Gordon F. Tomaselli
Eric J. Topol
Lilliam Valdes-Cruz
Frans J.J. Van de Werf
Dorothy E. Vatner
Stephen F. Vatner
Douglas E. Vaughan
Renu Virmani
Robert A. Vogel
Lars Wallentin
Carole A. Warnes
Hugh Watkins
Karl T. Weber
Keith Webster
William S. Weintraub
James N. Weiss
Stephen Westaby
Josiah N. Wilcox
David O. Williams
Andrew L. Wit
Joseph L. Witzum
Magdi Yacoub
Edward T.H. Yeh
Paul G. Yock
Salim Yusuf
Barry L. Zaret
Andreas M. Zeiher
Michael R. Zile
Douglas P. Zipes
Jay L. Zweier

Special Section Editors

Book Reviews

Thomas J. Ryan

Basic Science for Clinicians

John F. Keaney, Jr

Clinician Update and Patient Page

Elliott M. Antman

Contemporary Reviews in Cardiovascular Medicine

Ravin Davidoff and Michael A. Fifer

Controversies in Cardiovascular Medicine

Mark E. Josephson

Correspondence

Joseph A. Vita

Images in Cardiovascular Medicine

Warren J. Manning

New Drugs and Technologies

Dan M. Roden and Paul S. Teirstein

Scientific Publishing Committee

Ivor Benjamin, *Chair*

Andrew B. Buroker

N.A. Mark Estes III

Jerome Kassirer

Jeffrey Robbins

Diane Scott-Lichter

Robert Shirrell, *Vice Chair*

William Bryson Campbell

Janet H. Fisher

Rita Redberg

Christine E. Seidman

Linda S. Williams

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION

Editorial Correspondence

 (Additional information available on the Internet at <http://www.circulationaha.org/misc/about.shtml>)

EDITORIAL CORRESPONDENCE should be sent to Joseph Loscalzo, MD, PhD, Editor, *Circulation*, 560 Harrison Ave, Suite 502, Boston, MA 02118. Phone 617-414-8780. Fax 617-414-8781. E-mail circ@bu.edu

INSTRUCTIONS TO AUTHORS appear in the first issue of each month. Authors should consult these instructions before submitting manuscripts to *Circulation*.

AUTHOR COSTS include page charges, cost of color figures, and cost of reprints, if ordered.

Statements, opinions, and results of studies published in *Circulation* are those of the authors and do not reflect the policy or position of the American Heart Association, and the American Heart Association provides no warranty as to their accuracy or reliability.

Business Correspondence

 (Additional information available on the Internet at <http://www.circulationaha.org/misc/about.shtml>)

BUSINESS CORRESPONDENCE should be sent to American Heart Association, Lippincott Williams & Wilkins Subscription Fulfillment Dept, 351 West Camden Street, Baltimore, MD 21201-2436. Telephone 800-787-8984. Fax 800-787-8985. E-mail: custserv@lww.com; outside the U.S.: Telephone 410-361-8080; Fax 410-361-8048.

CHANGE OF ADDRESS: Please supply old and new addresses. Allow 4 weeks for changes.

ANNUAL SUBSCRIPTION RATES: U.S.: *Personal* \$303.00; *Institutional* \$576.00; *Single copy* \$17.00. **Outside the U.S.:** *Personal* \$519.00; *Institutional* \$781.00; *Single copy* \$17.00.

Medical professionals and scientists in training may subscribe for \$162 in the United States and \$271 outside the United States if payment is accompanied by a letter from the department chair verifying post held and completion date.

Prepayment is required. Make check, draft, or money order payable to the American Heart Association in US dollars drawn on a US bank, with *Circulation* on the face of the check. To charge with a credit card, include account number, expiration date, and name as it appears on card.

REPRINTS: Authors may purchase reprints of their article by dialing Kathryn Welch, Author Reprint Department, Lippincott Williams & Wilkins at 1-800-341-2258; or, via E-mail at ReprintsGroup@lww.com; an Author Reprint Order Form may be accessed through the Internet at <http://www.services.lww.com/periodicals/author-reprints>. Single copies of reprints are available from the corresponding authors. Reprints in large quantities, for commercial or academic use, may be purchased from the publisher. For information and prices: telephone 410-528-4121, E-mail kgray@lww.com

Advertising Correspondence

 (Additional information available on the Internet at <http://www.circulationaha.org/misc/about.shtml>)

US ADVERTISING SALES: The Jackson-Gaeta Group, Inc., 33 Small Avenue, Caldwell, NJ 07006. Telephone 973-403-7677. Fax 973-403-7795. Contact Joe Jackson, Paul Nalbandian, or Charles Novak. For media kits, contact Tina Auletta.

JAPAN ADVERTISING SALES: Biomedis International LTD, ASK Ginza Bldg, 10-6, 7-Chome Ginza, Chuo-ku, Tokyo 104-0061, Japan. Telephone (03) 3575-1611. Fax (03) 3569-0155. Contact Hiromu Furukawa. E-mail hiromu-f@biomedis.co.jp

REST OF WORLD SALES: Fran Grega, 3205 Douglas Point Court, Riva, MD 21140. Telephone 410-956-9861. Fax 410-956-9862. E-mail fgrega@comcast.net

ADVERTISING PRODUCTION: Contact Randy Ezell, Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201-2436, telephone 410-528-8533, E-mail rezell@lww.com

SPONSORED GIFT SUBSCRIPTIONS: Contact Carol Bak, Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201-2436, telephone 410-528-4163, fax 410-528-4305, E-mail cbak@lww.com

CLASSIFIED ADVERTISING: Contact Jennifer Williams, Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201-2436, telephone 1-800-528-1843, E-mail jwilliam@lww.com

Advertisements in this issue have been reviewed to comply with the principles governing advertising in American Heart Association publications. A copy of these principles is available on request. **The appearance of an advertisement in an AHA publication is neither an AHA guarantee nor endorsement of the product or service or the claims for the product or service made by the advertiser.**

Secondary Services

INDEXED OR ABSTRACTED in *Biological Abstracts*, *CABS*, *CINAHL*, *Chemical Abstracts*, *Current Contents*, *EMBASE/Excerpta Medica*, and *Index Medicus*.

AVAILABLE ONLINE from Ovid Online, 333 Seventh Avenue, 4th Floor, New York, NY 10001. Telephone 800-950-2035.

MICROFORM EDITION available from University Microfilms International, 300 North Zeeb Road, Ann Arbor, MI 48106-1346. Telephone 313-761-4700. Also available from Princeton Microfilm Corporation, Alexander Road, Box 2073, Princeton, NJ 08540. Telephone 609-452-2066.

AUTHORIZATION TO PHOTOCOPY items from this publication for personal and internal use, the personal or internal use of specific clients, or for educational use, is granted by the American Heart Association on the condition that the copier pay the appropriate fee to the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923. Telephone 978-750-8400. Fax 978-750-4744. This consent does not extend to copying for advertising or promotional purposes, for creating new collective works, or for resale. Individuals may make single photocopies for personal, noncommercial use without obtaining permission. For all other use, permission should be sought directly from the Rights & Permissions Desk, Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201-2436; telephone 410-528-4016; fax 410-528-8550; E-mail journalpermissions@lww.com

Circulation (ISSN 0009-7322) is published weekly by Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201-2436.

GST Registration Number 89552 4239 RT.

Printed in the USA. © 2004 American Heart Association, Inc.


of membranous RhoA and phosphorylated ERM in brainstem were greater both in angiotensin II-treated rats and SHR than in WKY. Valsartan reduced the expression levels of membranous RhoA in angiotensin II-treated rats and SHR. In addition, Y-27632 or valsartan reduced the expression levels of phosphorylated ERM in both groups. Subcutaneous infusion of phenylephrine increased SBP to the same level of angiotensin II infusion in WKY. However, it did not alter the expression levels of membranous RhoA and phosphorylated ERM. Conclusions: These results suggest that 1) the pressor response induced by central infusion of angiotensin II is substantially mediated by activation of Rho/Rho-kinase pathway in brainstem via AT1 receptors, 2) this pathway may also be involved in hypertensive mechanism in SHR.

1412

Endothelial Nitric Oxide and Hypertension in Autonomic Failure

Alfredo Gamboa, Cyndya Shiao, Andre Diedrich, Bonnie K Black, Ginnie Farley, Satish R Raj, David Robertson, Italo Biaggioni; Vanderbilt Univ, Nashville, TN

More than half of patients with autonomic failure (AF) have severe supine hypertension despite low or unresponsive norepinephrine levels and often undetectable plasma renin activity. Supine hypertension is related to increased vascular resistance but the mechanism is not known. To test the hypothesis that nitric oxide deficiency contributes to supine hypertension we blocked endogenous nitric oxide synthase with L-NMMA in 5 AF patients and 7 normal controls (supine SBP 173 ± 6 and 107 ± 5 mmHg, respectively). Systolic blood pressure (SBP) was normalized to 110 mmHg in AF with graded head-up tilt, and baroreflexes were eliminated with trimethaphan in normal controls to mimic autonomic failure. The pressor response to graded doses of L-NMMA was shifted to the left in AF (Figure); The dose necessary to increase SBP by 30 mmHg was 3.4-fold lower in AF compared to controls (136 ± 24 and 465 ± 103 $\mu\text{g}/\text{k}/\text{min}$ respectively, $p < 0.02$). In conclusion, contrary to our original hypothesis, our results suggest an increased tonic release of nitric oxide in AF. Thus, NO deficiency does not contribute to supine hypertension in autonomic failure. On the contrary, this enhanced tonic NO may contribute to orthostatic hypotension in these patients.



1413

Oral Administration of a Mineralocorticoid Receptor Antagonist Reduces Brain, Heart, and Blood-borne Proinflammatory Cytokines in Heart Failure

Yu-Ming Kang, Carver College of Med, Univ of Iowa, Iowa City, IA; Ralph F Johnson, Univ of Iowa, Iowa City, IA; Zhi-Hua Zhang, Carver College of Med, Univ of Iowa, Iowa City, IA; Robert M Weiss, Carver College of Med, Univ of Iowa and VA Med Ctr, Iowa City, IA; Alan K Johnson, Univ of Iowa, Iowa City, IA; Robert B Felder, Carver College of Med, Univ of Iowa and VA Med Ctr, Iowa City, IA

Introduction: Brain and blood-borne cytokines may contribute to neurohumoral excitation in heart failure (HF). We previously reported that blockade of mineralocorticoid receptors (MR) in the central nervous system with spironolactone (SL) reduces circulating tumor necrosis factor (TNF)- α in HF rats. The effect of SL on proinflammatory cytokines (PIC) in the brain and on other important circulating PIC - interleukin (IL)-1 β and IL-6 - was not determined. **Hypothesis:** Chronic treatment with oral SL will reduce brain and blood-borne PIC in rats with HF following MI. **Methods and Results:** Rats underwent coronary artery ligation to induce MI ($48.2 \pm 2.0\%$ of left ventricle, with ejection fraction of $35.5 \pm 4.1\%$ by echocardiography), or sham surgery (SHAM). Six weeks later, immunohistochemistry of the paraventricular nucleus (PVN) of hypothalamus, a region critical to cardiovascular regulation, revealed more PVN neurons (MI vs SHAM, $**P < 0.01$) positive for TNF- α ($59.5 \pm 3.3\%$ vs $10.8 \pm 0.9\%$) and IL-1 β ($70.7 \pm 3.9\%$ vs $13.8 \pm 1.2\%$) in MI ($n=6$) than in SHAM ($n=6$) rats. Double staining demonstrated that these neurons were distributed among PVN neurons expressing Fra-like immunoreactivity, indicating chronic neuronal activation. MI rats ($n=6$) treated with SL (1 mg/kg/day orally for 6 weeks) had fewer (MI+SL vs MI, $\#P < 0.01$) Fra-like positive PVN neurons ($85.5 \pm 5.4\%$ vs 183.8 ± 5.0), and fewer PVN neurons positive for TNF- α ($22.4 \pm 1.8\%$ vs $32.4 \pm 1.7\%$) and IL-1 β ($19.1 \pm 1.3\%$ vs $38.4 \pm 2.1\%$). Levels of TNF- α , IL-1 β and IL-6 in brain and heart tissues and in plasma were also lower in MI rats treated with SL (see table). **Conclusion:** In rats with ischemia-induced heart failure, orally administered SL has a global inhibitory influence on the appearance of proinflammatory cytokines in brain, heart and plasma. The beneficial influence of MR antagonism in patients with HF may result at least in part from blocking aldosterone-induced cytokine synthesis. (Table: $*P < 0.05$ MI+SL vs MI+VEH)

Group	plasma IL-1 β (pg/ml)	plasma IL-6 (pg/ml)	heart IL-1 β (pg/mg protein)	hypothalamus IL-1 β (pg/mg protein)	hypothalamus TNF- α (pg/mg protein)	brain-stem TNF- α (pg/mg protein)	brain-stem IL-6 (pg/mg protein)	cortex IL-1 β (pg/mg protein)	heart/BW Ratio (mg/g)	lung/BW (mg/g)
MI+VEH (n=7)	131.1 \pm 10.9	119.5 \pm 9.7	53.4 \pm 6.5	47.1 \pm 7.9	6.8 \pm 0.7	6.1 \pm 0.9	67.1 \pm 10.1	26.4 \pm 4.1	7.2 \pm 0.2	13.5 \pm 0.6
MI+SL (n=7)	57.5 \pm 3.1*	53.3 \pm 8.6*	32.6 \pm 5.9*	29.4 \pm 5.1*	3.8 \pm 0.6*	2.8 \pm 0.5*	39.4 \pm 5.5*	25.8 \pm 5.2	6.4 \pm 0.3*	12.1 \pm 0.6*
SHAM+SL (n=7)	48.2 \pm 3.6	33.7 \pm 2.8	23.7 \pm 6.5	17.4 \pm 4.8	2.8 \pm 0.7	2.1 \pm 0.6	25.3 \pm 4.8	24.3 \pm 4.9	3.3 \pm 0.1	5.2 \pm 0.3
SHAM+VEH (n=6)	53.9 \pm 2.4	36.1 \pm 2.6	27.6 \pm 5.4	18.8 \pm 4.0	3.2 \pm 0.9	2.4 \pm 0.8	28.0 \pm 8.7	23.6 \pm 6.5	3.4 \pm 0.1	5.1 \pm 0.3

Pulmonary Arterial Hypertension: New Therapies

Subspecialty: Integrative Biology

Wednesday

Ernest N Morial Convention Center, Hall 12

Abstracts 1414-1418

1414

Inhaled Treprostinil Sodium (TRE) For the Treatment of Pulmonary Hypertension

Robert Voswinckel, Beate Enke, Andre Kreckel, Frank Reichenberger, Stefanie Krick, Henning Gall, Tobias Gessler, Thomas Schmehl, Markus G Kohstall, Friedrich Grimminger, Hossein A Ghofrani, Werner Seeger, Horst Olschewski; Univ Hosp Giessen, Giessen, Germany

Objective: To evaluate the effects of inhaled TRE on pulmonary hemodynamics and gas exchange in severe pulmonary hypertension (PH) and to assess safety, tolerability and clinical efficacy in patients with severe PH. **Background:** TRE is a stable prostacyclin analogue that has been approved for treatment of pulmonary arterial hypertension as a continuous subcutaneous infusion. Iloprost, another prostacyclin analogue, has been shown to be efficacious in a randomised controlled study as repetitive inhalation. **Methods:** In an open-label study a preservative free solution of inhaled TRE was applied to 17 patients with severe pulmonary hypertension during Swan-Ganz catheter investigation. Patients received a TRE inhalation by use of the pulsed OptiNeb ® ultrasound nebulizer (3 single breaths, TRE solution 600 $\mu\text{g}/\text{ml}$). Hemodynamics were observed for 2 hours. Two patients with idiopathic PAH received compassionate treatment with 4 inhalations of TRE per day after the acute test. **Results:** Patients (male/female = 4/13) suffered from iPAH ($n=5$), PAH other ($n=8$) and CTEPH ($n=4$); PVR 948 ± 112 $\text{dyn}\cdot\text{s}\cdot\text{cm}^{-5}$, PAP 48.3 ± 2.7 mmHg, PAWP 8.9 ± 0.5 mmHg, CVP 10.8 ± 1.6 mmHg, CO 3.8 ± 0.3 l/min, SvO 2 61.8 ± 1.8 %. TRE inhalation resulted in a sustained, highly pulmonary selective vasodilatation over 120 minutes. Maximum PVR decrease was -31.2 ± 4.5 % after 30 min. PVR and SVR at 120 minutes after inhalation were 89.2 ± 4.2 % and 101.0 ± 4.0 % of the baseline values, respectively. The AUC for the observation period (120min) was -22.9 ± 3.8 % for PVR and -4.9 ± 3.2 % for SVR. The compassionate use patients have been treated for more than 3 months. In both patients NYHA class improved (from IV to III and from III to II), and six minute walk increased (from 0 m (bedridden) to 143 m, and from 310 m to 486 m, respectively). No side effects have been observed by the patients during long-term treatment. **Conclusion:** Inhaled TRE shows strong pulmonary selective vasodilatory efficacy with a long duration of effect following single acute dosing. Tolerability is excellent even at high drug concentrations and short inhalation times (3 breaths). Long-term treatment effects are very promising. The current results warrant controlled studies investigating this approach in a larger series of patients. Supported by Lung RX

1415

Rho-kinase in Pulmonary Hypertension

Ken Ishikura, Norikazu Yamada, Akihiro Tsuji, Satoshi Ota, Mashio Nakamura, Masaaki Ito, Naoki Isaka, Takeshi Nakano; Mie Univ Sch of Med, Tsu, Japan

Objectives: Pulmonary hypertension (PH) is a poor prognostic disease with limited treatment. Rho-kinase is involved in the pathophysiology of several diseases underlying smooth muscle hypercontraction. But the role of is unknown. The purpose of this preliminary report was to indicate the efficacy of fasudil, a Rho-kinase inhibitor in patients with pulmonary hypertension using interventional hemodynamic assessment. **Methods:** Fasudil was intravenously injected in 10 patients (9 female, mean \pm SD, 46 ± 15 years, NYHA II $n=2$, III $n=7$, IV $n=1$) with primary ($n=5$) and secondary ($n=5$) PH who were not received any vasodilator. Fasudil was administered 30mg with 1mg/min. Hemodynamic data were measured using Swan-Ganz catheter until 60 minutes after starting administration of fasudil. Hemodynamic and arterial blood gas data of baseline and the lowest total pulmonary resistance (TPR) time were compared. **Results:** The lowest TPR time was within 30 to 60 minutes after administration. Administration of fasudil significantly decreased TPR from 13.6 ± 6.8 U to 10.3 ± 4.9 U (-23.2 ± 9.2 %, $p < 0.001$) and mean pulmonary arterial pressure (mPAP) from 43.6 ± 14.5 mmHg to 38.8 ± 13.9 mmHg (-11.6 ± 10.4 %, $p < 0.02$). Cardiac index (CI) was significantly increased from 2.39 ± 0.66 L/min/m 2 to 2.74 ± 0.73 L/min/m 2 ($+16.5 \pm 15.1$ %, $p < 0.01$). Although TPR was equally decreased in both primary and secondary PH, the changes in the parameters that prescribed TPR, namely CI and mPAP, were different between the two groups subjects. Increased CI was a major factor into reducing TPR in primary PH, while reduced mPAP

- von der Thüsen, Jan, 846
 von Erffa, Johannes, 2452
 von Eye, Alexander A, 1962
 von Haehling, Stephan, 3016, 3475
 von Harsdorf, Rüdiger, 260
 von Lewinski, Dirk, 1285
 von Lueder, Thomas G, 1094
 von Lukowicz, Tobias, 3743
 von Mering, Gregory O, 2445, 3441, 3620
 von Scheidt, Wolfgang, 2730
 von Wilmowsky, Hubertus, 2444
 von Ziegler, Franz, 2291, 2292
 Voncken, Willem, 1265
 Vonder Muhll, Isabelle, 3625
 Vonderscher, Jacky, 1011
 Vondriska, Thomas M, 980
 Vondriska, Tom, 126
 Vonhof, Stefan, 3730
 Vooletich, Mary, 1761
 Voon, Wen-Choi, 677
 Vora, R, 1309
 Vorlova, Sandra, 956, 1407
 Voros, Szilard, 1670, 3174
 Vorpahl, Marc, 773
 Vos, Marc A, 1531
 Voskas, Daniel, 943
 Voskuil, Michiel, 2179
 Voskuyil, Jan, 1673
 Voss, Gemma B Sr, 3808
 Voss, Reinhard, 1057
 Voswinckel, Robert, 258, 1414
 Votaw, V. Scott, 933
 Voudris, Vassilis, 3402
 Vourvouri, Eleni C, 1996, 2538
 Vranckx, Roger, 281
 Vratsista, Ekaterini, 3643
 Vredevoe, Donna, 1963
 Vricella, Luca A, 2977
 Vrints, Christiaan, 3753
 Vrints, Christiaan J, 235, 1807
 Vu, Jack B, 2441
 Vu, John D, 2441
 Vupputuri, Suma, 3521
 Vyas, Anant K, 2361
 W Hamm, Christian, 2099
 Waagstein, Finn, 631
 Wachtell, Kristian, 2961, 3160, 3375, 3383, 3560
 Wacker, Christian, 2563
 Wackers, Frans J, 3168
 Wada, Atsuyuki, 896
 Wada, Hiroshi, 319, 942, 945
 Wada, Nozomi, 2733, 3307
 Wadanoli, Michael, 709
 Wadghiri, Youssef Z, 274
 Wadsworth, Scott, 1278
 Wagenknecht, Lynne E, 3854
 Waggoner, Alan D, 3561
 Wagner, Anja, 3140
 Wagner, Bettina, 2046
 Wagner, Daniel R, 2089
 Wagner, Erwin F, 1585
 Wagner, Galen S, 1933, 3089
 Wagner, Mary B, 1885
 Wagner, Roger A, 82
 Wagner, Stefan, 282
 Wagoner, Lynne, 2415
 Wagoner, Lynne E, 1874, 2244, 3473
 Wahbi, Karim, 1727
 Wahi, Sudhir, 2736
 Wahl, Denis, 3724
 Wakami, Kazuaki, 1730
 Wakamiya, Nobutaka, 1075
 Wake, Ryotaro, 531, 2728
 Wakimoto, Koji, 482
 Wakino, Shu, 434, 790, 1162
 Wakisaka, Yuko, 3215
 Waksman, Ron, 250, 1817, 2473, 3067
 Walch, Laurence, 887, 1683
 Wald, Ron, 2679
 Waldenström, A, 1726, 1732
 Waldenström, Anders, 40
 Waldo, Albert L, 1395
 Waldron, Michele M, 2835
 Waleffe, André, 3447
 Walienga, Jeanine, 2873
 Walenga, Jeanine M, 2872
 Walenta, Katrin, 2379, 3572
 Walker, Brian R, 1275
 Walker, Esteban, 3115
 Walker, Jamie M, 1575
 Walker, Jennifer D, 1710
 Walker, Johnny E, 3424
 Walker, Joseph C, 489
 Walker, Robert, 1959
 Wall, Barry M, 3343
 Wall, E., 217
 Wallace, Arthur W, 489
 Wallenstein, Sylvan, 3702
 Wallentin, Lars, 1947, 2522, 2675
 Wallick, Don W, 140
 Wallis, Julie, 970
 Walls, Dermot, 573
 Walraevens, Maarten, 2272
 Walsh, Helen A, 3365
 Walsh, Kenneth, 15, 207, 739, 1039, 1073, 1333
 Walsh, Kenneth A, 1392
 Walsh, Kevin, 2631
 Walsh, Mary N, 3131
 Waltenberger, Johannes, 449, 1520, 2299, 3675
 Walter, Dirk H, 715, 1058, 1338, 1679
 Walter, Glenn, 50
 Walther, Claudia, 3574
 Walther, Susanne, 3290
 Walther, Thomas, 1090
 Wamhoff, Brian R, 1249
 Wan, Tina C, 130
 Wang, Andrew, 2601, 2640
 Wang, Bao-Wei, 38
 Wang, Bing, 385, 1769
 Wang, Bing-yin, 338, 3735
 Wang, Bingyin, 1048
 Wang, Bo, 2954
 Wang, Chao-Hung, 7
 Wang, Chunlei, 3111
 Wang, Da-Zhi, 1, 761
 Wang, Dachun, 653
 Wang, Dajun, 453, 992
 Wang, Danming, 378
 Wang, Daowen, 2282
 Wang, Desuo, 1914
 Wang, Duolao, 3478
 Wang, Gang W, 126
 Wang, Guan-Ying, 1389
 Wang, Guangwu, 980
 Wang, Hao, 966
 Wang, Hong, 134, 147, 484, 565, 880
 Wang, Hong-Sheng, 622
 Wang, Huashan, 1637
 Wang, Hui-Lian, 322
 Wang, Huili, 889
 Wang, Jen Yu, 2086
 Wang, Ji-Quan, 2055
 Wang, Jianglan, 2493
 Wang, Jianwen, 524
 Wang, Jie, 14, 2794, 3299, 3370
 Wang, Jiguang, 2276
 Wang, Jinglan, 1631, 1856, 1952, 2485, 2491
 Wang, Jingxiong, 921
 Wang, Kai, 218
 Wang, Lai, 334
 Wang, Li, 107, 1438
 Wang, LiChun, 1580
 Wang, Lingli, 624
 Wang, Lixing, 3405
 Wang, Liyong, 3799
 Wang, Lu, 3687, 3710
 Wang, Mei, 2761, 3305
 Wang, Meijing, 1711
 Wang, Miao, 843
 Wang, Min, 3223
 Wang, Ming-Dong, 1301
 Wang, Nan, 1304, 1306
 Wang, Ning-Ping, 805, 1423, 1706
 Wang, Ningyuan, 3639
 Wang, Ou-Li, 646, 985
 Wang, Paul, 2896
 Wang, Ping, 828
 Wang, Qian, 793
 Wang, Qing, 3434
 Wang, Ray, 1097
 Wang, Renwei, 3685, 3769, 3770
 Wang, Sam C, 3, 632, 1259
 wang, Shan, 1661
 Wang, Shirley, 2366
 Wang, Steven, 1697
 Wang, Sufen, 2489
 Wang, Ting, 1914
 Wang, Tommy J, 3527
 Wang, Wei, 189, 339, 1125, 1268, 1535
 Wang, Wenli, 1383
 Wang, X, 1309
 Wang, Xiangyuan, 2090
 Wang, Xiao, 417
 Wang, Xiao-Li, 1595
 Wang, Xiaofei, 1008
 Wang, Xiaohong, 389
 Wang, Xiaoyin, 806, 1133, 3807
 Wang, Xiaozhen, 3
 Wang, Xin Qun, 1606
 Wang, Xing Li, 3818
 Wang, Xuejun (XJ), 764, 1138
 Wang, Xuerong, 1383
 Wang, Yang, 1439
 Wang, Yi, 2842
 Wang, Yibin, 202, 298, 914, 1081, 1371
 Wang, Yigang, 18, 1339, 1453
 Wang, Yongfei, 3530, 3774
 Wang, Zhigao, 1
 Wang, Zhiguo, 66, 921
 Wang-Polagruto, Janice F, 3587
 Wani, Maqsood, 1339, 1561
 Warabi, Eiji, 1358
 Ward, R. Parker, 1893, 1983
 Ward, Simone M, 448
 Warda, Hazem M, 3229
 Ware, Molly, 2007
 Ware, Stephanie M, 1744
 Warembourg, Henri, 2561a
 Warita, Shun-ichiro, 2315
 Warjngarten, Mauricio, 1500
 Warman, Eduardo, 1646
 Warner, John J, 3849
 Warnes, Carole A, 3022
 Warnholtz, Ascan, 376
 Warnica, J. Wayne, 2415
 Warnick, Russell G, 1477
 Warsy, Irfan A, 2895
 Was, Tomasz, 3074
 Washam, Jeffrey B, 3056, 3235
 Washington, Jason W, 1156
 Wasler, Andrae, 2814
 Wassar, Dawn, 3756
 Wasserman, Bruce A, 3684
 Wasserman, Hal S, 2068
 Wasserman, Karlman, 2296, 2495, 2496
 Wassmann, Sven, 181, 696, 1241
 Wasywich, Cara A, 1720, 3365
 Watanabe, Go, 2922
 Watanabe, Hideki, 2537, 2542
 Watanabe, Hidetsuna, 949
 Watanabe, Ichiro, 2174, 2709
 Watanabe, Ikuyoshi, 1640, 1698
 Watanabe, Kazuhiro, 1640, 1842, 1843
 Watanabe, Kazunori, 2228
 Watanabe, Keisuke, 1607, 1775
 Watanabe, Kenichi, 505
 Watanabe, Mari, 3092
 Watanabe, Masaki, 2197
 Watanabe, Michiko, 444
 Watanabe, Miho, 1544
 Watanabe, Miki, 3269
 Watanabe, Noboru, 2422, 3477
 Watanabe, Nozomi, 2733, 3307
 Watanabe, Sachiro, 2315
 Watanabe, Satoshi, 2588
 Watanabe, Sayaka, 1842, 1843
 Watanabe, Teruo, 1008
 Watanabe, Tetsu, 1224
 Watanabe, Tetsuya, 300, 755
 Watanabe, Tsuyoshi, 3159
 Watano, Keiko, 1311
 Waters, Christopher M, 419
 Waters, David, 1946
 Waters, Richard, 529, 825
 Wathen, Mark S, 1653, 2085, 2362
 Watkins, Craig, 331
 Watkins, Hugh, 54, 57, 970
 Watkins, Hugh C, 2664
 Watkins, Matthew W, 1979
 Watkins, Stanley, 1910
 Watkinson, William P, 1917
 Watson, Karol E, 245, 3708
 Wattanakit, Keattiyaoat, 3610
 Watts, Katie, 3595
 Wawrousek, Eric F, 764
 Way, Kerrie, 104
 Wazni, Oussama, 2703
 Wazni, Oussama M, 1871, 1889, 2712, 3205, 3253
 Weatherall, Mark, 1827
 Weatherley, Beth D, 3610
 Weaver, Joan A, 1911
 Weaver, W Douglas, 3706, 3779
 Weaver, W Douglas, 1751
 Webb, Carolyn M, 2666
 Webb, Carson S, 2958
 Webb, Gary, 3023, 3630
 Webb, Gary D, 3019
 Webb, John G, 2115
 Webb, Randy, 477
 Webb, Randy L, 2407
 Weber, Alberto, 2481
 Weber, Artur A, 381
 Weber, Artur-Aron, 353
 Weber, Christian, 847
 Weber, Karl T, 925, 3343
 Weber, Michael W, 1927
 Weber, Simon, 1727
 Weber, Stefan, 2690
 Weber, Thomas, 3603
 Webster, Keith A, 268, 900
 Webster, Linda, 1760
 Webster, Mark W, 1720
 Wecker, Andrea, 8, 200, 369, 655, 2186
 Weeks, David, 2614
 Wegscheider, Karl, 3489
 Wehinger, Andreas, 557
 Wehrens, Xander H, 307, 610, 1091
 Wei, Chiming, 14, 2095, 3270
 Wei, Heng, 297, 1462
 Wei, Jeanne Y, 1378
 Wei, Kevin, 1606, 3315, 3323, 3459
 Wei, Qin, 577, 578, 580
 Weichenhan, Dieter, 491
 Weichert, Jiri, 1970
 Weidemann, Frank, 2563
 Weidinger, Franz, 1791
 Weidman, William H, 1835
 Weiglein, Andreas H, 1866
 Weihrauch, Julia, 672
 Weil, Joachim, 1110, 1503
 Weil, Max H, 1631, 1856, 1858, 1952, 1958, 2484, 2485, 2491, 2492
 Weil, Max Harry, 2486, 2493
 Weinberg, Alan, 657, 1467
 Weinberg, Kenneth, 2363
 Weinbrenner, Christof, 2253
 Weinert, Lynn, 2759, 3038, 3322, 3449
 Weinheimer, Carla J, 110
 Weinmann, Hanns J, 3175
 Weinmann, Hanns-Joachim, 177
 Weinsaft, Johnathan W, 3490
 Weinsaft, Jonathan W, 3486, 3628
 Weintraub, Neal L, 1344
 Weintraub, William S, 1753, 1961, 2306, 2742, 3026, 3185, 3531
 Weis, Michael, 1565, 1571, 2730
 Weisbrod, Cara J, 3288

- Weisbrod, Robert, 1068
 Weisbrod, Robert M, 1594
 Weisel, John W, 347
 Weisel, Richard D, 7, 1451, 2182
 Weisman, Gary A, 695
 Weisnagel, S. John, 869
 Weiss, Daiana, 1455
 Weiss, Dominik R, 599, 674
 Weiss, Elisabeth, 387
 Weiss, Harvey R, 1433
 Weiss, Ingo, 3260
 Weiss, James N, 458
 Weiss, Melvin B, 1349
 Weiss, Robert G, 1548, 2835, 2993
 Weiss, Robert M, 394, 1404, 1409, 1413
 Weiss, Stephen J, 1458
 Weissbach, Herbert, 900
 Weisskoff, Robert M, 3180
 Weissleder, Ralph, 658, 832, 3179
 Weissman, Neil, 2458
 Weissman, Neil J, 2308, 2724, 3398
 Weisz, Giora, 3013
 Weitzman, Jonathan B, 1276
 Weivoda, Peggy, 2087
 Weldner, Paul W, 1979
 Wellens, Francis, 2272, 2727
 Wellner, Maren, 477
 Wellnhofer, Ernst, 3408
 Welsh, Robert C, 2515
 Welten, Rob J, 1777
 Welty, Francine K, 684, 3590
 Welty, Thomas K, 3544
 Wen, Hong Yuan, 742
 Wen, Jing, 634, 897, 2949
 Wenaweser, Peter, 2875
 Wendel, Hans P, 2368
 Wenderow, Tal, 2075
 Weng, Chih Yuan, 1849
 Weng, Yuguo, 1999, 2936, 2940
 Wenger, Nanette K, 1961
 Wengle, James G, 3019
 Wennbo, Håkan, 591, 1182
 Wennerblom, Bertil Sr, 2929
 Wensel, Roland, 2764
 Wenzel, Beate, 3260
 Werenowicz, Stanislaw, 2571
 Wergeland, Ragnhild, 3263
 Werner, Lael, 99
 Werner, Nikos, 696, 698, 1177, 2379, 3572
 Wernet, Peter, 1141
 Wernovsky, Gil, 3541
 Wessel, Armin, 3633
 Wessel, David L, 2876, 2890
 Wessel, Timothy R, 1685, 2445, 3441, 3620, 3792
 Wessely, Rainer, 773, 1832
 West, Matthew B, 1130
 West, Mike, 3799
 West, Teena, 3291
 Westenberg, Jos J, 2738
 Westendorp, Bart, 1420
 Westerhof, Nico, 522
 Westerhout, Cynthia M, 2513, 2523
 Westermann, Dirk, 1090, 1459, 2963, 3641
 Westfall, Amanda, 2147
 Westfall, Margaret V, 226, 1537
 Westheim, Arne, 1758
 Westin-Eriksson, Annika, 1182
 Westlake, Cheryl, 3415
 Westlake, Cheryl A, 1966
 Weston, Susan A, 3059, 3719
 Wettwer, Erich, 611
 Wexler, Dov, 1200
 Weyand, Cornelia M, 1186, 1187
 Weyand, Michael, 1286
 Weyrich, Andrew, 345
 Whalin, Matthew K, 1455
 Whalley, Gillian A, 3365
 Whang, Brian, 315, 320, 651, 794, 814, 816
 Whang, William, 3604
 Wharton, J M, 3100, 3245
 Wharton, J Marcus, 3357
 Wharton, Thomas P, 1979
 Wheatcroft, Stephen, 416
 Wheatcroft, Stephen B, 162
 Wheeler, Michael B, 72
 Whelton, Paul K, 3588, 3613
 Whisnant, Jack P, 3680
 Whitbourn, Robert, 1804, 2623, 3502, 3852
 White, Anthony R, 378
 White, Audrey, 3284
 White, C. M, 2907
 White, C. Michael, 3268
 White, C. Roger, 993, 1167
 White, Charles, 1194, 2546
 White, Christopher J, 2851, 2998
 White, Harvey, 2682, 2985
 White, Harvey D, 1720
 White, Jennifer K, 3108
 White, Linda W, 3084
 White, Melanie, 1540
 White, Richard D, 1783, 1911
 White, Roger D, 2136
 White, Thomas E, 2982
 White-Williams, C, 3237
 Whiteheart, Sidney W, 349
 Whiting, Bryant M, 3424
 Whitlow, Patrick L, 2068, 3496
 Whitman, Donna, 1817
 Whitmore, Harry, 1392
 Whitney, Edwin J, 3785
 Whittaker, Peter, 309, 1428
 Whyte, Greg, 3366
 Wians, Frank H Jr, 3654
 Wichmann, H.-Erich, 1101, 1103
 Wichter, Thomas, 1737, 1782
 Wickline, Samuel A, 1284, 1463
 Wiebers, David O, 3680
 Wiczorek, David F, 455
 Wiedemann, Michael, 2692
 Wiegand, Stanley, 771
 Wiegand, Uwe K, 1645
 Wieggers, Susan E, 2655
 Wieland, Thomas, 913
 Wielander, Christoph, 3826
 Wienbergen, Harm, 2916
 Wieneke, Heinrich, 1824
 Wiener, Isaac, 1870
 Wier, W G, 792
 Wiesfeld, Ans C, 3255, 3428
 Wiesmann, Frank, 2664, 3793
 Wiethoff, Andrea J, 3180
 Wiggins, S. Scott, 3246
 Wight, Thomas N, 1020
 Wight, Tom, 2383
 Wijesundera, Duminda, 2373
 Wijffels, Maurits C, 2019
 Wijns, William, 318, 1555, 2469, 3006, 3009, 3011, 3398, 3437
 Wik, Lars, 2139
 Wiklund, Olov, 1348
 Wikstrand, John, 3648
 Wikström, G, 1732
 Wikström, Johannes, 591, 1785
 Wilber, David J, 1869, 2530
 Wilcox, Josiah N, 805, 1423
 Wilcox, Robert, 2863
 Wildberger, Joachim Ernst, 2284, 2450
 Wilde, Arthur, 1101
 Wilde, Arthur A, 1254, 1105, 1735, 1741, 2893, 3208
 Wilders, Ronald, 73
 Wildman, Rachel P, 3613, 3740
 Wilensky, Michael, 2468
 Wilensky, Robert, 3187, 3189
 Wilensky, Robert L, 1677, 1702, 1936, 2866, 3267, 3508
 Wilhelm, Francois, 683
 Williams, Richard V, 2076
 Wilke, Andreas, 2424
 Wilkins, Benjamin, 32
 Wilkins, Martin R, 79, 1418
 Wilkins-Haug, Louise E, 1929
 Wilkinson, James, 2631
 Wilkoff, Bruce L, 1648, 1783, 3353
 Will, Melissa L, 3515
 Willecke, Klaus, 720
 Willems, Stephan, 1668, 2527, 2708
 Willenbockel, Christian, 1095, 1421
 Willenbrock, Roland, 800
 Willenheimer, Ronnie, 3796
 Willerson, James T, 549, 653, 892
 Willett, DuWayne, 3709
 Willett, Walter, 3562
 Willett, Walter C, 3582, 3591, 3604, 3739, 3775
 Willette, Robert N, 1158
 Willey, Vincent J, 3708
 William, Maged, 932
 Williams, Alan J, 472
 Williams, Dale, 3655
 Williams, David, 53, 2070, 3187, 3189
 Williams, David L, 1303
 Williams, David O, 1702, 3397
 Williams, G, 2503, 3359
 Williams, G. Melville, 2095
 Williams, Gordon H, 89, 2043
 Williams, Ismee A, 2483
 Williams, Jeff T, 3818
 Williams, Jill, 3596
 Williams, Kathryn, 2435
 Williams, Ken, 3854
 Williams, Kevin J, 3435
 Williams, Lynne, 3146
 Williams, Matthew L, 41, 1987, 2732
 Williams, Monica, 3783
 Williams, Paul T, 3782
 Williams, R. Sanders, 1372, 1379
 Williams, Randy, 2515
 Williams, Richard V, 1838, 2077
 Williams, Ronald B, 1719, 2661, 2962
 Williams, Simon C, 621, 930
 Williams, Steve, 416
 Williams, Stuart K, 902
 Williams, Todd A, 1463
 Williams, Ursula, 2391, 3318
 Williams, William, 3411
 Williams, William G, 2482
 Williamson, Jeff D, 3580
 Willott, Ruth, 57
 Willoughby, Scott, 3261
 Wilmore, Jack H, 435
 Wilson, Alan C, 2042, 3831
 Wilson, Amanda, 3233
 Wilson, Andrew, 3634, 3852
 Wilson, Anthony, 3634, 3852
 Wilson, David, 3790
 Wilson, Hadley, 2322
 Wilson, Kate, 1430
 Wilson, Lance D, 461
 Wilson, Neil, 3024
 Wilson, Peter W, 3581, 375, 3527, 3583, 3746, 3761
 Windecker, Stephan, 2626, 2875, 3011
 Wingate, Suzanne J, 2973
 Wink, David A, 142
 Winkelmann, Bernhard R, 558
 Winkelmann, Kim, 125, 1536
 Winkler, Karl, 558
 Winkler, Ralf, 2916
 Winnier, Deidre A, 3818
 Winowich, Stephen, 2931
 Winowich, Steve, 2713
 Winter, Patrick M, 1463
 Wintermantel, Erich, 1832
 Winters, Stephen L, 2901, 2902
 Wintersperger, Bernd J, 1554
 Wisdom, Paul, 428
 Wissing, Karl Martin, 3368
 Wit, Andrew, 1526
 Witkowski, Piotr, 708
 Witkowski, Piotr P, 1097
 Witt, Christian, 3025
 Witt, Sandra, 3543
 Witt, Sandra A, 1724
 Witt, Tyra A, 697
 Witters, Lee A, 1549
 Wittkowski, Knut M, 3332
 Witzenbichler, Bernhard, 1459
 Witzke, Christian, 714
 Witzum, Joseph, 247
 Witzum, Joseph L, 870
 Wiviott, Stephen D, 2340, 3222
 Wiviott, Steven D, 2124, 3233, 3848
 Wnendt, Stephan, 1558
 Wodnicki, Jan, 2795
 Wofe, Matthew J, 2584
 Wöhrle, Jochen, 1818
 Wojakowski, Wojciech, 238, 1143
 Wojcik, Kimberly, 1637
 Wojnicz, Romuald, 2795
 Wojnowski, Leszek, 3730
 Wojta, Johann, 3257
 Wojta, Johannes, 1157
 Wolak, Arik, 3073
 Wolf, Christopher J, 1794
 Wolf, Cordula M, 1108
 Wolf, David, 1199, 1441
 Wolf, Jean-Pierre, 3064
 Wolf, Matthew J, 1222
 Wolfe, Megan, 242
 Wolfe, Megan L, 684, 3439
 Wolfe, Robert R, 1835
 Wolfe, Rory, 3606
 Wolfel, Eugene E, 2714
 Wolfert, Robert L, 1477, 3786
 Wolfram, Roswitha, 1497
 Wolfram, Roswitha M, 3003
 Wolgemuth, Debra J, 811, 2090
 Wolin, Michael S, 700, 1195, 1236
 Wolinski, David, 3473
 Wolinsky, Claire, 771
 Wolk, Robert, 2058
 Wolkowicz, Paul, 1136
 Wollert, Kai C, 127, 1095, 1145, 1148, 1421, 2051
 Wolpert, Christian, 1106, 2346
 Wolska, Beata M, 1085
 Wolters, Sander L, 2053
 Wolzt, Michael, 1157
 Women's Health Initiative Research Group, for the, 3570
 Wong, Amy J, 1050
 Wong, Cindy W, 179
 Wong, Jack, 1393
 Wong, Johnny, 218
 Wong, K K, 1584
 Wong, K.k., 1583
 Wong, Kenneth Y, 2697
 Wong, Louise, 875
 Wong, Michelle H, 844
 Wong, Nathan, 2613
 Wong, Nathan D, 2441, 3164, 3682, 3683, 3838
 Wong, Philip, 445
 Wong, Pierre C, 2083
 Wong, S C, 2518, 3066
 Wong, S. C, 1923, 2461
 Wong, S. Chiu, 3015, 3716
 Wong, Shing Chiu, 1702
 Wong, Tony, 845
 Woo, Jean L, 3765
 Woo, Kam S, 3765
 Woo, Mary A, 3422
 Woo, Y. J, 1886, 1993
 Wood, Katherine C, 667
 Wood, Mark A, 1656, 1896, 3242
 Woodard, Pamela K, 1909
 Woodard, Sarah, 5
 Woodiwiss, Angela, 2262
 Woodrum, Julie E, 1489
 Woods, Christopher, 2640
 Woods, David, 111
 Woodward, Mark, 3784
 Woolson, Robert F, 2372
 Word, Ruth A, 701
 Worldwide WAVE Study Investigators, for the, 1650
 the working group of cardiac rehabili-
 on behalf of, Error In Author-
 Please Check3823
 Worrall-Carter, Linda, 3417
 Worrall-Carter, Linda, 3420