

(12) United States Patent Daly et al.

(54) VEGF-BINDING FUSION PROTEINS AND NUCLEIC ACIDS ENCODING THE SAME

(75) Inventors: **Thomas J. Daly**, New City, NY (US);

Nicholas J. Papadopoulos,

LaGrangeville, NY (US); Margaret Karow, Putnam Valley, NY (US)

(73) Assignee: Regeneron Pharmaceuticals, Inc.,

Tarrytown, NY (US)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 297 days.

(21) Appl. No.: 11/204,709

(22)Filed: Aug. 16, 2005

(65)**Prior Publication Data**

> US 2006/0058234 A1 Mar. 16, 2006

Related U.S. Application Data

- (60) Continuation-in-part of application No. 11/016,097, filed on Dec. 17, 2004, and a continuation-in-part of application No. 11/016,503, filed on Dec. 17, 2004, which is a division of application No. 10/009,852, filed as application No. PCT/US00/14142 on May 23, 2000, now Pat. No. 7,070,959, application No. 11/204,709, and a continuation-in-part of application No. 10/880, 021, filed on Jun. 29, 2004, now Pat. No. 7,279,159, which is a continuation-in-part of application No. 10/609,775, filed on Jun. 30, 2003, now Pat. No. 7,087, 411, which is a continuation-in-part of application No. 10/009,852, filed on Dec. 6, 2001, now Pat. No. 7,070, 959.
- Provisional application No. 60/138,133, filed on Jun. 8, 1999.
- (51) Int. Cl. C12N 15/62

(2006.01)C12N 15/63 (2006.01)

US 7,396,664 B2 (10) **Patent No.:** Jul. 8, 2008 (45) Date of Patent:

A61K 38/18 (2006.01)C07K 14/00 (2006.01)C07K 14/71 (2006.01)

(52) **U.S. Cl.** **435/69.7**; 424/134.1; 424/192.1; 435/71.1; 435/252.3; 435/320.1; 435/325; 435/358; 514/2; 514/12; 530/350; 530/387.3;

536/23.4

(58) Field of Classification Search None See application file for complete search history.

(56)References Cited

U.S. PATENT DOCUMENTS

6,897,294 B2 5/2005 Davis-Smyth et al.

FOREIGN PATENT DOCUMENTS

WO WO 00/75319 12/2000

OTHER PUBLICATIONS

Davis-Smyth et al. (1996). The second immunoglobulin-like domain of the VEGF tyrosine kinase receptor Flt-1 determines ligand binding and may initiate a signal transduction cascade. The EMBO Journal. 15(18):4919-4927.*

Davis-Smyth, T., et al, (1998) J. Bio. Chem. 273(6):3216-3222. Heidaran, M., et al., (1990) J. Bio. Chem. 265(31):18741-18744. Holash, J., et al., (2002) PNAS 99(17):11393-11398. Heidaran, M., et al., (1995) FASEB J. 9:140-145. Wulff, C., et al., (2002) Endocrinology 143(7):2797-2807.

* cited by examiner

Primary Examiner—Christine J Saoud Assistant Examiner-Jon M Lockard (74) Attorney, Agent, or Firm—Valeta Gregg, Esq.

(57)ABSTRACT

Nucleic acid molecules and multimeric proteins capable of binding vascular endothelial growth factor (VEGF). VEGF traps are disclosed which are therapeutically useful for treating VEGF-associated conditions and diseases, and are specifically designed for local administration to specific organs, tissues, and/or cells.

11 Claims, 3 Drawing Sheets



Observed	Observed Masses	Lypeoted	Monogonopaide	Dronoced Sugar
VGT C04003 M500	VGT C04008 M500	Masses	Compositions	Chain Structures
1356.4	1356.0	1.356.1	(Man)2(Man)3(Glc)2	
1380.9	1381.6	1381.2	(Glc)(Man)3(Glc)2(Fuc)	T
1397.2	1397.2	1397.3	(Gal)(Glc)(Man)3(Glc)2	
1438.6	1438.2	1438.2	(Glc)2(Man)3(Glc)2	
1526.7	1525.9	1527.2	(Glc)(Man)3(Glc)2(Fuc)2	
1543.4	1542.8	1543.4	(Gal)(Glc)(Man)3(Glc)2(Fuc)	
1559.7	1559.4	1559.4	(Gal)(Glc)(Man)(Man)3(Glc)2	
1584.6	1584.0	1584.4	(Glc)2(Man)3(Glc)2(Fuc)	
1600.5	0.0091	1600.4	(Gal)(Glc)2(Man)3(Glc)2	
1646.5	1646.6			Adduct of 1584.4
1688.5	1688.0	1688.4	(SA)(Gal)(Glc)(Man)3(Glc)2	*
1721.5	1720.8	1721.5	(Gal)(Glc)(Man)2(Man)3(Glc)2	

Figure 1A

Observed	Observed Masses	Lycotop	Monocochanide	Proposed Sugar
VGT C04003 M500	VGT C04008 M500	Masses	Compositions	Chain Structures
1746.7	1746.2	1746.5	(Gal)(Glc)2(Man)3(Glc)2(Fuc)	
1762.8	1762.6	1762.1	(Gal)2(Glc)2(Man)3(Glc)2	1
1809.5	1809.1			Adduct of 1746.5
1834.8	1834.7	1834.5	(SA)(Gal)(Glc)(Man)3(Glc)2(Fuc)	**************************************
1849.9	0.1881	1850.5	(SA)(Gal)2(Glc)(Man)3(Glc)2	
6.1681	1.1981	9.1681	(SA)(Gal)(Glc)2(Man)3(Glc)2	* • • • • • • • • • • • • • • • • • • •
9.8061	1908.2	9.8061	(Gal)(Glc)2(Man)3(Glc)2(Fuc)	
1996.9	2:9661	9.9661	(\$A)(Gal)2(Glc)(Man)3(Glc)2(Fuc)	
2012.8	2012.3	2012.8	(SA)(Gal) (Glc)2(Man)2(Man)3(Glc)2	
2037.7	2037.3	2037.7	(SA)(Gal)(Glc)2(Man)3(Glc)2(Fuc)	* • • • • • • • • • • • • • • • • • • •
2053.8	2053.2	2053.8	(SA)(Gal)2(Glc)2(Man)3(Glc)2	***
2075.8	2074.9			Adduct of 2053.8
2116.0	2115.7			Adduct of 2053.8

igure 1B



Observed Masses	d Masses	Fynected	Monosaccharide	Proposed Sugar
VGT C04003 M500	VGT C04008 M500	Masses	Compositions	Chain Structures
2200.0	2199.3	2199.9	(SA)(Gal)2(Glc)2(Man)3(Glc)2(Fuc)	*
2221.5	2221.5			Na ⁺ Adduct of 2199.9
2262.5	2261.3			Adduct of 2199.9
2345.6	2345.6	2345.0	(SA)2(Gal)2(Glc)2(Man)3(Glc)2	
2407.0	2407.2			Adduct of 2345.0
2492.5	2492.0	2491.2	(SA)2(Gal)2(Glc)2(Man)3(Glc)2(Fuc)	
2554.3	2553.1			Adduct of 2491.2
2564.6	2564.0	2565.2	(SA)(Gal)3(Glc)3(Man)3(Glc)2(Fuc)	
2857.2	2857.3	2856.5	(SA)2(Gal)3(Glc)3(Man)3(Glc)2(Fuc)	
2918.1	2918.9			Adduct of 2856.5
3148.5	3148.5	3147.8	(SA)3(Gal)3(Glc)3(Man)3(Glc)2(Fuc)	
3211.0	3211.0			Adduct of 3147.8

Figure 1C

1

VEGF-BINDING FUSION PROTEINS AND NUCLEIC ACIDS ENCODING THE SAME

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of application Ser. Nos. 11/016,097 filed 17 Dec. 2004 and 11/016,503 filed 17 Dec. 2004, which are divisionals of Ser. No. 10/009,852 filed 6 Dec. 2001, now U.S. Pat No. 7,070,959, which is the 10 National Stage of International Application No. PCT/US00/ 14142 filed 23 May 2000, which claims the benefit under 35 USC § 119(e) of U.S. provisional application No. 60/138,133 filed 8 Jun. 1999; and Application Ser. No. 10/880,021 filed 29 Jun. 2004, now U.S. Pat. No. 7,279,159 which is a continuation-in-part of Ser. No. 10/609,775 filed 30 Jun. 2003 now U.S. Pat. No. 7,087,411, which is a continuation-in-part of Ser. No. 10/009,852 filed 6 Dec. 2001, now U.S. Pat. No. 7,070,959, which is the National Stage of International Application No. PCT/US00/14142 filed 23 May 2000, which 20 claims the benefit under 35 USC § 119(e) of U.S. provisional Application No. 60/138,133 filed 8 Jun. 1999, which applications are herein specifically incorporated by reference in their entireties.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The invention encompasses fusion polypeptides capable of binding vascular endothelial cell growth factor (VEGF), VEGF family members, and splice variants with specifically desirable characteristics, as well as therapeutic methods of use.

2. Brief Summary of the Invention

In a first aspect, the invention features an isolated nucleic acid molecule encoding a fusion polypeptide comprising receptor components R1-R2-F, wherein R1 is vascular endothelial cell growth factor (VEGF) receptor component Ig domain 2 of Flt-1 (Flt1D2), R2 is VEGF receptor component Ig domain 3 of Flk-1 (Flk1D3) (also known as KDR), and F is a fusion component.

In a related second aspect, the invention features a VEGF-binding fusion polypeptide comprising VEGF receptor components R1-R2-F, wherein R1, R2, and F are as defined above. The components may be connected directly to each other or connected via one or more spacer sequences. In a preferred embodiment, R1 and R2 are the only receptor components present. In a specific embodiment, the VEGF-binding fusion polypeptide is amino acids 27-129 (R1) and 130-231 (R2) of SEQ ID NO:8, or a variant thereof.

The fusion component F is selected from the group consisting of a multimerizing component, a serum protein, or a molecule capable of binding a serum protein. In a preferred embodiment, F is a multimerizing component capable of 55 interacting with a multimerizing component on another fusion polypeptide to form a multimeric structure, e.g., a dimer or trimer. Most preferably, the F is selected from the group consisting of (i) a multimerizing component comprising a cleavable region (C-region), (ii) a truncated multimer- 60 izing component, (iii) an amino acid sequence between 1 to about 200 amino acids in length having at least one cysteine residue, (iv) a leucine zipper, (v) a helix loop motif and (vi) a coil-coil motif. Preferably, the multimerizing component is an immunoglobulin domain. In one embodiment, F is a full- 65 length or truncated immunoglobulin domain consisting of amino acids 232-458, 232-457, or 352-458 of SEQ ID NO:8.

2

The receptor components may be arranged in different orders, for example, R1R2F; R2R1F; R1FR2; R2FR1; FR1R2; FR2R1, etc. The components of the fusion polypeptide may be connected directly to each other, or connected via a spacer sequence.

In a third aspect, the invention features a multimeric VEGF-binding protein, comprising two or more fusion polypeptides of the invention (also called a VEGF "trap"). A VEGF trap composed of two fusion polypeptides having at least one truncated multimerizing component is termed a "truncated mini-trap." The multimeric VEGF-binding protein of the invention is capable of binding VEGF with an affinity (Kd) of at least 1×10⁻¹⁰ M, preferably at least 1×10⁻¹¹ M, even more preferably at least 1×10⁻¹² M, as measured by BIACORE™ based assays.

The C-region may be created in the multimerizing component by insertion, deletion, or mutation, such that an enzymatically or chemically cleavable site is created. The C-region may be created in any multimerizing component and at any position within; preferably, the C-region is created in a full-length Fc domain, or a fragment thereof, or a C_{H3} domain. The C-region may be a site cleavable by an enzyme, such as, thrombin, ficin, pepsin, matrilysin, or prolidase or cleavable chemically by, for example, formic acid or CuCl₂.

In all embodiments of the VEGF-binding fusion polypeptides of the invention (including full length VEGF-binding fusion polypeptides, truncated VEGF-binding fusion polypeptides, etc.), a signal sequence (S) may be included at the beginning (or N-terminus) of the fusion polypeptide of the invention. The signal sequence may be native to the cell, recombinant, or synthetic.

The components of the fusion polypeptide may be connected directly to each other or be connected via spacers. In specific embodiments, one or more receptor and/or fusion partner components of the fusion polypeptide are connected directly to each other without spacers. In other embodiments, one or more receptor and/or fusion partner components are connected with spacers.

In a fourth aspect, the invention encompasses vectors comprising the nucleic acid molecules of the invention, including expression vectors comprising the nucleic acid molecule operatively linked to an expression control sequence. In a fifth aspect, the invention encompasses host-vector systems for the production of a fusion polypeptide which comprise the expression vector, in a suitable host cell; host-vector systems wherein the suitable host cell is a bacterial, yeast, insect, mammalian cell; an E. Coli cell, or a COS or CHO cell. Additional encompassed in a sixth aspect are VEGF-binding fusion polypeptides of the invention modified by acetylation or pegylation, and other post-translational modifications resulting from expression in a mammalian cell line. Methods for acetylating or pegylating a protein are well known in the art. In specific embodiments, the fusion polypeptide of the invention expressed in a mammalian cell line such as a CHO cell comprises amino acids 27-457 of SEQ ID NO:8 and is glycosylated at Asn residues 62, 94, 149, 222 and 308.

In a related seventh aspect, the invention features a method of producing a VEGF-binding fusion polypeptides of the invention, comprising culturing a host cell transfected with a vector comprising a nucleic acid sequence of the invention, under conditions suitable for expression of the protein from the host cell, and recovering the fusion polypeptides so produced.

The VEGF-binding fusion polypeptides of the invention are therapeutically useful for treating any disease or condition which is improved, ameliorated, or inhibited by removal, inhibition, or reduction of VEGF. A non-exhaustive list of



DOCKET

Explore Litigation Insights



Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

