

# RETINA

THE JOURNAL OF RETINAL AND VITREOUS DISEASES

5 for Impact Factor  
Worldwide  
ing Ophthalmology Journals

**COMPARISON OF TWO DOSES OF INTRAVITREAL BEVACIZUMAB AS PRIMARY TREATMENT FOR MACULAR EDEMA SECONDARY TO BRANCH RETINAL VEIN OCCLUSIONS: RESULTS OF THE PAN AMERICAN COLLABORATIVE RETINA STUDY GROUP AT 24 MONTHS**

*Wu, Arevalo, Berrocal, Maia, Roca, Morales-Cantón, Alezzandrini, Díaz-Llopis*

**VASCULAR EVENTS IN PATIENTS WITH AGE-RELATED MACULAR DEGENERATION TREATED WITH INTRAOCULAR BEVACIZUMAB**

*Sheybani, Kymes, Schlieff, Apte*

**A PROSPECTIVE STUDY OF BLOOD PRESSURE AND INTRAOCULAR PRESSURE CHANGES IN HYPERTENSIVE AND NONHYPERTENSIVE PATIENTS AFTER INTRAVITREAL BEVACIZUMAB INJECTION**

*Lee, Yang, Lim, Lew*

**INTRAVITREAL BEVACIZUMAB FOR CHOROIDAL NEOVASCULARIZATION SECONDARY TO PRESUMED OCULAR HISTOPLASMOSIS SYNDROME**

*Ehrlich, Ciulla, Maturi, Kheradiya, Hrisomalos, Shulman, Guess, Coyle, Harris*

**"TREAT AND EXTEND" DOSING OF INTRAVITREAL ANTIVASCULAR ENDOTHELIAL GROWTH FACTOR THERAPY FOR TYPE 3 NEOVASCULARIZATION/RETINAL ANGIOMATOUS PROLIFERATION**

*Engelbert, Zweifel, Freund*

**PHACOEMULSIFICATION WITH INTRAVITREAL BEVACIZUMAB INJECTION IN DIABETIC PATIENTS WITH MACULAR EDEMA AND CATARACT**

*Akinci, Batman, Ozkiloglu, Altinsoy*

**THE COURSE OF RESPONSE TO FOCAL/GRID PHOTOCOAGULATION FOR DIABETIC MACULAR EDEMA**

*The Diabetic Retinopathy Clinical Research Network*

**A 1-YEAR RETROSPECTIVE REVIEW OF RANIBIZUMAB FOR NAÏVE NONSUBFOVEAL CHOROIDAL NEOVASCULARIZATION SECONDARY TO AGE-RELATED MACULAR DEGENERATION**

*Arias, Ruiz-Moreno, Gómez-Ulla, Fernández, Montero*

**PHOTODYNAMIC EFFECTS ON RETINAL OXYGEN SATURATION, BLOOD FLOW, AND ELECTROPHYSIOLOGICAL FUNCTION IN PATIENTS WITH NEOVASCULAR AGE-RELATED MACULAR DEGENERATION**

*Tsuchihashi, Mori, Peyman, Shimada, Yoneya*

**INTRAOPERATIVE USE OF HANDHELD SPECTRAL DOMAIN OPTICAL COHERENCE TOMOGRAPHY IMAGING IN MACULAR SURGERY**

*Dayani, Maldonado, Farsiu, Toth*

**ENHANCED DEPTH IMAGING OPTICAL COHERENCE TOMOGRAPHY OF THE CHOROID IN CENTRAL SEROUS CHORIORETINOPATHY**

*Imamura, Fujiwara, Margolis, Spaide*

**RETINAL DETACHMENT AFTER PRIMARY RHYTHOXYMYOPIA**

*Mudvari, Ravage, Rezaei*

 **Wolters Kluwer | Lippincott Williams**  
Health



PROPERTY OF THE  
**NATIONAL  
LIBRARY OF  
MEDICINE**

Full Text  
**OID**

ISSN 0275-004X

This material was prepared at the NLM and may be Subject US Copyright Laws

# RETINA®

THE JOURNAL OF RETINAL AND VITREOUS DISEASES

EDITOR-IN-CHIEF

ALEXANDER J. BRUCKER  
Philadelphia, PA

ASSOCIATE EDITORS

HARRY W. FLYNN, JR.  
Miami, FL  
WILLIAM R. FREEMAN  
La Jolla, CA

PHILIP J. ROSENFELD  
Miami, FL  
CAROL L. SHIELDS  
Philadelphia, PA  
RICHARD F. SPAIDE  
New York, NY

CHARLES P. WILKINSON  
Baltimore, MD  
GEORGE A. WILLIAMS  
Royal Oak, MI

Abstract Editor

JOSE S. PULIDO  
Rochester, MN

SECTION EDITORS

Photo Essay Editor

LEE M. JAMPOL  
Chicago, IL

Diagnostic and Therapeutic Editor

H. RICHARD McDONALD  
San Francisco, CA

Surgical Technique Editor

GEORGE A. WILLIAMS  
Royal Oak, MI

EDITORIAL BOARD

GARY W. ABRAMS  
Detroit, MI  
LLOYD P. AIELLO  
Boston, MA  
SOPHIE J. BAKRI  
Rochester, MN  
MARK S. BLUMENKRANZ  
Stanford, CA  
NEIL M. BRESSLER  
Baltimore, MD  
SUSAN B. BRESSLER  
Baltimore, MD  
STANLEY CHANG  
New York, NY  
STEVEN T. CHARLES  
Memphis, TN  
EMILY Y. CHEW  
Bethesda, MD  
D. JACKSON COLEMAN  
New York, NY  
GABRIEL J. COSCAS  
Paris, France  
DONALD J. D'AMICO  
Boston, MA  
FREDERICK L. FERRIS, III  
Bethesda, MD  
HOWARD F. FINE  
New York, NY  
GERALD A. FISHMAN  
West Chester, IL

JAMES C. FOLK  
Iowa City, IA  
J. DONALD M. GASS\*  
Nashville, TN  
\*1928-2005  
KURT GITTER  
New Orleans, LA  
W. RICHARD GREEN  
Baltimore, MD  
JULIA A. HALLER  
Baltimore, MD  
FRANK G. HOLZ  
Bonn, Germany  
DOUGLAS A. JABS  
New York, NY  
GLENN J. JAFFE  
Durham, NC  
MARK W. JOHNSON  
Ann Arbor, MI  
PETER K. KAISER  
Cleveland, OH  
ANSELM KAMP'IK  
Munich, Germany  
BARUCH D. KUPPERMANN  
Irvine, CA  
HARVEY A. LINCOFF  
New York, NY  
BROOKS W. McCUEN  
Durham, NC

RONALD G. MICHELS\*  
Baltimore, MD  
\*1943-1991  
DAVID C. MUSCH  
Ann Arbor, MI  
DAVID H. ORTH  
Chicago, IL  
ARNALL PATZ  
Baltimore, MD  
GHOLAM A. PEYMAN  
Tucson, AZ  
INGRID SCOTT  
Miami, FL  
JERRY A. SHIELDS  
Philadelphia, PA  
LAWRENCE J. SINGERMAN  
Cleveland, OH  
JASON S. SLAKTER  
New York, NY  
PAUL STERNBERG, JR.  
Nashville, TN  
YASUO TANO\*  
Osaka, Japan  
\*1948-2009  
JOHN T. THOMPSON  
Baltimore, MD  
CYNTHIA A. TOTH  
Durham, NC  
LAWRENCE A. YANNUZZI  
New York, NY

MANAGING EDITOR: TERRY ROTHSTEIN BRUCKER

ASSOCIATE MANAGING EDITOR: GENE SEABOLT

RETINA® The Journal of Retinal and Vitreous Diseases is indexed in *Biological Abstracts*, *EMBASE/Excerpta Medica*, *Index Medicus*, and *Current Contents*. RETINA® The Journal of Retinal and Vitreous Diseases (ISSN 0275-004X) is published ten times a year for the Ophthalmic Communications Society, Inc., by Lippincott Williams & Wilkins, 16522 Hunters Green Parkway, Hagerstown, MD 21740-2116. Business offices are located at 530 Walnut Street, Philadelphia, PA 19106-3621. Production offices are located at 351 W. Camden Street, Baltimore, MD 21201-2436. Periodicals postage paid at Hagerstown, MD and at additional mailing offices. Copyright © 2009 by Ophthalmic Communications Society, Inc.

Address for subscription information, orders, or changes of address (except Japan, India, Bangladesh, Sri Lanka, Nepal, and Pakistan): 16522 Hunters Green Parkway, Hagerstown, MD 21740-2116; phone: 1-800-638-3030; fax: 301-223-2400; in Maryland, call collect, 301-223-2300. In Japan, contact LWJ Igaku-Shoin Ltd., 3-23-14 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan; phone: 81-3-5689-5400; fax: 81-3-5689-5402. In India, Bangladesh, Sri Lanka, Nepal, and Pakistan, contact Globe Publication Pvt. Ltd. B-13, 3rd Fl. A Block, Shopping Complex, Naraina Vihar, Ring Road, New Delhi 110028, India; phone: 91-11-579-3211; fax: 91-11-579-8876.

Annual subscription rates worldwide: \$368.00 Individual Domestic, \$494.00 Individual International, \$834.00 Institutional Domestic, \$920.00 Institutional International, \$177.00 Residents Domestic, \$177.00 Residents International. All prices include a handling charge. United States residents of AL, CO, DC, FL, GA, HI, IA, ID, IN, KS, KY, LA, MD, MO, ND, NM, NV, PR, RI, SC, SD, UT, VT, WA, WV add state sales tax. (The Canadian GST tax of 7% will be added to the subscription price of all orders shipped to Canada. Lippincott Williams & Wilkins' GST Identification Number is 895524239. Publications Mail Agreement #864927.) Subscriptions outside the United States must be prepaid. Subscriptions outside North America must add \$15.00 for airfreight delivery. Single copies, when available, may be ordered from the publisher. Single copies \$92.00. Prices subject to change without notice. Copies will be replaced without charge if the publisher receives a request within 90 days of the mailing date, both in the U.S. and worldwide. Visit us on-line at [www.lww.com](http://www.lww.com). Web site: [www.retinajournal.com](http://www.retinajournal.com)

Individual and in-training subscription rates include print and access to the online version. Institutional rates are for print only; online subscriptions are available via Ovid. Institutions can choose to purchase a print and online subscription together for a discounted rate. Institutions that wish to purchase a print subscription, please contact Lippincott Williams & Wilkins, 16522 Hunters Green Parkway, Hagerstown, MD 21740-2116; phone 800-638-3030 (outside the United States 301-223-2300); fax 301-223-2400. Institutions that wish to purchase an online subscription or online with print, please contact the Ovid Regional Sales Office near you or visit [www.ovid.com/site/index.jsp](http://www.ovid.com/site/index.jsp) and select Contact and Locations.

POSTMASTER: Send address changes to RETINA® The Journal of Retinal and Vitreous Diseases, P.O. Box 1550, Hagerstown, MD 21740.

Advertising inquiries: Bob Williams, National Account Manager, Neurology & Ophthalmology, Lippincott Williams & Wilkins, 530 Walnut Street, Philadelphia, PA 19106; telephone: 215-521-8394; fax: 215-521-8411; e-mail: [bob.williams@wolterskluwer.com](mailto:bob.williams@wolterskluwer.com) This material was copied

Ⓢ Text printed on acid-free paper.

at the NLM and may be  
Subject US Copyright Laws



# “TREAT AND EXTEND” DOSING OF INTRAVITREAL ANTIVASCULAR ENDOTHELIAL GROWTH FACTOR THERAPY FOR TYPE 3 NEOVASCULARIZATION/RETINAL ANGIOMATOUS PROLIFERATION

MICHAEL ENGELBERT, MD, PhD,\*† SANDRINE A. ZWEIFEL, MD,\*†‡  
K. BAILEY FREUND, MD\*†

---

**Purpose:** The purpose of this study was to analyze long-term outcomes for the treatment of type 3 neovascularization/retinal angiomatous proliferation using a “Treat and Extend” dosing regimen for anti-vascular endothelial growth factor therapy.

**Methods:** This was a retrospective analysis of visual acuity and optical coherence tomography data of 11 eyes of 10 consecutive patients with newly diagnosed type 3 neovascularization/retinal angiomatous proliferation treated with intravitreal bevacizumab and/or ranibizumab with at least a 12-month follow-up. Three monthly injections were followed by continued treatment at intervals increasing by 2 weeks per visit, to a maximum of 10 weeks, unless clinical or optical coherence tomography evidence of persistent or recurrent fluid was present, in which case, the interval was shortened.

**Results:** Mean baseline Snellen visual acuity was 20/80, improved to 20/40 at 1 month, and was maintained throughout the 36-month period ( $n = 11$  at 12 months,  $n = 10$  at 24 months, and  $n = 8$  at 36 months) ( $P < 0.04$ , paired  $t$ -test). The mean center point optical coherence tomography thickness decreased from 320  $\mu\text{m}$  to 180–230  $\mu\text{m}$ , and was maintained during the study period ( $P < 0.02$ ). The mean number of injections was seven in the first year, six in the second year, and seven in the third year.

**Conclusion:** “Treat and Extend” anti-vascular endothelial growth factor dosing in type 3 neovascularization/retinal angiomatous proliferation delivers promising outcomes at a reduced burden for the patient and health care system compared with monthly and optical coherence tomography-guided dosing regimens.

RETINA 29:1424–1431, 2009

---

Type 3 neovascularization (otherwise known as retinal angiomatous proliferation [RAP])<sup>1,2</sup> is a subtype of neovascular age-related macular degeneration (AMD) with distinct angiographic and optical coherence tomography (OCT) features related to intraretinal proliferation of the abnormal vessels with associated retinal–retinal and retinal–choroidal anastomosis. Its natural course is

typically worse than other more frequent lesion types such as subretinal pigment epithelium neovascularization (type 1)/occult choroidal neovascularization or subneurosensory neovascularization (type 2)/well-defined (classic) choroidal neovascularization.<sup>3–5</sup> Many different treatment strategies<sup>6</sup> such as photocoagulation,<sup>3,4,7</sup> transpupillary thermotherapy,<sup>3,8</sup> photodynamic therapy (PDT),<sup>9–13</sup> intravitreal anti-vascular endothelial growth factor (anti-VEGF) agents,<sup>6,14,15</sup> intravitreal triamcinolone acetonide, surgical excision,<sup>16,17</sup> and many combinations of the above<sup>18–21</sup> have been tried in small case series with limited follow-up.

Monthly injections of antiangiogenic agents have become the standard of care for the treatment of neovas-

---

From the \*LuEsther T. Mertz Retinal Research Center, Manhattan Eye, Ear and Throat Hospital; †Vitreous-Retina-Macula Consultants of New York, New York, New York; and ‡University Hospital Zurich, Department of Ophthalmology, Frauenklinikstrasse 24, 8032 Zurich, Switzerland.

Supported by The Macula Foundation Inc.

Reprint requests: K. Bailey Freund, MD, 460 Park Avenue, 5th Floor, NY 10022; e-mail: kbfn@aol.com

cular AMD<sup>22,23</sup> but are expensive and difficult to sustain in this elderly patient population. However, the less frequent dosing in the PIER trial,<sup>24</sup> in which patients received quarterly injections after an initial series of three monthly injections, could not reproduce the excellent results obtained in trials using monthly dosing. The PrONTO Study<sup>25,26</sup> attempted to tailor the dosing to the individual needs of the patient based on acuity decline, clinical findings, or OCT evidence of disease activity and was able to demonstrate good visual results after a >24-month period.

Although PrONTO-style dosing has become widely adopted in the retinal community and seems to yield favorable results, this strategy does require monthly visits, clinical examinations, and OCTs with patients uncertain if or when they will need treatment. Because eyes with type 3 neovascularization/RAP typically manifest retinal–choroidal anastomosis, recurrent exudation may occur earlier and more frequently than with other neovascular lesion types. In our experience, some patients managed with this strategy will return for assessments having already developed macular hemorrhages in the injection-free interval with irreversible vision loss.<sup>27–29</sup> In theory, a dosing regimen that does not maintain the macula in a “dry” state could deny some patients the opportunity for further visual recovery.

The “Treat and Extend” dosing regimen is a strategy intended to resolve macular exudation and maintain the macula in this “dry” state indefinitely with, when possible, fewer patient visits and treatments than monthly dosing.<sup>30,31</sup> The strategy consists of an initial induction or “loading” sequence of at least three monthly injections. If stable visual acuity, an absence of macular hemorrhage, and a dry OCT have been achieved at this point, patients continue to receive regular maintenance injections at increasing intervals. At 6 weeks after the last of the three monthly injections, visual acuity, clinical findings, and OCT changes are recorded again, and patients receive an injection regardless of the presence or absence of disease activity. However, the interval to the next visit (and scheduled injection) is based on an observed change in the above parameters. If there are no changes, the next visit is scheduled for 8 weeks later. If there is a change, the patient comes for another scheduled injection and examination after 4 weeks. The observation and scheduled treatment interval is extended (hence the term “Treat and Extend”) to a maximum of 10 weeks. We report on 11 eyes of 10 patients with type 3 neovascularization/RAP managed with the “Treat and Extend” dosing regimen and with follow-up of between 12 and 36 months.

### Materials and Methods

A waiver of authorization for use of protected health information for the above-referenced research

and a waiver of consent for this retrospective chart review were obtained from the Institutional Review Board committee of the Manhattan Eye Ear and Throat Hospital, New York, NY.

The diagnosis of type 3 neovascularization/RAP was made by the treating physician (K.B.F.) based on the characteristic clinical, OCT, and angiographic features including intraretinal hemorrhage, cystoid macular edema, intraretinal vascular anastomosis, retinal–choroidal anastomosis, and in some cases, the presence of pigment epithelial detachment (PED) on OCT. Patients treated previously with thermal laser, PDT, or intravitreal pegaptanib (Macugen, Pfizer Inc., New York, NY), or who presented with subfoveal fibrosis or atrophy, a history of vitrectomy, aphakia or absence of posterior capsule, history of idiopathic or autoimmune associated uveitis in either eye, or diabetic retinopathy more severe than mild nonproliferative stage, were excluded from this study. Patients with preexisting cardiac or cerebrovascular conditions were not excluded from the study.

The treatment consisted of intravitreal injection of 1.25 mg of bevacizumab (Avastin, Genentech Inc., South San Francisco, CA) or 0.5 mg ranibizumab (Lucentis, Genentech Inc.) suspended in 0.05 mL. For the purpose of this analysis, no distinction between either antiangiogenic drug was made. Before intravitreal injection, topical anesthesia and surface disinfection with 5% povidone–iodine was performed. Intravitreal injections were administered at the time of diagnosis and subsequently followed a protocol we termed “Treat and Extend.” Patients all received at least 3 monthly injections followed by continued treatment at intervals increasing by 2 weeks per visit once visual acuity was stable, OCT showed an absence of intra- and subretinal fluid, and all hemorrhage had resolved. Resolution of PED was not required before treatment intervals were lengthened. The treatment interval was extended to a maximum of 10-week “maintenance” unless clinical examination or OCT detected new hemorrhage or persistent/recurrent fluid. In those cases, the interval was shortened by 2 weeks and maintained at that duration, provided this resolved the fluid.

The main outcome measure in this study was visual acuity after treatment. Decrease in retinal thickness, number of injections needed, and change in funduscopy or tomographic appearance were assessed as well.

Snellen visual acuity was measured by a certified ophthalmic technician. Snellen acuity was converted into logarithm of the minimum angle of resolution (logMAR) for statistical analysis at baseline and at 1, 3, 12, 24, and 36 months after injection of an antiangiogenic agent. Changes in logMAR-converted acuities were tested with a paired Student’s *t*-test and accepted as significant if  $P < 0.05$ . Also, the proportions of patients

approximately halving ( $\geq 0.3$  logMAR, but  $< 0.6$  logMAR-converted Snellen visual acuity improvement) or approximately quartering their visual angle ( $\geq 0.6$  logMAR-converted Snellen visual acuity improvement), as well as those that remained stable ( $< 0.3$  logMAR-converted visual acuity improvement) or lost lines on the Snellen chart compared with baseline, were reported.

The quantitative assessments of center point retinal thickness were made using Stratus OCT (Carl Zeiss Meditec, Dublin, CA) and Topcon OCT (Topcon 3D OCT-1000, Topcon Medical Systems, Paramus, NJ). The center point retinal thickness was defined as the distance between the internal limiting membrane and the retinal pigment epithelium under the fovea and did not include any fluid under the retinal pigment epithelium. For Topcon OCT images, the calipers provided by the Topcon image analysis software were used. The Stratus OCT measurements were made manually on the IMAGENet software on a single horizontal line scan through the fovea (Topcon Medical Systems), and the calculated data in pixels were multiplied with a conversion factor of 8  $\mu\text{m}$ /pixel. This conversion factor had been derived from previous comparisons of controls on the different imaging platforms (based on 20 normal eyes measured on the 2 platforms, Howard F. Fine, personal communication).

The qualitative assessment included identification of intraretinal cysts, neovascular complex within the retinal layers, and PED. Additional fundoscopic and tomographic changes and their development over time were recorded as well. Specifically, the presence of intraretinal hemorrhage or development of a pigment epithelial rip on funduscopy and the presence of intra- or subretinal fluid or PED on high-resolution B-scans were determined. Because staging of type 3 neovascularization/RAP is difficult and of controversial significance, this was not performed.

**Results**

Eleven eyes of 10 patients were included in this study. Eleven eyes completed the 12-month follow-up, 10 eyes completed the 24-month follow-up, and 8 eyes completed the 36-month follow-up.

Patient demographics, baseline, and follow-up visual acuity, center point retinal thickness data, and number of injections in the first, second, and third year are presented in Table 1. Median patient age was 85 years (range, 71–92 years). Seven of 10 patients were women. Two contralateral eyes had evidence of antecedent type 3 neovascularization/RAP lesions, and 2 patients developed disease in the contralateral eye during the study period. Only one of these two latter eyes was treated with a “Treat and Extend” protocol and included

Table 1. Summary of Patient Data

Patient	Gender	Age (Years)	Baseline			First Year			Second Year			Third Year		
			VA	OCT ( $\mu\text{m}$ )	Month 1 VA	Month 1 OCT ( $\mu\text{m}$ )	Month 12 VA	Month 12 OCT ( $\mu\text{m}$ )	Month 24 VA	Month 24 OCT ( $\mu\text{m}$ )	Month 36 VA	Month 36 OCT ( $\mu\text{m}$ )	No Inj	Year
1	OD	F	20/400	257	20/160	299	20/100	20/200	174	20/80	181	5	6	P
2	OS	F	20/50	241	20/50	183	20/60	20/40	177	20/40	163	5	7	P
3	OD	F	20/25	246	20/25	192	20/25	20/25	186	—	—	10	—	P
3	OS	—	20/200	433	20/40	289	20/25	20/30	108	20/40	136	8	10	L
4	OS	F	20/80	282	20/30	199	20/20	20/25	174	—	—	5	—	—
5	OS	M	20/200	410	20/70	174	20/100	—	232	—	—	—	—	L
6	OS	M	20/20	182	20/25	133	20/25	20/20	174	20/20	134	8	6	—
7	OS	F	20/50	548	20/30	374	20/50	20/30	196	20/30	199	5	7	—
8	OD	F	20/30	200	20/40	224	20/40	20/25	224	20/30	225	6	6	—
9	OS	M	20/200	388	20/70	206	20/40	20/40	224	20/40	218	6	5	—
10	OS	F	20/70	336	20/40	246	20/50	20/60	175	20/50	197	7	7	—

Inj, injections; P, at presentation; L, later.

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.