

VEGF Trap induces antiglioma effect at different stages of disease

Candelaria Gomez-Manzano, Jocelyn Holash, Juan Fueyo, Jing Xu, Charles A. Conrad, Kenneth D. Aldape, John F. de Groot, B. Nebiyu Bekele, and W. K. Alfred Yung

Departments of Neuro-Oncology (C.G.-M., J.F., J.X., C.A.C., J.F.G., W.K.A.Y.), Pathology (K.D.A.), and Biostatistics and Applied Mathematics (B.N.B.), The University of Texas M. D. Anderson Cancer Center, Houston, TX; Regeneron Pharmaceuticals, Inc., Tarrytown, NY (J.H.); USA

Pathological angiogenesis is a hallmark of cancer, specifically of glioblastomas, the most malignant and common primary brain tumor. Vascular endothelial growth factor (VEGF) is the key protein in the regulation of the hyper-vascular phenotype of primary malignant brain tumors. In this study, we tested VEGF Trap, a soluble decoy receptor for VEGF, in an intracranial glioma model. VEGF Trap was administered in short or prolonged schedules to animals bearing human gliomas at different stages of disease. Of importance, VEGF Trap treatment was efficacious in both initial and advanced phases of tumor development by significantly increasing overall survival. Furthermore, this effect was enhanced in animals treated with more prolonged regimens. In addition, we observed the emergence of a VEGF Trap-resistant phenotype characterized by tumor growth and increased invasiveness. Our results suggest that VEGF Trap will be effective in treating both patients with recurrent or progressive resectable glioblastoma and patients that have undergone extensive initial surgery. Finally, our results indicate that the clinical success of VEGF Trap may depend on a prolonged treatment in combined therapy aiming to simultaneously inhibit angiogenesis and tumor invasion. *Neuro-Oncology* 10, 940–945, 2008 (Posted to *Neuro-Oncology* [serial online], Doc. D08-00085, August 14, 2008. URL <http://neuro-oncology.dukejournals.org>; DOI: 10.1215/15228517-2008-061)

Received April 7, 2008; accepted June 4, 2008.

Address correspondence to C. Gomez-Manzano, Department of Neuro-Oncology, Unit 1002, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., Houston, TX 77030, USA (cmanzano@mdanderson.org).

Keywords: glioblastoma, therapy, VEGF, VEGF Trap

The striking induction of angiogenesis in glioblastoma multiforme (GBM) has fueled the speculation that progression to GBM requires the activation of angiogenesis, a finding that has stimulated significant efforts to develop angiogenesis-blocking agents. Vascular endothelial growth factor (VEGF) is critical for promoting the earliest stages of vasculogenesis, which includes endothelial cell proliferation, differentiation, migration, and tubular formation. Clinical trials of specific VEGF inhibitors for the treatment of patients with gliomas are ongoing, and preliminary analyses showed beneficial effects in patients with malignant gliomas.^{1–4} Recently, a new anti-VEGF agent, VEGF Trap/afibercept (henceforth referred to as VEGF Trap), has been developed by incorporating domains of both VEGF receptor 1 (VEGFR-1) and VEGFR-2 fused to the constant region of human immunoglobulin G1, which acts as a soluble decoy receptor for VEGF. VEGF Trap has very high affinity for all isoforms of VEGF-A (<1 pM), as well as placental growth factor, a closely related angiogenic factor.⁵ VEGF Trap was engineered to have minimal interactions with the extracellular matrix, and this property apparently accounts for its satisfying pharmacokinetic profile superior to soluble forms of VEGFR-1.⁵ Its efficacy has been proven in preclinical studies in several types of solid tumor^{5–9} and in a subcutaneous glioma model.¹⁰ Because tumor progression and angiogenesis are greatly dependent on the existent micro-environment of the tumor,^{11,12} we undertook this study to characterize the effect of VEGF Trap in an orthotopic glioblastoma model in several stages of the disease. We

Copyright 2008 by the Society for Neuro-Oncology

have previously described the development of growth patterns and angiogenesis in an intracranial U-87 MG human glioma model. Vessel cooption and remodeling were present at the early stages of disease, whereas the advanced stages are distinguished by high vascular density.¹³ These two phases were similar to stages described in other previous reports.^{14,15} Based on this tumoral angiogenesis and kinetic pattern, we administered VEGF Trap to animals bearing U-87 MG intracranial xenografts at several phases of tumor development. In the present study, we demonstrated that VEGF Trap treatment in animals bearing human gliomas resulted in significant prolonged survival. Of importance, our results indicate that VEGF Trap was equally effective against initial or advanced disease, and that the response was enhanced when VEGF Trap was administered in a prolonged schedule.

Material and Methods

Cell Line

The human glioma cell line U-87 MG was purchased from the American Type Culture Collection (Manassas, VA, USA). Cells were maintained in Dulbecco's modified Eagle/F12 medium (1:1, vol:vol) (The University of Texas M. D. Anderson Cancer Center Media Core Facility, Houston, TX, USA) supplemented with 10% fetal calf serum and 1% antibiotic/antimycotic agent (Invitrogen, Carlsbad, CA, USA) in a humidified atmosphere containing 5% CO₂ at 37°C.

Drugs

VEGF Trap and human Fc (hFc, constant region of human IgG1) were kindly provided by Regeneron Pharmaceuticals (Tarrytown, NY, USA). Stocks of 50 mg/ml in aqueous solution were kept at -80°C.

In Vivo Experiments

The U-87 MG human glioma cells (5×10^5) were engrafted in the caudate nucleus of athymic mice (Harlan Sprague Dawley Inc., Indianapolis, IN, USA), as previously described.¹³ At 0, 4, and 10 days after cell implantation, we administered VEGF Trap (2.5 mg/kg subcutaneously, twice a week, for a total of 3 or 6 weeks) to separate groups of 10–15 animals per treatment bearing U-87 MG intracranial xenografts. Either phosphate-buffered saline (PBS) or hFc was blindly administered as a control agent in randomly selected subgroups of glioma-bearing animals. Animals showing generalized or local symptoms of disease were euthanized. Brains were fixed in 4% formaldehyde for 24 h and embedded in paraffin. Slides were stained with hematoxylin and eosin. All animal studies were performed in the veterinary facilities of The M. D. Anderson Cancer Center in accordance with institutional guidelines.

Enzyme-Linked Immunosorbent Assays

Blood was collected from the tail vein of glioma-bearing mice 3 days after the initial dose of VEGF Trap, hFc, or vehicle, and VEGF Trap was quantified in the serum by enzyme-linked immunosorbent assays (ELISA), as previously reported.¹⁶

Statistical Analyses

The *in vivo* anticancer effect of different treatments was assessed by plotting Kaplan-Meier survival curves, and treatment groups were compared using the log-rank test. The effects of VEGF Trap when administered in different treatment schedules were analyzed using a permutation test.

Results and Discussion

Antiglioma Effect of VEGF Trap on Initial Disease

The VEGF Trap-mediated anti-glioma effect was assessed *in vivo* using an intracranial human glioma xenograft model. We selected the U-87 MG cell line for this study because it produces gliomas in nude mice with highly predictable growth kinetics and well-characterized pathological features¹³; in addition, U-87 MG cells express high levels of VEGF and, when implanted intracranially in immunocompromised mice, develop as highly vascularized tumors.^{11,13} Our group has previously characterized the kinetics of tumor growth and vascularization of human U-87 MG xenografts implanted intracranially in nude mice. Of interest to the present study, U-87 MG intracranial tumors exhibited initially minimal tumor growth, but changes in the host vessels were evident as soon as day 1 and definitely by day 4 after implantation; these changes included significant vessel co-option, as illustrated by the existence of engorged smooth-muscle actin (SMA)-positive vascular structures in the periphery of the xenograft.¹³

To test the effect of VEGF Trap in the initial phases of the disease, we planned two different treatment schedules (Figs. 1 and 2) consisting of the subcutaneous administration of 2.5 mg/kg VEGF Trap twice weekly over 3 weeks, starting on either day 0 (schedule A) or day 4 (schedule B) after the intracranial implantation of human glioma cells in nude mice. Control groups were treated with PBS or hFc at doses and volumes similar to those used for the test drug. The agents were administered in a double-blinded manner; that is, the identity of the test groups was concealed from both the personnel preparing the drugs and the animal caretakers.

Animals treated with VEGF Trap starting on day 0 or day 4 after implantation had significantly prolonged survival compared to the hFc- or PBS-treated animals ($p < 0.0001$ and $p < 0.005$, respectively). In animals treated with schedule A, the median overall survival of the control-treated animals (treated with either hFc or PBS) was 30 days, with all animals dying by day 33. Treatment with VEGF Trap prolonged the mean survival




Fig. 1. Schematic representation of the treatment schedule used with the anti-vascular endothelial growth factor (VEGF) agent VEGF Trap, which was based on our previous studies of the kinetics of growth and vascularization in the U-87 MG intracranial model. U-87 MG cells were implanted in the brains of the animals on day 0, and VEGF Trap was administered starting on day 0 (schedule A), day 4 (schedule B), or day 10 (schedules C₅ and C₆) after cell implantation. The two schedule C subgroups were treated in either a 3-week (schedule C₅) or 6-week (schedule C₆) schedule. Schedules A and B followed a 3-week treatment regimen. Animals were euthanized when signs of neurological or generalized disease appeared.

by 8 days. In animals treated with schedule B, the mean survival in the PBS- and hFc-treated animals was 27.5 and 30 days, respectively, but it was increased to 36 days in the group treated with VEGF Trap. No treatment-schedule-dependent differences in survival duration were observed in animals receiving VEGF Trap, suggesting VEGF Trap is efficacious in initial phases of disease that

were characterized by active vessel co-option and remodeling. Analysis performed 3 days after the first VEGF Trap doses were administered revealed high VEGF Trap levels (approximately >50 µg/ml) in the serum of all these animals, suggesting an efficient systemic biodistribution (data not shown).




Fig. 2. Effect of the anti-vascular endothelial growth factor (VEGF) agent VEGF Trap on initial phases of disease: survival analysis of glioma-bearing animals treated with VEGF Trap since day 0 (A) or day 4 (B), as pictured in Fig. 1. Kaplan-Meier survival curves begin on the day of U-87 MG intracranial implantation following the subcutaneous injection of VEGF Trap or of vehicle or human Fc (control). The *p*-values (determined by log-rank test) show significant overall survival differences between VEGF Trap-treated and control-treated animals. Abbreviations: E, events; N, number of animals.

Antiglioma Effect of VEGF Trap on Disease Burden

To test the effect of VEGF Trap on tumor burden, and based on our previous study of U-87 MG intracranial growth and angiogenesis, we decided to start treatment on day 10 after cell implantation in one subgroup of mice (Fig. 1, schedule C₃). According to our previous studies, by day 10, increased microvascular density (MVD) was associated with exponential tumor growth and a decrease in the rate of induced angiogenesis within the host and the tumor periphery.¹³ Twelve days after implantation, the tumors consisted of spherical masses of cells with a high MVD and large, distorted, SMA-positive vessels. The tumor limits were clearly defined, and the cancer cells did not exhibit the invasive pattern into host tissue seen in preceding days.¹³

In the present study, glioma cells were implanted intracranially, and 10 days later, VEGF Trap was administered subcutaneously at a dose of 25 mg/kg twice weekly for 3 weeks. Control groups were treated with PBS or hFc at doses and volumes similar to those of the test drug. Treatment of the glioma-bearing animals with VEGF Trap resulted in a significant increase in the survival of these animals ($p < 0.005$) (Fig. 3A). In particular, the median overall survival of control-treated (PBS or hFc) animals was 31 days, with all the animals dead by day 33, whereas the mean survival of VEGF Trap-treated animals was 45 days. We observed no significant difference in the effect of VEGF Trap on prolonging survival at different stages of the disease (comparing effects of schedules A and B with schedule C₃) ($p > 0.1$, permutation test), suggesting that VEGF Trap can be similarly effective in both the initial and burden disease stage. These data further suggest that targeting circulating levels of VEGF is equally effective in challenging tumor

growth under both initial and established tumoral vasculature phases.

Antiglioma Effect of Prolonged VEGF Trap Treatment

We next explored the effect in vivo of more prolonged VEGF Trap treatment. In this experiment, animals bearing intracranial human gliomas were treated with VEGF Trap (25 mg/kg) twice weekly for 6 weeks starting on day 10 after cell implantation (Fig. 1, schedule C₆). Control animals were treated with vehicle or hFc (25 mg/kg) twice weekly until they showed signs of disease, at which time they were euthanized according to institutional regulations. Animals treated with VEGF Trap for 6 weeks survived longer than did animals treated with hFc (median overall survival, 55 days and 21 days, respectively; Fig. 3B) ($p < 0.0001$). We also analyzed the difference in median survival times between the animals treated with VEGF Trap for 6 weeks and those treated for 3 weeks. Using the permutation test and after adjusting for overall survival on PBS-treated groups, we found the increase in survival obtained with the 6-week VEGF Trap treatment to be significantly greater than the increase in survival obtained with the 3-week treatment ($p < 0.05$). These data suggest that VEGF Trap is more effective in prolonging overall survival when administered in a prolonged treatment schedule.

Histological Examination of VEGF Trap-Treated Tumors

Microscopic analysis of histological sections from formalin-fixed, paraffin-embedded brains revealed that control- and VEGF Trap-treated animals eventually suffered from the lethal growth of their tumors. Because of




Fig. 3. Effect of the anti-vascular endothelial growth factor (VEGF) agent VEGF Trap on advanced glioma disease: survival analyses of glioma-bearing animals that were treated with VEGF Trap starting on day 10 after cell implantation in either a 3-week (schedule C₃) or 6-week (schedule C₆) regimen, as pictured in Fig. 1. Kaplan-Meier survival curves begin on the day of U-87 MG intracranial implantation following the subcutaneous injection of VEGF Trap or control agent (vehicle or human Fc). The p -values (determined by log-rank test) show significant overall survival differences between VEGF Trap-treated and control-treated animals. Abbreviations: E, events; N, number of animals.




Fig. 4. Histological examination of brain sections from animals treated with the anti-vascular endothelial growth factor (VEGF) agent VEGF Trap. (A) Hematoxylin-eosin staining of mouse brains bearing U-87 MG xenografts treated with human Fc (hFc) or VEGF Trap according to schedule B. No signs of hemorrhagic areas or an enhanced invasive phenotype were observed after VEGF Trap treatment. N, normal tissue; T, tumor tissue. Original magnification, $\times 100$. (B) Histological examination of brain sections from animals treated with VEGF Trap as described for schedule C. Sections stained with hematoxylin and eosin show the presence of an invasive phenotype with satellitosis characterized by glioma clustering around vascular vessels and accumulation of invasive glioma cells far from the main tumor mass (arrows). Original magnification: left, $\times 100$; right, $\times 200$.

previous studies describing that treatment with antiangiogenic agents may result in intracranial hemorrhages or enhance tumor invasion,^{2,17} we specifically examined the tumors for the presence of these adverse effects. Histological examination of the brains of the cohorts treated for 3 weeks did not reveal either phenomenon. Treated U-87 MG-derived tumors displayed a very well-defined border with the normal host parenchyma (Fig. 4A). However, examination of the brains of animals that received prolonged treatment (6 weeks) of VEGF Trap, which survived longer than those treated on a 3-week schedule, revealed the signs of mass effect and the presence of the so-called “secondary structures” or “satellitosis” consisting of aggregations of glioma cells in the perivascular regions, as well as the presence of glioma cells along the Virchow-Robin spaces (Fig. 4B). These data suggest that U-87 MG-derived xenografts acquired an invasive phenotype in response to anti-VEGF therapy. These results are in agreement with a similar pattern of

growth of intracranial G55 xenografts in animals treated with an antibody against mouse VEGFR-2, DC101,¹⁷ or a neutralizing VEGF antibody.¹⁸ These results may be likewise in agreement with those from clinical trials in patients with cancer treated with VEGF inhibitors, in that they survived longer but eventually exhibited resistance to the treatment.^{19,20} Of importance, the model described here offers us the possibility of testing combined therapies designed to counteract the emergence of a resistant phenotype to anti-VEGF therapies.

Taken together, our data show that treatment with VEGF Trap significantly prolonged the survival of glioma xenograft-bearing mice. Of great interest, initial/residual disease and disease burden were both similarly affected by the antiangiogenesis treatment. In addition, the prolonged use of VEGF Trap (over 6 weeks) improved outcomes significantly more than did treatment administered in a short schedule (over 3 weeks).

The traits for personalized medicine are emerging

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.