

Original article

# Pulse Oximetry in the Diagnosis of Acute Heart Failure

Josep Masip,\* Maria Gayà, Joaquim Páez, Antoni Betbesé, Francisco Vecilla, Ruben Manresa, and Pilar Ruíz

Unidad de Cuidados Intensivos, Hospital Sant Joan Despí Moisès Broggi, Consorci Sanitari Integral, Universidad de Barcelona, Sant Joan Despí, Barcelona, Spain

Article history:

Received 12 February 2012  
Accepted 18 February 2012  
Available online 4 July 2012

Keywords:

Pulse oximetry  
Oxygen saturation  
Acute heart failure  
Myocardial infarction

ABSTRACT

**Introduction and objectives:** Oxygen saturation by pulse oximetry is commonly used for monitoring critical patients, but its utility as a diagnostic marker of acute heart failure has not been assessed. This study analyzed the diagnostic role of oxygen saturation by pulse oximetry in a series of patients with acute myocardial infarction.

**Methods:** In a prospective observational cohort study of 220 consecutive patients with acute myocardial infarction, data collection included baseline oxygen saturation by pulse oximetry (without oxygen), physiologic measurements, Killip class and data from portable chest radiography, recorded at the same hour on each of the first three days after admission. Patients were followed up for one year.

**Results:** There were 612 assessments. Baseline oxygen saturation by pulse oximetry decreased progressively in relation to the presence and the severity of acute heart failure assessed by Killip classes 1 to 3 (mean: 95, 92 and 85, respectively;  $P < .001$ ) or by radiology score 0 to 4 (95, 94, 92, 89 and 83, respectively;  $P < .001$ ), with a correlation coefficient of 0.66 and 0.63, respectively. Receiver operating characteristic curves disclosed the cut-off of oxygen saturation by pulse oximetry  $< 93$  to have the greatest area, with a sensitivity of 65%, specificity 90%, and overall test accuracy 83%. Patients grouped according to lowest oxygen saturation by pulse oximetry showed significantly different rates of one-year mortality or rehospitalization for heart failure.

**Conclusions:** Baseline oxygen saturation by pulse oximetry is useful in establishing the diagnosis and severity of heart failure in acute settings such as myocardial infarction and may have prognostic implications. The diagnosis may be suspected when baseline oxygen saturation by pulse oximetry is  $< 93$ .

© 2012 Sociedad Española de Cardiología. Published by Elsevier España, S.L. All rights reserved.

## Pulsioximetría en el diagnóstico de insuficiencia cardiaca aguda

RESUMEN

**Introducción y objetivos:** La saturación de oxígeno mediante pulsioximetría se usa habitualmente en la monitorización de pacientes críticos, pero su utilidad como marcador diagnóstico de insuficiencia cardiaca aguda no ha sido evaluada. Este estudio analiza el papel diagnóstico de la saturación de oxígeno mediante pulsioximetría en una serie de pacientes con infarto agudo de miocardio.

**Métodos:** En un estudio observacional prospectivo, se incluyó a 220 pacientes consecutivos con infarto agudo de miocardio. Se registraron la saturación de oxígeno mediante pulsioximetría basal (sin oxígeno), las constantes fisiológicas, la clase Killip y la puntuación radiológica a la misma hora, durante los primeros 3 días del ingreso. Se siguió a los pacientes durante 1 año.

**Resultados:** Se obtuvieron 612 valoraciones. La saturación de oxígeno mediante pulsioximetría basal disminuyó de forma progresiva respecto a la presencia y la gravedad de la insuficiencia cardiaca, tanto valorada con la clasificación de Killip 1-3 (medias, 95, 92 y 85, respectivamente;  $p < 0,001$ ), como con la puntuación radiológica 0-4 (95, 94, 92, 89 y 83, respectivamente;  $p < 0,001$ ), con un cociente de correlación de 0,66 y 0,63 respectivamente. Las curvas *receiver operating characteristic* para la saturación de oxígeno mediante pulsioximetría mostraron que el punto de corte  $< 93$  tenía la mayor área, con sensibilidad del 65%, especificidad del 90% y precisión diagnóstica del 83%. Los pacientes agrupados según su saturación de oxígeno mediante pulsioximetría más baja, mostraron tasas significativamente distintas de mortalidad o rehospitalización con insuficiencia cardiaca.

**Conclusiones:** La saturación de oxígeno mediante pulsioximetría es útil para establecer el diagnóstico y la gravedad de la insuficiencia cardiaca en situaciones agudas como el infarto de miocardio y puede tener implicaciones pronósticas. El diagnóstico debe sospecharse cuando la saturación de oxígeno mediante pulsioximetría basal es  $< 93$ .

© 2012 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Palabras clave:

Pulsioximetría  
Saturación de oxígeno  
Insuficiencia cardiaca aguda  
Infarto agudo de miocardio

SEE RELATED ARTICLE:

<http://dx.doi.org/10.1016/j.rec.2012.05.002>, Rev Esp Cardiol. 2012;65:867–8

\* Corresponding author: Unidad de Cuidados Intensivos, Hospital Sant Joan Despí Moisès Broggi, Consorci Sanitari Integral, Universidad de Barcelona, Jacint Verdaguer 90, 08970 Sant Joan Despí, Barcelona, Spain.

E-mail address: [jmasip@ub.edu](mailto:jmasip@ub.edu) (J. Masip).

## Abbreviations

AHF: acute heart failure  
 AMI: acute myocardial infarction  
 LB-SpO<sub>2</sub>: lowest baseline SpO<sub>2</sub>  
 MB-SpO<sub>2</sub>: morning baseline SpO<sub>2</sub>  
 SpO<sub>2</sub>: oxygen saturation by pulse oximetry

## INTRODUCTION

Acute heart failure (AHF) is a frequent complication in patients with acute myocardial infarction (AMI).<sup>1,2</sup> The incidence may be rather variable depending on the diagnostic criteria.<sup>3</sup> In clinical practice, diagnosis is initially made on admission by anamnesis and physical examination.<sup>4</sup> However, identification of this complication is frequently difficult and it is necessary to obtain complementary data from portable chest radiography<sup>5</sup> or other techniques such as echocardiography,<sup>6</sup> pulmonary wedge catheter,<sup>7</sup> or biological markers.<sup>8</sup> Because AHF affects pulmonary function,<sup>9</sup> gas exchange could be altered even in mild forms. Although AHF has been associated with hypoxemia,<sup>10</sup> the diagnostic contribution of oxygen saturation by bedside pulse oximetry (SpO<sub>2</sub>) has not been analyzed. Most of the previous studies using this technique in AMI were limited to investigating the incidence of hypoxemia or nocturnal sleep apnea syndrome.<sup>11–15</sup> Therefore, we conducted an observational study to assess the value of baseline SpO<sub>2</sub> for the diagnosis and severity of AHF in patients with AMI, a homogeneous population with a high prevalence of this complication.

## METHODS

### Patient Population

Our institutional ethics committee approved the study design and all patients gave their informed consent. All consecutive patients of any age with ST and non-ST segment elevation AMI admitted to our critical/semi-critical intensive care unit were considered for inclusion. AMI was defined by prolonged chest pain with ischemic electrocardiographic changes and elevated myocardial necrosis biomarkers. Patients were included irrespective of the treatment received prior to admission (primary coronary angioplasty, fibrinolytic or conventional treatment). Patients with moderate to severe chronic obstructive pulmonary disease (GOLD classes II–III) and those who developed cardiogenic shock were excluded.

### Study Design

This was a prospective observational cohort study. Patients were continuously monitored by pulse oximetry during the first three days, using modular Hewlett-Packard (SpO<sub>2</sub>/Plet M1020A) pulse oximeters with a central station monitor. The target SpO<sub>2</sub> was “morning baseline SpO<sub>2</sub>” (MB-SpO<sub>2</sub>), which was registered by nurses in awake patients on the morning round, usually after chest radiography, and taking the most stable value on room air.

steady value was reached, after several minutes of oxygen discontinuation.

Physical examination data was recorded by staff physicians (Killip class)<sup>4</sup> on admission and daily morning rounds thereafter. Portable chest radiographs were also obtained daily during rounds and were evaluated using a radiology score by consensus of 2 observers and sometimes 3 in case of initial disagreement. This score was based on Battler et al. classification,<sup>5</sup> establishing 5 categories: (0) *normal*; (1) *vascular redistribution* to upper lung fields; (2) *interstitial edema* defined by B-Kerley lines or prominent and diffuse hilar contour with vascular redistribution; (3) *localized edema* with alveolar infiltrates in less than 50% of the lung fields, mainly hilo-basal, and (4) *diffuse alveolar edema* with alveolar infiltrates in 3 or 4 quadrants.

Data obtained on the first morning after admission was introduced as “first day”. Thus, the interval from the onset of AMI to that moment could be variable. However, data on the following days were also collected in the morning but at constant intervals up to 3 days. Oxygen therapy was administered if baseline SpO<sub>2</sub> was lower than 93%. Cardiac necrosis markers (MB fraction creatine kinase/troponin T) were obtained on admission and every 6 h, until peak value was identified. Natriuretic peptides were not available in our center at the time of study recruitment.

The variables registered during daily rounds were heart rate, respiratory rate, blood pressure, MB-SpO<sub>2</sub>, Killip class, and radiology score. The variable “lowest baseline SpO<sub>2</sub>” (LB-SpO<sub>2</sub>) was obtained retrospectively upon discharge from the intensive care unit (ICU), taking the lowest SpO<sub>2</sub> value among those recorded during the whole acute cardiac care stay.

Ventricular function was assessed by echocardiography, which was performed during hospitalization. After discharge, patients were followed up for one year and death or rehospitalization with AHF was registered.

### Diagnosis of Acute Heart Failure

Patients with Killip classes 2 or 3 were initially considered to have AHF in the Bayesian analysis. However, an abnormal chest radiography was required to confirm this diagnosis in such a way that patients who presented normal radiology (score 0) were considered not to have AHF (Killip false positive) regardless of the assigned Killip class. Conversely, patients initially evaluated as Killip 1 on physical examination who presented unequivocal signs of pulmonary congestion on radiology (score 3–4) were considered to have AHF (Killip false negative).

### Statistical Analysis

The total data registered during the 3 days of monitoring accounted for more than 600 determinations for every one of the 6 study variables (Killip class, radiology score, MB-SpO<sub>2</sub>, heart rate, respiratory rate and blood pressure). Receiver operating characteristic (ROC) curves from these simultaneous determinations were used to analyze the test accuracy for the diagnosis of AHF for different cut-off levels of MB-SpO<sub>2</sub>, heart, and respiratory rates.

When patients were analyzed, their LB-SpO<sub>2</sub> was individually assigned. Comparisons between means were analyzed using the Student *t* test or analysis of variance (ANOVA). In variables with no normal distribution or no homogeneous variances, a nonparametric test (Kruskal-Wallis or Mann-Whitney) was used. Pair-wise multiple comparisons were performed by least-significant differences or Tamhane test when variances were not equal. Non-continuous variables were analyzed by Fisher exact test. Continuous variables were reported as mean (standard deviation).

Independent factors were analyzed by logistic regression analysis. Kaplan-Meier event-free (death or rehospitalization with AHF) survival analysis was used for the follow-up evaluation, assigning patients to one of three groups, according to their LB-SpO<sub>2</sub>. The range of LB-SpO<sub>2</sub> for these groups was guided by the 95% confidence intervals (95%CI) of the mean SpO<sub>2</sub> in each Killip class group. Consequently, these groups were: group 1 (SpO<sub>2</sub>>93%); group 2 (SpO<sub>2</sub>=90%-93%), and group 3 (SpO<sub>2</sub><90%).

## RESULTS

Over a 2-year period, 220 consecutive patients with AMI, with and without ST segment elevation, were admitted to our intensive/semi-intensive care unit and accepted for participation in the study. The study was approved by our Ethics and Research Committee and was carried out in the year 2000. Fourteen patients (6%) developed cardiogenic shock or needed mechanical ventilation (invasive or noninvasive) and were subsequently excluded. In the remaining patients there were 122 with ST-segment elevation and 9 with left bundle branch block. Non-Q wave AMI was the final diagnosis in 106 patients.

The characteristics of the patients based on the incidence of AHF are shown in Table 1. Overall, patients with AHF were older, less predominantly men, with higher rates of previous hypertension, diabetes, myocardial infarction and heart failure than patients without AHF.

After 3 days the cumulative incidence of some degree of AHF was 35%. Two thirds of the patients presented AHF on admission whereas the rest of the patients developed this complication thereafter. Regarding symptoms, AHF was clinically silent in the majority of the patients and less than 20% showed some degree of shortness of breath. Mortality after 30 days and at 1 year was significantly greater in patients with AHF.

Physiologic measurements are also presented in Table 1. Patients with AHF showed significantly higher respiratory and

heart rates and lower LB-SpO<sub>2</sub> and left ventricular ejection fraction, while there were no differences in blood pressure.

Distribution of mean MB-SpO<sub>2</sub> values in relation to the degree of AHF is shown in Figures 1 and 2. Median MB-SpO<sub>2</sub> was significantly lower according to the degree of AHF measured either by Killip classes (n=612) or Battler scores (n=604). All intergroup pair-wise comparisons were also significantly different. There were also significant differences between respiratory rates for Killip classes 1 to 3 (19 [3], 21 [4], and 26 [8], respectively; *P*<.001) and for Battler scores 0 to 4 (19 [3], 20 [3], 21 [5], 24 [7], and 26 [7], respectively; *P*<.001). All intergroup pair-wise comparisons for respiratory rate were also significantly different.

## Diagnostic Accuracy

Regarding measurements (n=612), the correlation coefficients between MB-SpO<sub>2</sub> and Killip class and Battler score were -0.666 and -0.619, respectively (*P*<.001). Regarding patients, the correlation coefficient between LB-SpO<sub>2</sub> and left ventricular ejection fraction was 0.433; *P*<0.001.


The MB-SpO<sub>2</sub> measurements were plotted in a ROC curve for the diagnosis of AHF (Fig. 3). The cut-off level of SpO<sub>2</sub><93 showed the greatest area under the curve with the highest test accuracy (83%). The cut-off value for patients with chronic bronchitis (n=45) with greater area was SpO<sub>2</sub><93 as well. The ROC curves were also plotted for respiratory and heart rate. The cut-off with greater area for respiratory rate was >21 breaths/min (area, 0.64), with a test accuracy of 59% (odds ratio=2.19; 95%CI, 1.53-3.16). The cut-off value for heart rate was >76 bpm (area, 0.61) and the test accuracy 73% (odds ratio=4.21; 95%CI, 2.8-6.3).

There were 29 patients who were misclassified by Killip score, 20 (10%) false negatives and 9 (4%) false positives. The combination of MB-SpO<sub>2</sub><93 and Killip classes (≥2) or respiratory rate (>21 breaths/min) increased the specificity to 99.4% and 97% and the positive predictive value to 97.2% and 78%, respectively.


**Table 1**  
Characteristics of 206 Patients With Acute Myocardial Infarction According to the Presentation of Acute Heart Failure

Characteristics	No AHF (n = 133)	AHF (n = 73)	P
<i>Characteristics</i>			
Age, years	61.5±12.4	70±9.4	<.001
Male	105 (79)	47 (64)	.031
Diabetes	34 (26)	35 (48)	.002
Hypertension	59 (45)	48 (69)	.002
Prior myocardial infarction	19 (15)	28 (39)	<.001
Prior heart failure	6 (5)	22 (31)	<.001
LBBB	1 (0.8)	8 (11)	.001
ST-Segment elevation	84 (63)	48 (66)	.760
<i>Outcomes</i>			
30-day mortality	0	9 (12.3)	<.001
One-year mortality	6 (4.9)	18 (26.1)	<.001
<i>Physiologic measurements</i>			
Lowest SpO <sub>2</sub>	93.4±2.1	87.9±6.3	<.001
Systolic blood pressure, mmHg	118.6±14.6	118.9±17.5	.880
Heart rate, bpm	71.7±11.7	81.1±13.5	<.001
Respiratory rate, breaths/min	18.9±2.4	21.5±3.8	<.001
LVEF, %	60.1±8.7	45.8±10.5	<.001
End-diastolic diameter, mm	52.3±5.7	54.7±5.9	.010

AHF, acute heart failure; LBBB, left bundle branch block; LVEF, left ventricle ejection fraction; SpO<sub>2</sub>, oxygen saturation by pulse oximetry. Data are expressed as mean ± standard deviation or no. (%).



**Figure 1.** Baseline oxygen saturation by pulse oximetry values in relation to the degree of heart failure assessed by physical examination (Killip classes) for all measurements. Squares are means; bars (standard deviation). SpO<sub>2</sub>, oxygen saturation by pulse oximetry.



**Figure 2.** Baseline oxygen saturation by pulse oximetry values in relation to the degree of heart failure assessed by radiology score (Battler classes) for all measurements. Squares are means; bars (standard deviation). SpO<sub>2</sub>, oxygen saturation by pulse oximetry.


**Follow-up**

Mean follow-up was 13 months and was completed in 192 patients (93%). Multivariate logistic regression analysis showed age and AHF to be the independent predictive variables for 1-year death or rehospitalization for AHF (Table 2). LB-SpO<sub>2</sub> was only an independent factor when AHF was not introduced in the analysis.

Kaplan-Meier event-free (mortality and rehospitalization for AHF) survival curves for the 3 LB-SpO<sub>2</sub> groups are shown in Figure 4. There were significant differences in the event-free rates between the 3 groups as well as in pair-wise group comparisons: group 1 vs group 2 (90.7% vs 76.1%; P=.01), group 2 vs group 3 (76.1 vs 54.3%; P=.01) and group 1 vs group 3 (90.7 vs 54.3%; log rank test, P=.00005).

**DISCUSSION**

This study shows the utility of baseline SpO<sub>2</sub> as a complementary tool to establish the diagnosis and severity of AHF. We chose to perform the study in patients with AMI because of the high prevalence of AHF in this population. The main advantage of SpO<sub>2</sub> lies in the fact that is noninvasive, it can be monitored continuously, and it is not affected by interobserver or intraobserver variability. The finding of a baseline SpO<sub>2</sub> lower than 93 may be considered a signal of AHF and a warning to clinicians about this complication which may be especially useful between rounds. The lower the SpO<sub>2</sub>



SpO <sub>2</sub> , %	Sensitivity	Specificity	PPV	NPV	Accuracy	AUC-ROC	OR (95% CI)
<90	32	98.2	87.1	79.1	79.9	0.65	26 (12-54)
<91	42.6	97.3	85.7	81.6	82.1	0.69	27 (14-50)
<92	52.1	95.4	81.5	83.7	83.3	0.73	23 (13-39)
<93	64.5	90.1	71.7	86.7	83.2	0.77	17 (11-26)
<94	75.7	78.9	57.9	89.5	78	0.76	12 (8-18)
<95	85.8	63.9	47.5	92.2	69.9	0.74	11 (7-17)

**Figure 3.** Receiving operating characteristics curve for oxygen saturation by pulse oximetry and acute heart failure. A table with information about different cut-off points is attached. 95%CI, 95% confidence interval; AUC, area under curve; NPV, negative predictive value; OR, odds ratio; ROC: receiver operating characteristic; SpO<sub>2</sub>, oxygen saturation by pulse oximetry; PPV, positive predictive value.

The results also stress the role of measuring respiratory rate and heart rate. These findings are relevant because respiratory rate is not usually considered a diagnostic marker of AHF. Although heart rate is commonly assessed for the diagnosis, it can be altered by the use of beta-blockers. The association of a decreasing SpO<sub>2</sub> with an increasing respiratory rate may reinforce the suspicion of AHF in patients with doubtful diagnostic criteria.

Our study found good correlations between baseline SpO<sub>2</sub> and the degree of AHF or left ventricular ejection fraction. These findings were in accordance with previous trials analyzing oxygen saturation by CO-oximetry from arterial blood samples, which showed a close correlation between the defects in gas exchange and the severity of AHF or the pulmonary capillary wedge pressures,<sup>10,16-18</sup> although this correlation decreased in chronic patients.<sup>19</sup> Hypoxemia in AHF is explained by an increase in veno-arterial shunting and ventilation-perfusion imbalance.<sup>10,16-18</sup> Despite this well-known interaction, however, the determination of oxygen saturation (SaO<sub>2</sub>) by arterial puncture as a diagnostic marker of AHF is not a common practice and this valuable information is frequently missed. Although SpO<sub>2</sub> values are closely related to SaO<sub>2</sub> obtained by CO-oximetry, the accuracy of pulse oximetry may decrease in special situations such as severe hypoxemia, anemia, acidosis, pigmentation, low perfusion states, and the use of vasoactive drugs.<sup>20,21</sup> Therefore, SpO<sub>2</sub> and SaO<sub>2</sub> are not totally equivalent and the information in the literature about the role of SpO<sub>2</sub> in AHF is scarce. Some authors have reported their experience with SpO<sub>2</sub> in detecting Cheyne-Stokes respiration in patients with heart failure<sup>22</sup> but, to our knowledge, no previous

**Table 2**  
Risk Factors for One-year Death or Rehospitalization for Heart Failure

	Univariate analysis		Multivariate analysis	
	OR (95%CI)	P	OR (95%CI)	P
Acute heart failure	7.8 (3.6-17.1)	<.001	7.1 (2.7-19)	<.001
<b>Measurements</b>				
SpO <sub>2</sub> <sup>a</sup>	5.5 (2.6-11.8)	<.001		
Left ventricular function <sup>b</sup>	4.3 (1.9-9.6)	.001		
Heart rate <sup>b</sup>	2.5 (1.2-5)	.013		
Respiratory rate <sup>b</sup>	2.5 (1.2-5)	.02		
<b>Clinical characteristics</b>				
History of heart failure	5.5 (2.3-12.9)	<.001		
Prior myocardial infarction	3.5 (1.7-7.4)	.001		
Diabetes	2.5 (1.2-5)	.015		
Hypertension	2.2 (1.05-4.6)	.049		
Age <sup>c</sup>	1.13 (1.07-1.18)	<.001	1.12 (1.05-1.18)	<.001

95%CI, 95% confidence interval; OR, odds ratio; SpO<sub>2</sub>, oxygen saturation by pulse oximetry.

The OR reflects the odds for patients with the characteristic in question, as compared to those without the characteristic.


<sup>a</sup> SpO<sub>2</sub> is a categorical variable (cut-off <93) obtained by taking the lowest value registered on the rounds for each patient.

<sup>b</sup> Categorical variables (cut-off: ejection fraction <0.50; heart rate >76 bpm; respiratory rate >21 breaths/min).

<sup>c</sup> The OR for age represents the exponent for each year of age in the logistic equation.

The diagnosis of AHF may be difficult in patients with AMI because it is often silent.<sup>23</sup> In our series, the majority of patients with AHF did not show shortness of breath. Therefore, because AHF may be misdiagnosed in many cases by physical examination<sup>24,25</sup> (5%-10% of patients in our study were misclassified), the diagnostic criteria of AHF in this study required abnormal radiology because this technique enhances the sensitivity and diagnostic accuracy.<sup>3,23</sup> Chest radiography is considered a class I recommendation in the European Society of Cardiology guidelines on the diagnosis and treatment of AHF.<sup>26,27</sup> In these guidelines, SpO<sub>2</sub> also received the same consideration for monitoring, not for diagnosis.

In recent decades the evaluation of ventricular function and filling pressures by echocardiography has become a widespread practice in the assessment of AHF, often replacing invasive approaches like pulmonary catheter.<sup>27-29</sup> Nevertheless, the information provided by echocardiography is not continuous. Biochemical markers such as natriuretic peptides are frequently abnormally elevated on admission<sup>30</sup> and have better prognostic than diagnostic value in patients with AMI.<sup>8</sup> In this context, SpO<sub>2</sub> may become a complementary, continuous, and low-cost parameter for assessing AHF with prognostic implications, as shown in the follow-up.



Patients at risk at the end of period

SpO <sub>2</sub> >93%	86	80	71
SpO <sub>2</sub> 90-93%	71	56	51
SpO <sub>2</sub> <90%	35	20	16

**Figure 4.** Kaplan-Meier event-free survival curves for three groups of oxygen saturation by pulse oximetry. Events are death or rehospitalization with heart failure within the first year. Numbers are patients at risk in each group after 1,

## Limitations

This study had some limitations, including the lack of determination of natriuretic peptides, which were not commonly used in our institution when the study was performed, and the absence of a gold standard for the diagnosis of AHF in patients with AMI. Other limitations were the registry format that introduced data from each patient several times, the inability to achieve a completely blinded clinical assessment, and finally, the fact that patients with AHF were older and age might produce a decrease in SpO<sub>2</sub> by itself. Furthermore, we assessed the merits of SpO<sub>2</sub> as a diagnostic marker of AHF in patients with AMI, but this could not be tested in patients with significant chronic obstructive pulmonary disease or shock and it should be evaluated in the future in other acute care conditions. Finally, this series should not be considered as representative of the incidence of AHF in patients with acute coronary syndromes, as our population was recruited in a single community hospital with strongly biased admissions.

## CONCLUSIONS

In this study, baseline SpO<sub>2</sub> provided reliable information for establishing the diagnosis and severity of AHF as a complication of AMI with a warning cut-off value <93. The use of pulse oximetry for diagnosis purposes may be recommended when managing

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.