



(12) **United States Patent**
Naicker et al.

(10) **Patent No.:** **US 6,503,921 B2**
(45) **Date of Patent:** ***Jan. 7, 2003**

(54) **DEUTERATED RAPAMYCIN COMPOUNDS, METHODS AND USES THEREOF**

(75) Inventors: **Selvaraj Naicker**, Edmonton (CA);
Randall W. Yatscoff, Edmonton (CA);
Robert T. Foster, Edmonton (CA)

(73) Assignee: **Isotechnika, Inc.**, Edmonton (CA)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **09/987,812**

(22) Filed: **Nov. 16, 2001**

(65) **Prior Publication Data**

US 2002/0028827 A1 Mar. 7, 2002

Related U.S. Application Data

(63) Continuation of application No. 09/348,015, filed on Jul. 6, 1999, which is a continuation-in-part of application No. 09/148,623, filed on Sep. 4, 1998, now abandoned.

(60) Provisional application No. 60/057,632, filed on Sep. 5, 1997.

(51) **Int. Cl.**⁷ **C07D 491/16**; A61K 31/445

(52) **U.S. Cl.** **514/291**; 540/456

(58) **Field of Search** 540/456; 514/291

(56) **References Cited**
PUBLICATIONS

Dennis P. Curran, et al., Intramolecular Hydrogen Transfer Reaction of o-(Bromophenyl)dialkylsilyl Ethers. Preparation of Rapamycin-d₁, Tetrahedron Letters, vol. 33, No. 17, pp. 2295-2298.

Don Sticker, Senior Technical Information Specialist of Chemical Abstract Service, C.A.S., Commercial Database Search of Deuterated Rapamycin, search conducted Mar. 9, 2000.

Park et al., *Jr. Bio Chem.*, vol. 267, No. 5 (15) pp 3316-3324, 1992.

Connelly et al, *Biochemistry*, vol. 32, pp 5583-5590 (1993).

Primary Examiner—Bruck Kifle

(74) *Attorney, Agent, or Firm*—Burns Doane Swecker & Mathis LLP

(57) **ABSTRACT**

The synthesis of deuterated analogues of rapamycin is disclosed together with a method for use for inducing immunosuppression and in the treatment of transplantation rejection, graft vs host disease, autoimmune diseases, diseases of inflammation leukemia/lymphoma, solid tumors, fungal infections, hyperproliferative vascular disorders. Also described is a method for the synthesis of water soluble deuterated rapamycin compounds and their use as described above.

21 Claims, 2 Drawing Sheets




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5

DEUTERATED RAPAMYCIN COMPOUNDS, METHODS AND USES THEREOF

REFERENCE TO RELATED APPLICATIONS

This application is a continuation of Ser. No. 09/348,015 filed Jul. 6, 1999 which is a continuation-in-part of U.S. patent application Ser. No. 09/148,623, filed Sep. 4, 1998, now abandoned which is based on provisional patent application No. 60/057,632, filed Sep. 5, 1997 both of which are relied on and incorporated herein by reference.

BACKGROUND OF THE INVENTION

This invention relates to deuterated derivatives of rapamycin and a method for using them in the treatment of transplantation rejection, host vs. graft disease, graft vs. host disease, leukemia/lymphoma, hyperproliferative vascular disorders, autoimmune diseases, diseases of inflammation, solid tumors, and fungal infections.

Rapamycin, known as sirolimus, is a 31-membered macrolide lactone, $C_{51}H_{79}NO_{13}$, with a molecular mass of 913.6 Da. In solution, sirolimus forms two conformational trans-, cis-isomers with a ratio of 4:1 (chloroform) due to hindered rotation around the pipercolic acid amide bond. It is sparingly soluble in water, aliphatic hydrocarbons and diethyl ether, whereas it is soluble in alcohols, halogenated hydrocarbons and dimethyl sulfoxide. Rapamycin is unstable in solution and degrades in plasma and low-, and neutral-pH buffers at 37° C. with half-life of <10 h. the structures of the degradation products have recently been characterized. Rapamycin is a macrocyclic triene antibiotic produced by *Streptomyces hygroscopicus*, which was found to have antifungal activity, particularly against *Candida albicans*, both in vitro and in vivo [C. Vezina et al., J. Antibiot. 28, 721 (1975); S. N. Sehgal et al., J. Antibiot. 28, 727 (1975); H. A. Baker et al., J. Antibiot. 31, 539 (1978); U.S. Pat. Nos. 3,929,992; and 3,993,749].

Rapamycin alone (U.S. Pat. No. 4,885,171) or in combination with picibanil (U.S. Pat. No. 4,401,653) has been shown to have antitumor activity. R. Martel et al. [Can. J. Physiol. Pharmacol. 55, 48 (1977)] disclosed that rapamycin is effective in the experimental allergic encephalomyelitis model, a model for multiple sclerosis; in the adjuvant arthritis model, a model for rheumatoid arthritis; and effectively inhibited the formation of IgE-like antibodies.

The immunosuppressive effects of rapamycin have been disclosed in FASEB 3, 3411 (1989). Cyclosporin A and FK-506, other macrocyclic molecules, also have been shown to be effective as immunosuppressive agents, therefore useful in preventing transplant rejection [FASEB 3, 3411 (1989); FASEB 3, 5256 (1989); and R. Y. Calne et al., Lancet 1183 (1978)]. Although it shares structural homology with the immunosuppressant tacrolimus and binds to the same intracellular binding protein in lymphocytes, rapamycin inhibits S6p70-kinase and therefore has a mechanism of immunosuppressive action distinct from that of tacrolimus. Rapamycin was found to prolong graft survival of different transplants in several species alone or in combination with other immunosuppressants. In animal models its spectrum of toxic effects is different from that of cyclosporin or FK-506., comprising impairment of glucose homeostasis, stomach, ulceration, weight loss and thrombocytopenia, although no nephrotoxicity has been detected.

Mono- and diacylated derivatives of rapamycin (esterified at the 28 and 43 positions) have been shown to be useful as antifungal agents (U.S. Pat. No. 4,316,885) and used to make water soluble prodrugs of rapamycin (U.S. Pat. No.

4,650,803). Recently, the numbering convention for rapamycin has been changed; therefore according to Chemical Abstracts nomenclature, the esters described above would be at the 31- and 42-positions. Carboxylic acid esters (PCT application No. WO 92/05179), carbamates (U.S. Pat. No. 5,118,678), amide esters (U.S. Pat. No. 5,118,678), (U.S. Pat. No. 5,118,678) fluorinated esters (U.S. Pat. No. 5,100,883), acetals (U.S. Pat. No. 5,151,413), silyl ethers (U.S. Pat. No. 5,120,842), bicyclic derivatives (U.S. Pat. No. 5,120,725), rapamycin dimers (U.S. Pat. No. 5,120,727) and O-aryl, O-alkyl, O-alkylenyl and O-alkynyl derivatives (U.S. Pat. No. 5,258,389) have been described.

Rapamycin is metabolized by cytochrome P-450 3A to at least six metabolites. During incubation with human liver and small intestinal microsomes, sirolimus was hydroxylated and demethylated and the structure of 39-O-demethyl sirolimus was identified. In bile of sirolimus-treated rats >16 hydroxylated and demethylated metabolites were detected.

In rapamycin, demethylation of methoxy group at C-7 Carbon will lead to the change in the conformation of the Rapamycin due to the interaction of the released C-7 hydroxyl group with the neighbouring pyran ring system which is in equilibrium with the open form of the ring system. The C-7 hydroxyl group will also interact with the triene system and possibly alter the immunosuppressive activity of rapamycin. This accounts for the degradation of rapamycin molecule and its altered activity.

Stable isotopes (e.g., deuterium, ^{13}C , ^{15}N , ^{18}O) are non-radioactive isotopes which contain one additional neutron than the normally abundant isotope of the atom in question. Deuterated compounds have been used in pharmaceutical research to investigate the in vivo metabolic fate of the compounds by evaluation of the mechanism of action and metabolic pathway of the non deuterated parent compound. (Blake et al. J. Pharm. Sci. 64, 3, 367-391, 1975). Such metabolic studies are important in the design of safe, effective therapeutic drugs, either because the in vivo active compound administered to the patient or because the metabolites produced from the parent compound prove to be toxic or carcinogenic (Foster et al., Advances in drug Research Vol. 14, pp. 2-36, Academic press, London, 1985).

Incorporation of a heavy atom particularly substitution of deuterium for hydrogen, can give rise to an isotope effect that can alter the pharmacokinetics of the drug. This effect is usually insignificant if the label is placed in a molecule at the metabolically inert position of the molecule.

Stable isotope labeling of a drug can alter its physico-chemical properties such as pKa and lipid solubility. These changes may influence the fate of the drug at different steps along its passage through the body. Absorption, distribution, metabolism or excretion can be changed. Absorption and distribution are processes that depend primarily on the molecular size and the lipophilicity of the substance.

Drug metabolism can give rise to large isotopic effect if the breaking of a chemical bond to a deuterium atom is the rate limiting step in the process. While some of the physical properties of a stable isotope-labeled molecule are different from those of the unlabeled one, the chemical and biological properties are the same, with one important exception: because of the increased mass of the heavy isotope, any bond involving the heavy isotope and another atom will be stronger than the same bond between the light isotope and that atom. In any reaction in which the breaking of this bond is the rate limiting step, the reaction will proceed slower for the molecule with the heavy isotope due to kinetic isotope effect. A reaction involving breaking a C—D bond can be up

to 700 per cent slower than a similar reaction involving breaking a C—H bond.

More caution has to be observed when using deuterium labeled drugs. If the C—D bond is not involved in any of the steps leading to the metabolite, there may not be any effect to alter the behavior of the drug. If a deuterium is placed at a site involved in the metabolism of a drug, an isotope effect will be observed only if breaking of the C—D bond is the rate limiting step. There are evidences to suggest that whenever cleavage of an aliphatic C—H bond occurs, usually by oxidation catalyzed by a mixed-function oxidase, replacement of the hydrogen by deuterium will lead to observable isotope effect. It is also important to understand that the incorporation of deuterium at the site of metabolism slows its rate to the point where another metabolite produced by attack at a carbon atom not substituted by deuterium becomes the major pathway by a process called “metabolic switching”.

It is also observed that one of the most important metabolic pathways of compounds containing aromatic systems is hydroxylation leading to a phenolic group in the 3 or 4 position to carbon substituents. Although this pathway involves cleavage of the C—H bond, it is often not accompanied by an isotope effect, because the cleavage of this bond is mostly not involved in the rate-limiting step. The substitution of hydrogen by deuterium at the stereo center will induce a greater effect on the activity of the drug.

Clinically relevant questions include the toxicity of the drug and its metabolite derivatives, the changes in distribution or elimination (enzyme induction), lipophilicity which will have an effect on absorption of the drug. Replacement of hydrogen by deuterium at the site involving the metabolic reaction will lead to increased toxicity of the drug. Replacement of hydrogen by deuterium at the aliphatic carbons will have an isotopic effect to a larger extent. Deuterium placed at an aromatic carbon atom, which will be the site of hydroxylation, may lead to an observable isotope effect, although this is less often the case than with aliphatic carbons. But in few cases such as in penicillin, the substitution on the aromatic ring will induce the restriction of rotation of the ring around the C—C bond leading to a favorable stereo-specific situation to enhance the activity of the drug.

Approaching half a century of stable-isotope usage in human metabolic studies has been without documented significant adverse effect. Side-effects with acute D dosing are transitory with no demonstrated evidence of permanent deleterious action. The threshold of D toxicity has been defined in animals and is far in excess of concentrations conceivably used in human studies (Jones PJ, Leatherdale ST Clin Sci (Colch) April 1991; 80(4):277–280). The possibility that D may have additional beneficial pharmacological applications cannot be excluded. For isotopes other than D, evidence of observed toxicity remains to be produced even at dosages far in excess of the range used in metabolic studies. Absence of adverse effect may be attributable to small mass differences and the similar properties of tracer and predominantly abundant isotopes. The precision of extrapolating toxicity thresholds from animal studies remains unknown. However, should perturbation of the delicate homeostatic characteristic of living organisms occur with use of stable isotopes, it is almost undoubtedly at some level of administration greatly in excess of those administered currently in biomedical research.

In the prior art, no details are described regarding deuterated derivatives to improve the stability of rapamycin

molecule and also about glycosylated deuterated rapamycin to improve the stability and also the solubility of the molecule in order to increase the bio-availability of the drug. We therefore defined the global objective of preparing a rapamycin derivative which is more stable, less prone to degradation, and more water soluble to improve the bio-availability.

SUMMARY OF THE INVENTION

Deuteration of the rapamycin molecule results in altered physicochemical and pharmacokinetic properties which enhance its usefulness in the treatment of transplantation rejection, host vs. graft disease, graft vs. host disease, leukemia/lymphoma, hyperproliferative vascular disorders, autoimmune diseases, diseases of inflammation, solid tumors, and fungal infections.

Deuterium isotope is selected based on the fact that if ^{13}C , ^{15}N or another heavy isotope differing from the light one by less than 10% in mass is incorporated at the site of metabolism, there may be a small isotope effect. In addition to this, there are secondary isotope effects away from the site of isotope substitution due to changes in electronic environment.

Substitution of deuterium in methyl groups of rapamycin will result in a slower rate of oxidation of the C—D bond relative to the rate of oxidation of a non deuterium substituted C—H bond. The isotopic effect acts to reduce formation of demethylated metabolites and thereby alters the pharmacokinetic parameters of the drug. Lower rates of oxidation, metabolism and clearance result in greater and more sustained biological activity. Deuteration is targeted at various sites of the rapamycin molecule to increase the potency of drug, reduce toxicity of the drug, reduce the clearance of the pharmacologically active moiety and improve the stability of the molecule.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is the chemical structure of 7-deuteromethyl rapamycin showing sites of deuteration.

FIG. 2 is the chemical structure of epi-7 deuteromethyl rapamycin showing sites of deuteration.

FIG. 3 is the chemical structure of 7,43-d₆-rapamycin showing sites of deuteration.

FIG. 4 is the chemical structure of 31,42-d₂ showing sites of deuteration.

FIG. 5 illustrates the preparation of glycosylated deuterorapamycin.

DETAILED DESCRIPTION OF THE INVENTION

Substitution of deuterium for ordinary hydrogen and deuterated substrates for protio metabolites can produce profound changes in biosystems. Isotopically altered drugs have shown widely divergent pharmacological effects. Petersen et al., found increased anti-cancer effect with deuterated 5,6-benzylidene-dl-L-ascorbic acid (Zilascorb) [Anticancer Res. 12, 33 (1992)].

Substitution of deuterium in methyl groups of rapamycin will result in a slower rate of oxidation of the C—D bond relative to the rate of oxidation of a non deuterium substituted C—H bond. The isotopic effect acts to reduce formation of demethylated metabolites and thereby alters the pharmacokinetic parameters of the drug. Lower rates of oxidation, metabolism and clearance result in greater and more sustained biological activity. Deuteration is targeted at

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.