

Drug Facts

Active ingredients	Purpose
---------------------------	----------------

Naphazoline hydrochloride (0.012%).....reliever	Redness
Polyethylene glycol 300 (0.2%).....Lubricant	

Uses

- relieves redness of the eye due to minor eye irritations
- temporary relief of burning and irritation due to dryness of the eye
- protects against further irritation

Warnings

For external use only

Do not use

- if solution changes color or becomes cloudy

Ask a doctor before use if you have

- narrow angle glaucoma

When using this product

- do not touch tip of container to any

Drug Facts (continued)

surface to avoid contamination

- do not overuse as it may produce increased redness of the eye
- pupils may become enlarged temporarily
- remove contact lenses before using
- replace cap after use

Stop use and ask a doctor if

- you experience eye pain, changes in vision, continued redness or irritation of the eye
- condition worsens or persists for more than 72 hours

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- instill 1 to 2 drops in the affected eye(s) up to four times daily

Other information

- store at 15°-25°C (59°-77°F)
- keep tightly closed
- use before expiration date marked on the carton and bottle

Inactive ingredients

benzalkonium chloride (0.01%), boric acid, edetate disodium, sodium borate, sodium chloride. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

Drug Facts
(continued)

Questions
comments

(Call: 1-800-5