

Drug Facts

Active ingredients

Hypromellose (0.5%).....	Lubricant
Naphazoline hydrochloride (0.03%).....	Redness reliever

Purpose

Uses

- temporary relief of redness and discomfort due to:
 - minor eye irritations
 - exposure to wind or sun
 - dryness of the eye
- prevents further irritation

Warnings

For external use only

Do not use

- if solution changes color or becomes cloudy

Ask a doctor before use if you have

- narrow angle glaucoma

When using this product

- do not touch tip of container to any surface to avoid contamination
- do not overuse as it may produce increased redness of the eye
- pupils may become enlarged temporarily
- remove contact lenses before using
- replace cap after use

Stop use and ask a doctor if

- you experience eye pain, changes in vision, continued redness or irritation of the eye

Drug Facts (continued)

- condition worsens or persists for more than 72 hours

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- instill 1 to 2 drops in the affected eye(s) up to four times daily

Other information

- store at 15°-25°C (59°-77°F)
- keep tightly closed
- use before expiration date marked on the carton and bottle

Inactive ingredients

benzalkonium chloride (0.01%), boric acid, edetate disodium, purified water, sodium borate, sodium chloride. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

Questions or comments?

☎ Call: 1-800-553-5340