

US009254338B2

(12) United States Patent Yancopoulos

(54) USE OF A VEGF ANTAGONIST TO TREAT ANGIOGENIC EYE DISORDERS

- (71) Applicant: REGENERON PHARMACEUTICALS, INC., Tarrytown, NY (US)
- (72) Inventor: George D. Yancopoulos, Yorktown Heights, NY (US)
- (73) Assignee: Regeneron Pharmaceuticals, Inc., Tarrytown, NY (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 132 days.
- (21) Appl. No.: 13/940,370
- (22) Filed: Jul. 12, 2013

Prior Publication Data

US 2013/0295094 A1 Nov. 7, 2013

Related U.S. Application Data

- (63) Continuation-in-part of application No. PCT/US2012/020855, filed on Jan. 11, 2012.
- (60) Provisional application No. 61/432,245, filed on Jan. 13, 2011, provisional application No. 61/434,836, filed on Jan. 21, 2011, provisional application No. 61/561,957, filed on Nov. 21, 2011.
- (51) Int. Cl.

RM

(65)

A61K 38/18	(2006.01)
C07K 14/71	(2006.01)
A61K 47/48	(2006.01)
A61K 38/17	(2006.01)
C07K 16/22	(2006.01)
A61K 39/00	(2006.01)

- (52) U.S. CI.
 CPC A61K 47/48415 (2013.01); A61K 38/179 (2013.01); C07K 14/71 (2013.01); C07K 16/22 (2013.01); A61K 2039/505 (2013.01)
- (58) Field of Classification Search None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,396,664 B2	7/2008	Daly et al.
2003/0171320 A1	9/2003	Guyer
2005/0163798 A1	7/2005	Papadopoulos et al.
2005/0260203 A1	11/2005	Wiegand et al.
2006/0058234 A1	3/2006	Daly et al.
2006/0172944 A1	8/2006	Wiegand et al.
2007/0190058 A1	8/2007	Shams

FOREIGN PATENT DOCUMENTS

IР	2010-509369	3/2010	
WO	00/75319	12/2000	
WO	2008/063932	5/2008	

(10) Patent No.: US 9,254,338 B2 (45) Date of Patent: Feb. 9, 2016

OTHER PUBLICATIONS

Anonymous "Lucentis (rangibizymab injection) Intravitreal Injection" pp. 103 (Jun. 2006).

Do et al., "An exploratory study of the safety, tolerability and bioactivity of a single intravitreal injection of vascular endothelial growth factor Trap-Eye in patients with diabetic macular oedema" Br J Opthamol. 93(2):144-1449 (Feb. 2009).

Do et al., "The DA VINCI Study: phase 2 primary results of VEGF Trap-Eye in patients with diabetic macular edema" Opthamology 118(9):1819-1826 (Sep. 2011).

The Eyetech Study Group, "Anti-Vascular Endothelial Growth Factor Therapy for Subfoveal Choroidal Neovascularization Secondary to Age-related Macular Degeneration" American Academy of Ophthamology, 110 (5):979-986 (May 2003).

Heieret al., "rhuFab V2 (anti-VEGF Antibody) for Treatment of Exudative AMD" Symposium 8:Experimental and Emerging Treatments for Choroidal Neovascularization, 10 pp (2002).

Heier et al., "RhuFab V2 in Wet AMD—6 Month Continued Improvement Following Multiple Intravitreal Injections" Invest Ophthalmol Vis Sci, 44:E-Abstract 972 (2003).

Krzystolik et al., "Prevention of Experimental Choroidal NEovascularization With Intravitreal Anti-Vascular Endothelial Growth Factor Antibody Fragment" Arch Ophthamol., 120:338-346 (Mar. 2002).

Nguyen et al., "A Phase I Study of Intravitreal Vascular Endothelial Growth Factor Trap-Eye in Patients with Neovascular Age-Related Macular Degeneration" Opthamology, J.B. Lippincott Co., Philadelphia, PA, US, 116(11):2141-2148 (Nov. 1, 2009).

Nichols, Earl R., "AAO: Ranibizumab (rhuRab) May Improve Vision in Age-Related Macular Degeneration" Doctor's Guide Global Edition, www.pslgroup.com/dg/23f2aa.htm, pp. 1-2 (Nov. 24, 20013). Pai et al., "Current concepts in intravitreal drug therapy for diabetic retinopathy" Saudi Journal of Opthamology 24(4):143-149 (Jun. 30, 2010).

Stewart, "THe expanding role of vascular endothelial growth factor inhibitors in opthamology" Mayo Clin Proc. 87 (1):77-88 (Jan. 2012).

Thomas Reuters Integrity "VEGF Trap-Eye final phase II results in age-related macular degeneration presented at 2008 Retina Society Meeting" (Sep. 28, 2008). Nguyen et al., "A phase I trial of an IV-administered vascular

Nguyen et al., "A phase I trial of an IV-administered vascular endothelial growth factor trap for treatment in patients with choroidal neovascularization due to age-related macular degeneration" Ophthalmology (Sep. 2006) 113 (9):1522e1-1522e14 (epub Jul. 28, 2006).

Primary Examiner - Christine J Saoud

Assistant Examiner — Jon M Lockard

(74) Attorney, Agent, or Firm-Frank Cottingham; Karl Bozicevic

(57) ABSTRACT

The present invention provides methods for treating angiogenic eye disorders by sequentially administering multiple doses of a VEGF antagonist to a patient. The methods of the present invention include the administration of multiple doses of a VEGF antagonist to a patient at a frequency of once every 8 or more weeks. The methods of the present invention are useful for the treatment of angiogenic eye disorders such as age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, branch retinal vein occlusion, and corneal neovascularization.

26 Claims, 1 Drawing Sheet

D

Α

OCKF'

Δ

R

Μ



Mylan Exhibit 1019

15

USE OF A VEGF ANTAGONIST TO TREAT ANGIOGENIC EYE DISORDERS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of International Patent Application No. PCT/US2012/020855, filed on Jan. 11, 2012, which claims the benefit of U.S. Provisional Application Nos. 61/432,245, filed on Jan. 13, 2011, 61/434,836, filed on Jan. 21, 2011, and 61/561,957, filed on Nov. 21, 2011, the contents of which are hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to the field of therapeutic treatments of eye disorders. More specifically, the invention relates to the administration of VEGF antagonists to treat eye disorders caused by or associated with angiogenesis. 20

BACKGROUND

Several eye disorders are associated with pathological angiogenesis. For example, the development of age-related 25 macular degeneration (AMD) is associated with a process called choroidal neovascularization (CNV). Leakage from the CNV causes macular edema and collection of fluid beneath the macula resulting in vision loss. Diabetic macular edema (DME) is another eye disorder with an angiogenic 30 component. DME is the most prevalent cause of moderate vision loss in patients with diabetes and is a common complication of diabetic retinopathy, a disease affecting the blood vessels of the retina. Clinically significant DME occurs when fluid leaks into the center of the macula, the light-sensitive 35 part of the retina responsible for sharp, direct vision. Fluid in the macula can cause severe vision loss or blindness. Yet another eye disorder associated with abnormal angiogenesis is central retinal vein occlusion (CRVO). CRVO is caused by obstruction of the central retinal vein that leads to a back-up 40 of blood and fluid in the retina. The retina can also become ischemic, resulting in the growth of new, inappropriate blood vessels that can cause further vision loss and more serious complications. Release of vascular endothelial growth factor (VEGF) contributes to increased vascular permeability in the 45 eye and inappropriate new vessel growth. Thus, inhibiting the angiogenic-promoting properties of VEGF appears to be an effective strategy for treating angiogenic eye disorders

FDA-approved treatments of angiogenic eye disorders such as AMD and CRVO include the administration of an ⁵⁰ anti-VEGF antibody called ranibizumab (Lucentis®, Genentech, Inc.) on a monthly basis by intravitreal injection.

Methods for treating eye disorders using VEGF antagonists are mentioned in, e.g., U.S. Pat. Nos. 7,303,746; 7,306, 799; 7,300,563; 7,303,748; and US 2007/0190058. Nonethe-55 less, there remains a need in the art for new administration regimens for angiogenic eye disorders, especially those which allow for less frequent dosing while maintaining a high level of efficacy.

BRIEF SUMMARY OF THE INVENTION

The present invention provides methods for treating angiogenic eye disorders. The methods of the invention comprise sequentially administering multiple doses of a VEGF antagonist to a patient over time. In particular, the methods of the invention comprise sequentially administering to the patient a

DOCKE

2

single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonists. The present inventors have surprisingly discovered that beneficial therapeutic effects can be achieved in patients suffering from angiogenic eye disorders by administering a VEGF antagonist to a patient at a frequency of once every 8 or more weeks, especially when such doses are preceded by about three doses administered to the patient at a frequency of about 2 to 4 weeks. Thus, according to the methods of the present invention, each secondary dose of VEGF antagonist is administered 2 to 4 weeks after the immediately preceding dose, and each tertiary dose is administered at least 8 weeks after the immediately preceding dose. An example of a dosing regimen of the present invention is shown in FIG. 1. One advantage of such a dosing regimen is that, for most of the course of treatment (i.e., the tertiary doses), it allows for less frequent dosing (e.g., once every 8 weeks) compared to prior administration regimens for angiogenic eye disorders which require monthly administrations throughout the entire course of treatment. (See, e.g., prescribing information for Lucentis® [ranibizumab], Genentech, Inc.). The methods of the present invention can be used to treat

The methods of the present invention can be used to treat any angiogenic eye disorder, including, e.g., age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, corneal neovascularization, etc.

The methods of the present invention comprise administering any VEGF antagonist to the patient. In one embodiment, the VEGF antagonist comprises one or more VEGF receptor-based chimeric molecule(s), (also referred to herein as a "VEGF-Trap" or "VEGFT"). An exemplary VEGF antagonist that can be used in the context of the present invention is a multimeric VEGF-binding protein comprising two or more VEGF receptor-based chimeric molecules referred to herein as "VEGFR1R2-FcAC1(a)" or "aflibercept."

Various administration routes are contemplated for use in the methods of the present invention, including, e.g., topical administration or intraocular administration (e.g., intravitreal administration).

Aflibercept (EYLEATM, Regeneron Pharmaceuticals, Inc) was approved by the FDA in November 2011, for the treatment of patients with neovascular (wet) age-related macular degeneration, with a recommended dose of 2 mg administered by intravitreal injection every 4 weeks for the first three months, followed by 2 mg administered by intravitreal injection once every 8 weeks.

Other embodiments of the present invention will become apparent from a review of the ensuing detailed description.

BRIEF DESCRIPTION OF THE FIGURE

FIG. **1** shows an exemplary dosing regimen of the present invention. In this regimen, a single "initial dose" of VEGF antagonist ("VEGFT") is administered at the beginning of the treatment regimen (i.e. at "week 0"), two "secondary doses" are administered at weeks 4 and 8, respectively, and at least six "tertiary doses" are administered once every 8 weeks thereafter i.e. at weeks 16, 24, 32, 40, 48, 56, etc.)

⁶⁰ thereafter, i.e., at weeks 16, 24, 32, 40, 48, 56, etc.).

DETAILED DESCRIPTION

Before the present invention is described, it is to be understood that this invention is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the

Mylan Exhibit 1019

terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein, the term "about," when used in reference to a particular recited numerical value, means that the value may vary from the recited value by no more than 10 1%. For example, as used herein, the expression "about 100" includes 99 and 101 and all values in between (e.g., 99.1, 99.2, 99.3, 99.4, etc.).

Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described.

Dosing Regimens

The present invention provides methods for treating angiogenic eye disorders. The methods of the invention comprise 20 sequentially administering to a patient multiple doses of a VEGF antagonist. As used herein, "sequentially administering" means that each dose of VEGF antagonist is administered to the patient at a different point in time, e.g., on different days separated by a predetermined interval (e.g., hours, 25 days, weeks or months). The present invention includes methods which comprise sequentially administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonist.

The terms "initial dose," "secondary doses," and "tertiary doses," refer to the temporal sequence of administration of the VEGF antagonist. Thus, the "initial dose" is the dose which is administered at the beginning of the treatment regimen (also referred to as the "baseline dose"); the "secondary doses" are 35 the doses which are administered after the initial dose; and the "tertiary doses" are the doses which are administered after the secondary doses. The initial, secondary, and tertiary doses may all contain the same amount of VEGF antagonist, but will generally differ from one another in terms of frequency 40 of administration. In certain embodiments, however, the amount of VEGF antagonist contained in the initial, secondary and/or tertiary doses will vary from one another (e.g., adjusted up or down as appropriate) during the course of treatment.

In one exemplary embodiment of the present invention, each secondary dose is administered 2 to 4 (e.g., 2, 21/2, 3, 31/2, or 4) weeks after the immediately preceding dose, and each tertiary dose is administered at least 8 (e.g., 8, 8¹/₂, 9, 9¹/₂, 10, 10¹/₂, 11, 11¹/₂, 12, 12¹/₂, 13, 13¹/₂, 14, 14¹/₂, or more) weeks 50 after the immediately preceding dose. The phrase "the immediately preceding dose," as used herein, means, in a sequence of multiple administrations, the dose of VEGF antagonist which is administered to a patient prior to the administration of the very next dose in the sequence with no intervening 55 doses

In one exemplary embodiment of the present invention, a single initial dose of a VEGF antagonist is administered to a patient on the first day of the treatment regimen (i.e., at week 0), followed by two secondary doses, each administered four 60 weeks after the immediately preceding dose (i.e., at week 4 and at week 8), followed by at least 5 tertiary doses, each administered eight weeks after the immediately preceding dose (i.e., at weeks 16, 24, 32, 40 and 48). The tertiary doses may continue (at intervals of 8 or more weeks) indefinitely 65 during the course of the treatment regimen. This exemplary administration regimen is depicted graphically in FIG. 1.

DOCKE

The methods of the invention may comprise administering to a patient any number of secondary and/or tertiary doses of a VEGF antagonist. For example, in certain embodiments, only a single secondary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) secondary doses are administered to the patient. Likewise, in certain embodiments, only a single tertiary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) tertiary doses are administered to the patient.

In embodiments involving multiple secondary doses, each secondary dose may be administered at the same frequency as the other secondary doses. For example, each secondary dose may be administered to the patient 4 weeks after the immediately preceding dose. Similarly, in embodiments involving multiple tertiary doses, each tertiary dose may be administered at the same frequency as the other tertiary doses. For example, each tertiary dose may be administered to the patient 8 weeks after the immediately preceding dose. Alternatively, the frequency at which the secondary and/or tertiary doses are administered to a patient can vary over the course of the treatment regimen. For example, the present invention includes methods which comprise administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by at least 5 tertiary doses of the VEGF antagonist, wherein the first four tertiary doses are administered 8 weeks after the immediately preceding dose, and wherein each subsequent tertiary dose is administered from 8 to 12 (e.g., $8, 8\frac{1}{2}$). 9, 9¹/₂, 10, 10¹/₂, 11, 11¹/₂, 12) weeks after the immediately preceding dose. The frequency of administration may also be adjusted during the course of treatment by a physician depending on the needs of the individual patient following clinical examination.

VEGF Antagonists

45

The methods of the present invention comprise administering to a patient a VEGF antagonist according to specified dosing regimens. As used herein, the expression "VEGF antagonist" means any molecule that blocks, reduces or interferes with the normal biological activity of VEGF.

VEGF antagonists include molecules which interfere with the interaction between VEGF and a natural VEGF receptor, e.g., molecules which bind to VEGF or a VEGF receptor and prevent or otherwise hinder the interaction between VEGF and a VEGF receptor. Specific exemplary VEGF antagonists include anti-VEGF antibodies, anti-VEGF receptor antibodies, and VEGF receptor-based chimeric molecules (also referred to herein as "VEGF-Traps").

VEGF receptor-based chimeric molecules include chimeric polypeptides which comprise two or more immunoglobulin (Ig)-like domains of a VEGF receptor such as VEGFR1 (also referred to as Flt1) and/or VEGFR2 (also referred to as Flk1 or KDR), and may also contain a multimerizing domain (e.g., an Fc domain which facilitates the multimerization [e.g., dimerization] of two or more chimeric polypeptides). An exemplary VEGF receptor-based chimeric molecule is a molecule referred to as VEGFR1R2-Fc∆C1(a) which is encoded by the nucleic acid sequence of SEQ ID NO:1. VEGFR1R2-Fc Δ C1(a) comprises three components: (1) a VEGFR1 component comprising amino acids 27 to 129 of SEQ ID NO:2; (2) a VEGFR2 component comprising amino acids 130 to 231 of SEQ ID NO:2; and (3) a multimerization component ("Fc Δ C1(a)") comprising amino acids 232 to 457 of SEQ ID NO:2 (the C-terminal amino acid of SEQ ID NO:2 [i.e., K458] may or may not be included in the VEGF antago-

Mylan Exhibit 1019

nist used in the methods of the invention; see e.g., U.S. Pat. No. 7,396,664). Amino acids 1-26 of SEQ ID NO:2 are the signal sequence.

The VEGF antagonist used in the Examples set forth herein below is a dimeric molecule comprising two VEGFR1R2- 5 $Fc\Delta C1$ (a) molecules and is referred to herein as "VEGFT." Additional VEGF receptor-based chimeric molecules which can be used in the context of the present invention are disclosed in U.S. Pat. Nos. 7,396,664, 7,303,746 and WO 00/75319. 10

Angiogenic Eye Disorders

The methods of the present invention can be used to treat any angiogenic eye disorder. The expression "angiogenic eye disorder," as used herein, means any disease of the eve which is caused by or associated with the growth or proliferation of 15 blood vessels or by blood vessel leakage. Non-limiting examples of angiogenic eye disorders that are treatable using the methods of the present invention include age-related macular degeneration (e.g., wet AMD, exudative AMD, etc.), retinal vein occlusion (RVO), central retinal vein occlusion 20 (CRVO; e.g., macular edema following CRVO), branch retinal vein occlusion (BRVO), diabetic macular edema (DME), choroidal neovascularization (CNV; e.g., myopic CNV), iris neovascularization, neovascular glaucoma, post-surgical fibrosis in glaucoma, proliferative vitreoretinopathy (PVR), 25 optic disc neovascularization, corneal neovascularization, retinal neovascularization, vitreal neovascularization, pannus, pterygium, vascular retinopathy, and diabetic retinopathies.

Pharmaceutical Formulations

The present invention includes methods in which the VEGF antagonist that is administered to the patient is contained within a pharmaceutical formulation. The pharmaceutical formulation may comprise the VEGF antagonist along with at least one inactive ingredient such as, e.g., a pharma- 35 ceutically acceptable carrier. Other agents may be incorporated into the pharmaceutical composition to provide improved transfer, delivery, tolerance, and the like. The term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in 40 the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the antibody is administered. A multitude of appropriate formulations can be found in the 45 formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, Pa., 1975), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lip- 50 ids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTINTM), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing 55 carbowax. Any of the foregoing mixtures may be appropriate in the context of the methods of the present invention, provided that the VEGF antagonist is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. See also 60 Powell et al. PDA (1998) J Pharm Sci Technol. 52:238-311 and the citations therein for additional information related to excipients and carriers well known to pharmaceutical chemists

Pharmaceutical formulations useful for administration by 65 injection in the context of the present invention may be prepared by dissolving, suspending or emulsifying a VEGF

DOCKE

6

antagonist in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor

oil)], etc. As the oily medium, there may be employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared can be filled in an appropriate ampoule if desired.

Modes of Administration

The VEGF antagonist (or pharmaceutical formulation comprising the VEGF antagonist) may be administered to the patient by any known delivery system and/or administration method. In certain embodiments, the VEGF antagonist is administered to the patient by ocular, intraocular, intravitreal or subconjunctival injection. In other embodiments, the VEGF antagonist can be administered to the patient by topical administration, e.g., via eye drops or other liquid, gel, ointment or fluid which contains the VEGF antagonist and can be applied directly to the eye. Other possible routes of administration include, e.g., intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral.

Amount of VEGF Antagonist Administered

Each dose of VEGF antagonist administered to the patient 30 over the course of the treatment regimen may contain the same, or substantially the same, amount of VEGF antagonist. Alternatively, the quantity of VEGF antagonist contained within the individual doses may vary over the course of the treatment regimen. For example, in certain embodiments, a first quantity of VEGF antagonist is administered in the initial dose, a second quantity of VEGF antagonist is administered in the secondary doses, and a third quantity of VEGF antagonist is administered in the tertiary doses. The present invention contemplates dosing schemes in which the quantity of VEGF antagonist contained within the individual doses increases over time (e.g., each subsequent dose contains more VEGF antagonist than the last), decreases over time (e.g., each subsequent dose contains less VEGF antagonist than the last), initially increases then decreases, initially decreases then increases, or remains the same throughout the course of the administration regimen.

The amount of VEGF antagonist administered to the patient in each dose is, in most cases, a therapeutically effective amount. As used herein, the phrase "therapeutically effective amount" means a dose of VEGF antagonist that results in a detectable improvement in one or more symptoms or indicia of an angiogenic eye disorder, or a dose of VEGF antagonist that inhibits, prevents, lessens, or delays the progression of an angiogenic eye disorder. In the case of an anti-VEGF antibody or a VEGF receptor-based chimeric molecule such as VEGFR1R2-Fc Δ C1(a), a therapeutically effective amount can be from about 0.05 mg to about 5 mg, e.g., about 0.05 mg, about 0.1 mg, about 0.15 mg, about 0.2 mg, about 0.25 mg, about 0.3 mg, about 0.35 mg, about 0.4 mg, about 0.45 mg, about 0.5 mg, about 0.55 mg, about 0.6 mg, about 0.65 mg, about 0.7 mg, about 0.75 mg, about 0.8 mg, about 0.85 mg, about 0.9 mg, about 1.0 mg, about 1.05 mg, about 1.1 mg, about 1.15 mg, about 1.2 mg, about 1.25 mg, about 1.3 mg, about 1.35 mg, about 1.4 mg, about 1.45 mg, about 1.5 mg, about 1.55 mg, about 1.6 mg, about 1.65 mg, about 1.7 mg, about 1.75 mg, about 1.8 mg, about 1.85

Mylan Exhibit 1019

DOCKET



Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

