



US005359030A

United States Patent [19]

[11] Patent Number: 5,359,030

Ekwuribe

[45] Date of Patent: Oct. 25, 1994

[54] CONJUGATION-STABILIZED POLYPEPTIDE COMPOSITIONS, THERAPEUTIC DELIVERY AND DIAGNOSTIC FORMULATIONS COMPRISING SAME, AND METHOD OF MAKING AND USING THE SAME

ethylene Glycol Derivatized Superoxide Dismutase," Pharm. Res. Comm., 1982 14: 11-120.
(List continued on next page.)

[75] Inventor: Nnochiri N. Ekwuribe, Southfield, Mich.

Primary Examiner—Jeffrey E. Russel
Assistant Examiner—Nancy J. Gromet
Attorney, Agent, or Firm—Steven J. Hultquist; Fran S. Wasserman

[73] Assignee: Protein Delivery, Inc., Durham, N.C.

[57] ABSTRACT

[21] Appl. No.: 59,701

A stabilized conjugated peptide complex comprising a peptide conjugatively coupled to a polymer including lipophilic and hydrophilic moieties, wherein the peptide may for example be selected from the group consisting of insulin, calcitonin, ACTH, glucagon, somatostatin, somatotropin, somatomedin, parathyroid hormone, erythropoietin, hypothalamic releasing factors, prolactin, thyroid stimulating hormones, endorphins, enkephalins, vasopressin, non-naturally occurring opioids, superoxide dismutase, interferon, asparaginase, arginase, arginine deaminase, adenosine deaminase, ribonuclease, trypsin, chymotrypsin, and papain. In a particular aspect, the invention comprises an insulin composition suitable for parenteral as well as non-parenteral administration, preferably oral or parenteral administration, comprising insulin covalently coupled with a polymer including (i) a linear polyalkylene glycol moiety and (ii) a lipophilic moiety, wherein the insulin, the linear polyalkylene glycol moiety and the lipophilic moiety are conformationally arranged in relation to one another such that the insulin in the composition has an enhanced in vivo resistance to enzymatic degradation, relative to insulin alone. One, two, or three polymer constituents may be covalently attached to the insulin molecule, with one polymer constituent being preferred. The conjugates of the invention are usefully employed in therapeutic as well as non-therapeutic, e.g., diagnostic, applications, and the peptide and polymer may be covalently coupled to one another, or alternatively may be associatively coupled to one another, e.g., by hydrogen bonding or other associative bonding relationship.

[22] Filed: May 10, 1993

[51] Int. Cl.⁵ C07K 7/40; C07K 7/36; C07K 17/08; C08H 1/00

[52] U.S. Cl. 530/303; 530/307; 530/309; 530/322; 530/345; 530/402; 530/351; 530/409; 530/410; 530/411; 435/188; 424/85.1; 424/85.4; 424/94.3

[58] Field of Search 435/188; 514/3, 4, 12; 530/303, 307, 324, 309, 345, 322, 402, 326, 409, 410, 411, 325, 351; 424/85.1, 85.2, 85.4, 85.5, 85.6, 85.9, 94.3

[56] References Cited

U.S. PATENT DOCUMENTS

4,003,792	1/1977	Mill et al.	530/303
4,094,196	8/1977	Hüper et al.	526/271
4,179,337	12/1979	Davis et al.	435/181
4,585,754	4/1986	Meisner et al.	514/8
4,849,405	7/1989	Ecanow	514/3
4,963,367	10/1990	Ecanow	424/485
5,013,556	5/1991	Woodle et al.	424/450

OTHER PUBLICATIONS


Nucci, et al. "The Therapeutic Value of Poly(ethylen Glycol)—Modified Proteins" Ac. Drug. Del. Rev. 6: 133-151 1991.

Conradi, R. A., et al., "The Influence of Peptide Structure on Transport Across Caco-2 Cells," Pharm. Res., 1991, 8 (12): 1453-1459.

Abuchowski, A. and F. F. Davis, "Soluble Polymer--Enzyme Adducts," pp. 368-383, Enzymes as Drugs, J. S. Holcenberg, John Wiley, 1981.


Boccu, E. et al., "Pharmacokinetic Properties of Poly-


33 Claims, 2 Drawing Sheets



OTHER PUBLICATIONS

- Igarashi, R. et al, "Biologically Active Peptides Conjugated with Lecithin for DDS" Proceed. Intern. Symp. Cont. Rel. Bioactiv. Mater. 1990, 17 367-368.
- Taniguchi, T. et al, "Synthesis of Acyloyl Lysozyme and Improvement of its Lymphatic Transport Following Small Intestinal Administration in Rats" Proceed. Intern. Symp. Control. Rel. Bioactiv. Mater., 1992, 19: 104-105.
- Russell-Jones, G. J. "Vitamin B12 Drug Delivery", Proceed. Intern. Symp. Control. Rel. Bioactive. Mater., 1992, 19: 102-103.
- Baudys, M. et al, "Synthesis and Characteristics of Different Glycosylated Derivatives of Insulin" Proceed. Intern. Symp. Cont. Rel. Bioactive. Mater., 1992, 19: 210-211.
- Chien, Y. W., Novel Drug Delivery Systems, pp. 678-679, Marcell Deffer, Inc., New York, N.Y., 1992.
- Santiago, N. et al, "Oral Immunization of Rats with Influenza Virus M Protein (M1) Microspheres," Proceed. Intern. Symp. Cont. Rel. Bioactive. Mater., 1992, 19: 116-117.
- Banting, R. G., et al, "Pancreatic Extracts in the Treatment of Diabetes Mellitus," The Canadian Med. Assoc. J. 1922, 12: 141-146.
- Brange, J. et al, "Chemical Stability of Insulin. 1. Hydrolytic Degradation During Storage of Pharmaceutical Preparations," Pharm. Res., 1992, 9 (6): 715-726.
- Brange, J. et al, "Chemical Stability of Insulin. 2. Formation of Higher Molecular Weight Transformation Products During Storage of Pharmaceutical Preparations," Pharm. Res., 1992, 9 (6) 727-734.
- Robbins, D. C. et al, "Antibodies to Covalent Aggregates of Insulin in Blood of Insulin-Using Diabetic Patients" Diabetes, 1987, 36: 838-841.
- M. Maislos et al, "The Source of the Circulating Aggregate of Insulin in Type I Diabetic Patients is Therapeutic Insulin" J. Clin. Invest., 1986, 77: 717-723
- Ratner, R. E. et al, "Persistent Cutaneous Insulin Allergy Resulting from High-Molecular Weight Insulin Aggregates," Diabetes, 1990, 39: 728-733.
- Oka, K. et al, "Enhanced Intestinal Absorption of a Hydrophobic Polymer-conjugated Protein Drug, Smancs, in an Oily Formulation" Pharm. Res., 1990, 7 (8): 852-855.
- Saffran, M. et al, "A New Approach to the Oral Administration of Insulin and Other Peptide Drugs," Science, 1986, 233: 1081-1084.





F

**CONJUGATION-STABILIZED POLYPEPTIDE
COMPOSITIONS, THERAPEUTIC DELIVERY
AND DIAGNOSTIC FORMULATIONS
COMPRISING SAME, AND METHOD OF MAKING
AND USING THE SAME**

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to conjugation-stabilized (poly)peptide and protein compositions and formulations, and to methods of making and using same.

2. Description of the Related Art

The use of polypeptides and proteins for the systemic treatment of certain diseases is now well accepted in medical practice. The role that the peptides play in replacement therapy is so important that many research activities are being directed towards the synthesis of large quantities by recombinant DNA technology. Many of these polypeptides are endogenous molecules which are very potent and specific in eliciting their biological actions.

A major factor limiting the usefulness of these substances for their intended application is that they are easily metabolized by plasma proteases when given parenterally. The oral route of administration of these substances is even more problematic because in addition to proteolysis in the stomach, the high acidity of the stomach destroys them before they reach their intended target tissue. Polypeptides and protein fragments, produced by the action of gastric and pancreatic enzymes, are cleaved by exo and endopeptidases in the intestinal brush border membrane to yield di- and tripeptides, and even if proteolysis by pancreatic enzymes is avoided, polypeptides are subject to degradation by brush border peptidases. Any of the given peptides that survive passage through the stomach are further subjected to metabolism in the intestinal mucosa where a penetration barrier prevents entry into the cells.

In spite of these obstacles, there is substantial evidence in the literature to suggest that nutritional and pharmaceutical proteins are absorbed through the intestinal mucosa. On the other hand, nutritional and drug (poly)peptides are absorbed by specific peptide transporters in the intestinal mucosa cells. These findings indicate that properly formulated (poly)peptides and proteins may be administered by the oral route, with retention of sufficient biological activity for their intended use. If, however, it were possible to modify these peptides so that their physiological activities were maintained totally, or at least to a significant degree, and at the same time stabilize them against proteolytic enzymes and enhance their penetration capability through the intestinal mucosa, then it would be possible to utilize them properly for their intended purpose. The product so obtained would offer advantages in that more efficient absorption would result, with the concomitant ability to use lower doses to elicit the optimum therapeutic effect.

The problems associated with oral or parenteral administration of proteins are well known in the pharmaceutical industry, and various strategies are being used in attempts to solve them. These strategies include incorporation of penetration enhancers, such as the salicylates, lipid-bile salt-mixed micelles, glycerides, and acylcarnitines, but these frequently are found to cause

and inflammation of tissue. These problems arise because enhancers are usually coadministered with the peptide product and leakages from the dosage form often occur. Other strategies to improve oral delivery include mixing the peptides with protease inhibitors, such as aprotinin, soybean trypsin inhibitor, and amastatin, in an attempt to limit degradation of the administered therapeutic agent. Unfortunately these protease inhibitors are not selective, and endogenous proteins are also inhibited. This effect is undesirable.

Enhanced penetration of peptides across mucosal membranes has also been pursued by modifying the physicochemical properties of candidate drugs. Results indicate that simply raising lipophilicity is not sufficient to increase paracellular transport. Indeed it has been suggested that cleaving the peptide-water hydrogen bonds is the main energy barrier to overcome in obtaining peptide diffusion across membranes (Conradi, R. A., Hilgers, A. R., Ho, N. F. H., and Burton, P. S., "The influence of peptide structure on transport across Caco-2 cells", *Pharm. Res.*, 8, 1453-1460, (1991)). Protein stabilization has been described by several authors. Abuchowski and Davis ("Soluble polymers-Enzyme adducts", In: *Enzymes as Drugs*, Eds. Holcenberg and Roberts, J. Wiley and Sons, New York, N.Y., (1981)) disclosed various methods of derivatization of enzymes to provide water soluble, non-immunogenic, in vivo stabilized products.

A great deal of work dealing with protein stabilization has been published. Abuchowski and Davis disclose various ways of conjugating enzymes with polymeric materials (Ibid). More specifically, these polymers are dextrans, polyvinyl pyrrolidones, glycopeptides, polyethylene glycol and polyamino acids. The resulting conjugated polypeptides are reported to retain their biological activities and solubility in water for parenteral applications. The same authors, in U.S. Pat. No. 4,179,337, disclose that polyethylene glycol rendered proteins soluble and non-immunogenic when coupled to such proteins. These polymeric materials, however, did not contain fragments suited for intestinal mucosa binding, nor did they contain any moieties that would facilitate or enhance membrane penetration. While these conjugates were water-soluble, they were not intended for oral administration.

Meisner et al., U.S. Pat. No. 4,585,754, teaches that proteins may be stabilized by conjugating them with chondroitin sulfates. Products of this combination are usually polyanionic, very hydrophilic, and lack cell penetration capability. They are usually not intended for oral administration.

Mill et al., U.S. Pat. No. 4,003,792, teaches that certain acidic polysaccharides, such as pectin, alginate, hyaluronic acid and carrageenan, can be coupled to proteins to produce both soluble and insoluble products. Such polysaccharides are polyanionic, derived from food plants. They lack cell penetration capability and are usually not intended for oral administration.

In Pharmacological Research Communication 14, 11-120 (1982), Boccu et al. disclosed that polyethylene glycol could be linked to a protein such as superoxide dismutase ("SOD"). The resulting conjugated product showed increased stability against denaturation and enzymatic digestion. The polymers did not contain moieties that are necessary for membrane interaction

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.