

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/078683 A1

(51) International Patent Classification⁷: A61K 31/00, 38/00, A01N 61/00, C12Q 1/00, C07K 2/00, 4/00, 5/00, 7/00, 14/00, 16/00, 17/00, G01N 33/53, 33/48, 33/567, 33/574

(74) Agents: SANZO, Michael, A. et al.; Pillsbury Winthrop LLP, 1600 Tysons Boulevard, McLean, VA 22102 (US).

(21) International Application Number: PCT/US02/09551

(22) International Filing Date: 28 March 2002 (28.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:		
60/279,438	29 March 2001 (29.03.2001)	US
60/279,437	29 March 2001 (29.03.2001)	US
60/300,850	27 June 2001 (27.06.2001)	US
60/303,806	10 July 2001 (10.07.2001)	US
60/307,358	25 July 2001 (25.07.2001)	US
60/348,646	17 January 2002 (17.01.2002)	US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: SYNERGY PHARMACEUTICALS, INC. [US/US]; Suite 450, Two Executive Drive, Somerset, NJ 08873 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(72) Inventors: SHAILUBHAI, Kunwar; 600 Wick Lane, Blue Bell, PA 19422 (US). NIKIFOROVICH, Gregory; 751 Aramis Drive, St. Louis, MO 63141 (US). JACOB, Gary, S.; 12541 Mason Forest Drive, Creve Coeur, MO 63141 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/078683 A1

(54) Title: GUANYLATE CYCLASE RECEPTOR AGONISTS FOR THE TREATMENT OF TISSUE INFLAMMATION AND CARCINOGENESIS

(57) Abstract: A method of treatment of inflamed, pre-cancerous or cancerous tissue or polyps in a mammalian subject is disclosed. The treatment involves administration of a composition of at least one peptide agonist of a guanylate cyclase receptor and/or other small molecules that enhance intracellular production of cGMP. The at least one peptide agonist of a guanylate cyclase receptor may be administered either alone or in combination with an inhibitor of cGMP-dependent phosphodiesterase. The inhibitor may be a small molecule, peptide, protein or other compound that inhibits the degradation of cGMP. Without requiring a particular mechanism of action, this treatment may restore a healthy balance between proliferation and apoptosis in the subject's population of epithelial cells, and also suppress carcinogenesis. Thus, the method may be used to treat, <i>inter alia</i>, inflammation, including gastrointestinal inflammatory disorders, general organ inflammation and asthma, and carcinogenesis of the lung, gastrointestinal tract, bladder, testis, prostate and pancreas, or polyps.

Guanylate Cyclase Receptor Agonists for the Treatment of Tissue Inflammation and Carcinogenesis

Cross Reference to Related Applications

5 The present application claims the benefit of U.S. provisional application nos. 60/279,438, filed on March 29, 2001; 60/279,437, filed on March 29, 2001; 60/300,850, filed on June 27, 2001; 60/303,806, filed on July 10, 2001; 60/307,358, filed on July 25, 2001; and 60/348,646, filed on January 17, 2002.

10 Field of the Invention

 The present invention relates to the therapeutic use of guanylate cyclase receptor agonists as a means for enhancing the intracellular production of cGMP. The agonists may be used either alone or in combination with inhibitors of cGMP-specific phosphodiesterase to prevent or treat cancerous, pre-cancerous and metastatic growths, particularly in the
15 gastrointestinal tract and lungs. In addition, the agonists may be used in the treatment of inflammatory disorders such as ulcerative colitis and asthma.

Background of the Invention

 Uroguanylin, guanylin and bacterial ST peptides are structurally related peptides that
20 bind to a guanylate cyclase receptor and stimulate intracellular production of cyclic guanosine monophosphate (cGMP) (1-6). This results in the activation of the cystic fibrosis transmembrane conductance regulator (CFTR), an apical membrane channel for efflux of chloride from enterocytes lining the intestinal tract (1-6). Activation of CFTR and the subsequent enhancement of transepithelial secretion of chloride leads to stimulation of sodium
25 and water secretion into the intestinal lumen. Therefore, by serving as paracrine regulators of CFTR activity, cGMP receptor agonists regulate fluid and electrolyte transport in the GI tract (1-6; US patent 5,489,670).

 The process of epithelial renewal involves the proliferation, migration, differentiation,
30 senescence, and eventual loss of GI cells in the lumen (7,8). The GI mucosa can be divided into three distinct zones based on the proliferation index of epithelial cells. One of these zones, the proliferative zone, consists of undifferentiated stem cells responsible for providing a constant source of new cells. The stem cells migrate upward toward the lumen to which they

are extruded. As they migrate, the cells lose their capacity to divide and become differentiated for carrying out specialized functions of the GI mucosa (9). Renewal of GI mucosa is very rapid with complete turnover occurring within a 24-48 hour period (9). During this process mutated and unwanted cells are replenished with new cells. Hence, homeostasis of the GI
5 mucosa is regulated by continual maintenance of the balance between proliferation and apoptotic rates (8).

The rates of cell proliferation and apoptosis in the gut epithelium can be increased or decreased in a wide variety of different circumstances, *e.g.*, in response to physiological stimuli such as aging, inflammatory signals, hormones, peptides, growth factors, chemicals and
10 dietary habits. In addition, an enhanced proliferation rate is frequently associated with a reduction in turnover time and an expansion of the proliferative zone (10). The proliferation index has been observed to be much higher in pathological cases of ulcerative colitis and other GI disorders (11). Thus, intestinal hyperplasia is the major promoter of gastrointestinal inflammation and carcinogenesis.

15

In addition to a role for uroguanylin and guanylin as modulators of intestinal fluid and ion secretion, these peptides may also be involved in the continual renewal of GI mucosa. Previously published data in WO 01/25266 suggests a peptide with the active domain of uroguanylin may function as an inhibitor of polyp development in the colon and may constitute
20 a treatment of colon cancer. However, the mechanism by which this is claimed to occur is questionable in that WO 01/25266 teaches uroguanylin agonist peptides that bind specifically to a guanylate cyclase receptor, termed GC-C, that was first described as the receptor for *E. coli* heat-stable enterotoxin (ST) (4). Knockout mice lacking this guanylate cyclase receptor show resistance to ST in intestine, but effects of uroguanylin and ST are not disturbed in the
25 kidney *in vivo* (3). These results were further supported by the fact that membrane depolarization induced by guanylin was blocked by genistein, a tyrosine kinase inhibitor, whereas hyperpolarization induced by uroguanylin was not effected (12,13). Taken together these data suggest that uroguanylin also binds to a currently unknown receptor, which is distinct from GC-C.

30

Other papers have reported that production of uroguanylin and guanylin is dramatically decreased in pre-cancerous colon polyps and tumor tissues (14-17). In addition, genes for both uroguanylin and guanylin have been shown to be localized to regions of the genome frequently

associated with loss of heterozygosity in human colon carcinoma (18-20). Taken together, these findings indicate that uroguanylin, guanylin and other peptides with similar activity may be used in the prevention or treatment of abnormal colon growths. This proposal is bolstered by a recent study demonstrating oral administration of uroguanylin inhibits polyp formation in mice (15,16).

Uroguanylin and guanylin peptides also appear to promote apoptosis by controlling cellular ion flux. Alterations in apoptosis have been associated with tumor progression to the metastatic phenotype. While a primary gastrointestinal (GI) cancer is limited to the small intestine, colon, and rectum, it may metastasize and spread to such localities as bone, lymph nodes, liver, lung, peritoneum, ovaries, brain. By enhancing the efflux of K^+ and influx of Ca^{++} , uroguanylin and related peptides may promote the death of transformed cells and thereby inhibit metastasis.

One of the clinical manifestations of reduced CFTR activity is the inflammation of airway passages (21). This effect may be due to CFTR regulating the expression of NF-kB, chemokines and cytokines (22-25). Recent reports have also suggested that the CFTR channel is involved in the transport and maintenance of reduced glutathione, an antioxidant that plays an important role in protecting against inflammation caused by oxidative stress (39). Enhancement of intracellular levels of cGMP by way of guanylate cyclase activation or by way of inhibition of cGMP-specific phosphodiesterase would be expected to down-regulate these inflammatory stimuli. Thus, uroguanylin-type agonists should be useful in the prevention and treatment of inflammatory diseases of the lung (*e.g.*, asthma), bowel (*e.g.*, ulcerative colitis and Crohn's disease), pancreas and other organs.

Overall, it may be concluded that agonists of guanylate cyclase receptor such as uroguanylin have potential therapeutic value in the treatment of a wide variety of inflammatory conditions, cancer (particularly colon cancer) and as anti-metastatic agents. The development of new agonists is therefore of substantial clinical importance.

Summary of the Invention

The present invention is based upon the development of new agonists of guanylate cyclase receptor, and new uses of naturally occurring agonists. The agonists are analogs of

uroguanylin, many of which have superior properties either in terms of improved receptor activation, stability, activity at low pH or reduced adverse effects. The peptides may be used to treat any condition that responds to enhanced intracellular levels of cGMP. Intracellular levels of cGMP can be increased by enhancing intracellular production of cGMP and/or by inhibition
5 of its degradation by cGMP-specific phosphodiesterases. Among the specific conditions that can be treated or prevented are inflammatory conditions, cancer, polyps, and metastasis.

In its first aspect, the present invention is directed to a peptide consisting essentially of the amino acid sequence of any one of SEQ ID NOs:2-21 and to therapeutic compositions
10 which contain these peptides. The term “consisting essentially of” includes peptides that are identical to a recited sequence identification number and other sequences that do not differ substantially in terms of either structure or function. For the purpose of the present application, a peptide differs substantially if its structure varies by more than three amino acids from a peptide of SEQ ID NOs:2-21 or if its activation of cellular cGMP production is reduced or
15 enhanced by more than 50%. Preferably, substantially similar peptides should differ by no more than two amino acids and not differ by more than about 25% with respect to activating cGMP production. The most preferred peptide is a bicycle having the sequence of SEQ ID NO:20.

20 The peptides may be in a pharmaceutical composition in unit dose form, together with one or more pharmaceutically acceptable excipients. The term “unit dose form” refers to a single drug delivery entity, *e.g.*, a tablet, capsule, solution or inhalation formulation. The amount of peptide present should be sufficient to have a positive therapeutic effect when administered to a patient (typically, between 100 µg and 3 g). What constitutes a “positive
25 therapeutic effect” will depend upon the particular condition being treated and will include any significant improvement in a condition readily recognized by one of skill in the art. For example, it may constitute a reduction in inflammation, a shrinkage of polyps or tumors, a reduction in metastatic lesions, etc.

30 The invention also encompasses combination therapy utilizing a guanylate cyclase receptor agonist administered either alone or together with an inhibitor of cGMP-dependent phosphodiesterase, an anti-inflammatory agent or an anticancer agent. These agents should be present in amounts known in the art to be therapeutically effective when administered to a

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.