

Technology), and IDH2 antibody (Abcam). Metabolites were extracted from cultured cells and from tissue samples according to close variants of a previously reported method (Lu, W., Kimball, E. & Rabinowitz, J. D. *J Am Soc Mass Spectrom* 17, 37-50 (2006)), using 80% aqueous methanol (-80 °C) and either tissue scraping or homogenization to disrupt cells. Enzymatic activity in cell lysates was assessed by following a change in NADPH fluorescence over time in the presence of isocitrate and NADP, or  $\alpha$ KG and NADPH. For enzyme assays using recombinant IDH1 enzyme, proteins were produced in *E. coli* and purified using Ni affinity chromatography followed by Sephacryl S-200 size-exclusion chromatography. Enzymatic activity for recombinant IDH1 protein was assessed by following a change in NADPH UV absorbance at 340 nm using a stop-flow spectrophotometer in the presence of isocitrate and NADP or  $\alpha$ KG and NADPH. Chirality of 2HG was determined as described previously (Struys, E. A., Jansen, E. E., Verhoeven, N. M. & Jakobs, C. *Clin Chem* 50, 1391-5 (2004)). For crystallography studies, purified recombinant IDH1 (R132H) at 10 mg/mL in 20 mM Tris pH 7.4, 100 mM NaCl was pre-incubated for 60 min with 10 mM NADPH, 10 mM calcium chloride, and 75 mM  $\alpha$ KG. Crystals were obtained at 20°C by vapor diffusion equilibration using 3  $\mu$ L drops mixed 2:1 (protein:precipitant) against a well-solution of 100 mM MES pH 6.5, 20% PEG 6000. Patient tumor samples were obtained after informed consent as part of a UCLA IRB-approved research protocol. Brain tumor samples were obtained after surgical resection, snap frozen in isopentane cooled by liquid nitrogen and stored at -80 C. The IDH1 mutation status of each sample was determined using standard molecular biology techniques as described previously (Yan, H. et al. *N Engl J Med* 360, 765-73 (2009)). Metabolites were extracted and analyzed by LC-MS/MS as described above. Full methods are available in the supplementary material.

### ***Supplementary methods***

**Cloning, Expression, and Purification of ICDH1 wt and mutants in *E. coli*.** The open reading frame (ORF) clone of human isocitrate dehydrogenase 1 (cDNA) (IDH1; ref. ID NM\_005896) was purchased from Invitrogen in pENTR221 (Carlsbad, CA) and Origene Inc. in pCMV6 (Rockville, MD). To transfect cells with wild-type or mutant IDH1, standard molecular biology mutagenesis techniques were utilized to alter the DNA sequence at base pair 395 of the ORF in pCMV6 to introduce base pair change from guanine to adenine, which resulted in a change in the amino acid code at

position 132 from arginine (wt) to histidine (mutant; or R132H), and confirmed by standard DNA sequencing methods. For 293T cell transfection, wild-type and R132H mutant IDH1 were subcloned into pCMV-Sport6 with or without a carboxy-terminal Myc-DDK-tag. For stable cell line generation, constructs in pCMV6 were used. For expression in *E. coli*, the coding region was amplified from pENTR221 by PCR using primers designed to add NDEI and XHO1 restrictions sites at the 5' and 3' ends respectively. The resultant fragment was cloned into vector pET41a (EMD Biosciences, Madison, WI) to enable the *E. coli* expression of C-terminus His8-tagged protein. Site directed mutagenesis was performed on the pET41a-ICHD1 plasmid using the QuikChange® MultiSite-Directed Mutagenesis Kit (Stratagene, La Jolla, CA) to change G395 to A, resulting in the Arg to His mutation. R132C, R132L and R132S mutants were introduced into pET41a-ICHD1 in an analogous way.

Wild-type and mutant proteins were expressed in and purified from the *E. coli* Rosetta™ strain (Invitrogen, Carlsbad, CA) as follows. Cells were grown in LB (20 µg/ml Kanamycin) at 37°C with shaking until OD600 reaches 0.6. The temperature was changed to 18°C and protein expression was induced by adding IPTG to final concentration of 1 mM. After 12-16 hours of IPTG induction, cells were resuspended in Lysis Buffer (20mM Tris, pH7.4, 0.1% Triton X-100, 500 mM NaCl, 1 mM PMSF, 5 mM β-mercaptoethanol, 10 % glycerol) and disrupted by microfluidation. The 20,000g supernatant was loaded on metal chelate affinity resin (MCAC) equilibrated with Nickel Column Buffer A (20 mM Tris, pH7.4, 500mM NaCl, 5 mM β-mercaptoethanol, 10% glycerol) and washed for 20 column volumes. Elution from the column was effected by a 20 column-volume linear gradient of 10% to 100% Nickel Column Buffer B (20 mM Tris, pH7.4, 500 mM NaCl, 5 mM β-mercaptoethanol, 500 mM Imidazole, 10% glycerol) in Nickel Column Buffer A). Fractions containing the protein of interest were identified by SDS-PAGE, pooled, and dialyzed twice against a 200-volume excess of Gel Filtration Buffer (200 mM NaCl, 50 mM Tris 7.5, 5 mM β-mercaptoethanol, 2 mM MnSO<sub>4</sub>, 10% glycerol), then concentrated to 10 ml using Centricon (Millipore, Billerica, MA) centrifugal concentrators. Purification of active dimers was achieved by applying the concentrated eluent from the MCAC column to a Sephacryl S-200 (GE Life Sciences, Piscataway, NJ) column equilibrated with Gel Filtration Buffer and eluting the column with 20 column volumes of the same buffer. Fractions corresponding to the retention time of the dimeric protein were identified by SDS-PAGE and pooled for storage at -80°C.

**Cell lines and Cell Culture.** 293T cells were cultured in DMEM (Dulbecco's modified Eagles Medium) with 10% fetal bovine serum and were transfected using pCMV-6-based IDH-1 constructs in six-well plates with Fugene 6 (Roche) or Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions. Parental vector pCMV6 (no insert), pCMV6-wt IDH1 or pCMV6-R132H were transfected into human glioblastoma cell lines (U87MG; LN-18 (ATCC, HTB-14 and CRL-2610; respectively) cultured in DMEM with 10 % fetal bovine serum. Approximately 24 hrs after transfection, the cell cultures were transitioned to medium containing G418 sodium salt at concentrations of either 500 ug/ml (U87MG) or 750 ug/ml (LN-18) to select stable transfectants. Pooled populations of G418 resistant cells were generated and expression of either wild-type IDH1 or R132 IDH1 was confirmed by standard Western blot analysis.

**Western blot.** For transient transfection experiments in 293 cells, cells were lysed 72 hours after transfection with standard RIPA buffer. Lysates were separated by SDS-PAGE, transferred to nitrocellulose and probed with goat-anti-IDHc antibody (Santa Cruz Biotechnology sc49996) or rabbit-anti-MYC tag antibody (Cell Signaling Technology #2278) and then detected with HRP-conjugated donkey anti-goat or HRP-conjugated goat-anti-rabbit antibody (Santa Cruz Biotechnology sc2004). IDH1 antibody to confirm expression of both wild-type and R132H IDH1 was obtained from Proteintech. The IDH2 mouse monoclonal antibody used was obtained from Abcam.

**Detection of isocitrate,  $\alpha$ KG, and 2HG in purified enzyme reactions by LC-MS/MS.** Enzyme reactions performed as described in the text were run to completion as judged by measurement of the oxidation state of NADPH at 340 nm. Reactions were extracted with eight volumes of methanol, and centrifuged to remove precipitated protein. The supernatant was dried under a stream of nitrogen and resuspended in H<sub>2</sub>O. Analysis was conducted on an API2000 LC-MS/MS (Applied Biosystems, Foster City, CA). Sample separation and analysis was performed on a 150 x 2 mm, 4 uM Synergi Hydro-RP 80 A column, using a gradient of Buffer A (10

mM tributylamine, 15 mM acetic acid, 3% (v/v) methanol, in water) and Buffer B (methanol) using MRM transitions.

**Cell lysates based enzyme assays.** 293T cell lysates for measuring enzymatic activity were obtained 48 hours after transfection with M-PER lysis buffer supplemented with protease and phosphatase inhibitors. After lysates were sonicated and centrifuged at 12,000g, supernatants were collected and normalized for total protein concentration. To measure IDH oxidative activity, 3  $\mu$ g of lysate protein was added to 200  $\mu$ l of an assay solution containing 33 mM Tris-acetate buffer (pH 7.4), 1.3 mM  $MgCl_2$ , 0.33 mM EDTA, 100  $\mu$ M  $\beta$ -NADP, and varying concentrations of D-(+)-*threo*-isocitrate. Absorbance at 340 nm, reflecting NADPH production, was measured every 20 seconds for 30 min on a SpectraMax 190 spectrophotometer (Molecular Devices). Data points represent the mean activity of 3 replicates per lysate, averaged among 5 time points centered at every 5 min. To measure IDH reductive activity, 3  $\mu$ g of lysate protein was added to 200  $\mu$ l of an assay solution which contained 33 mM Tris-acetate (pH 7.4), 1.3 mM  $MgCl_2$ , 25  $\mu$ M  $\beta$ -NADPH, 40 mM  $NaHCO_3$ , and 0.6 mM  $\alpha$ KG. The decrease in 340 nm absorbance over time was measured to assess NADPH consumption, with 3 replicates per lysate.

**Recombinant IDH1 Enzyme Assays.** All reactions were performed in standard enzyme reaction buffer (150 mM NaCl, 20 mM Tris-Cl, pH 7.5, 10% glycerol, 5 mM  $MgCl_2$  and 0.03% (w/v) bovine serum albumin). For determination of kinetic parameters, sufficient enzyme was added to give a linear reaction for 1 to 5 seconds. Reaction progress was monitored by observation of the reduction state of the cofactor at 340 nm in an SFM-400 stopped-flow spectrophotometer (BioLogic, Knoxville, TN). Enzymatic constants were determined using curve fitting algorithms to standard kinetic models with the Sigmaplot software package (Systat Software, San Jose, CA).

**Determination of chirality of reaction products from enzyme reactions and tumors.** Enzyme reactions were run to completion and extracted with methanol as described above, then derivatized with enantiomerically pure tartaric acid before resolution and analysis by LC-MS/MS. After being thoroughly dried, samples were resuspended in freshly prepared 50 mg/ml (2*R*,3*R*)-(+)-Tartaric acid in dichloromethane:acetic acid (4:1) and incubated for 30 minutes at 75°C. After cooling

to room temperature, samples were briefly centrifuged at 14,000g, dried under a stream of nitrogen, and resuspended in H<sub>2</sub>O. Analysis was conducted on an API200 LC-MS/MS (Applied Biosystems, Foster City, CA), using an isocratic flow of 90:10 (2 mM ammonium formate, pH 3.6:MeOH) on a Luna C18(2) 150 x 2 mm, 5  $\mu$ M column. Tartaric-acid derivatized 2HG was detected using the 362.9/146.6 MRM transition and the following instrument settings: DP -1, FP -310, EP -4, CE-12, CXP-26. Analysis of the (R)-2HG standard, 2HG racemic mixture, and methanol-extracted tumor biomass (q.v.) was similarly performed.

**Crystallography conditions.** Crystals were obtained at 20°C by vapor diffusion equilibration using 3  $\mu$ L drops mixed 2:1 (protein:precipitant) against a well-solution of 100 mM MES pH 6.5, 20% PEG 6000.

**Protein characterization.** Approximately 90 mg of human cytosolic isocitrate dehydrogenase (HcIDH) was supplied to Xtal BioStructures by Agios. This protein was an engineered mutant form, R132S, with an 11-residue C-terminal affinity-purification tag (sequence SLEHHHHHHHH). The calculated monomeric molecular weight was 48.0 kDa and the theoretical pI was 6.50. The protein, at about 6 mg/mL concentration, was stored in 1-mL aliquots in 50 mM Tris-HCl (pH 7.4), 500 mM NaCl, 5 mM  $\beta$ -mercaptoethanol and 10% glycerol at -80°C. As shown in **FIG. 32A**, SDS-PAGE was performed to test protein purity and an anti-histidine Western blot was done to demonstrate the protein was indeed his-tagged. A sample of the protein was injected into an FPLC size-exclusion column to evaluate the sample purity and to determine the polymeric state in solution. **FIG. 32B** is a chromatogram of this run showing a single peak running at an estimated 87.6 kDa, suggesting IDH exists as a dimer at pH 7.4. Prior to crystallization, the protein was exchanged into 20 mM Tris-HCl (pH 7.4) and 100 mM NaCl using Amicon centrifugal concentrators. At this time, the protein was also concentrated to approximately 15 mg/mL. At this protein concentration and ionic strength, the protein tended to form a detectable level of precipitate. After spinning out the precipitate, the solution was stable at ~10 mg/mL at 4 °C.

**Initial attempts at crystallization.** The strategy for obtaining diffraction-quality crystals was derived from literature conditions, specifically “Structures of Human

Cytosolic NADP-dependent Isocitrate Dehydrogenase Reveal a Novel Self-regulatory Mechanism of Activity,” Xu, *et al.* (2005) *J.Biol.Chem.* 279: 33946-56. In this study, two crystal forms of HcIDH wildtype protein were produced. One contained their “binary complex”, IDH-NADP, which crystallized from hanging drops in the tetragonal space group  $P4_32_12$ . The drops were formed from equal parts of protein solution (15 mg/mL IDH, 10 mM NADP) and precipitant consisting of 100 mM MES (pH 6.5) and 12% PEG 20000. The other crystal form contained their “quaternary complex”, IDH-NADP/isocitrate/ $Ca^{2+}$ , which crystallized in the monoclinic space group  $P2_1$  using 100 mM MES (pH 5.9) and 20% PEG 6000 as the precipitant. Here they had added 10 mM DL-isocitrate and 10 mM calcium chloride to the protein solution. First attempts at crystallizing the R132S mutant in this study centered around these two reported conditions with little variation. The following lists the components of the crystallization that could be varied; several different combinations of these components were tried in the screening process.

In the protein solution:

HcIDH(R132S)	always ~10 mg/mL or ~0.2 mM
Tris-HCl (pH 7.4)	always 20 mM
NaCl	always 100 mM
NADP <sup>+</sup> /NADPH	absent or 5 mM NADP <sup>+</sup> (did not try NADPH)
DL-isocitric acid, trisodium salt	absent or 5 mM
calcium chloride	absent or 10 mM

In the precipitant: 100 mM MES (pH 6.5) and 12% PEG 20000

**OR**

100 mM MES (pH 6.0) and 20% PEG 6000

Drop size: always 3  $\mu$ L

Drop ratios: 2:1, 1:1 or 1:2 (protein:precipitant)

Upon forming the hanging drops, a milky precipitate was always observed. On inspection after 2-4 days at 20 °C most drops showed dense precipitation or phase separation. In some cases, the precipitate subsided and it was from these types of drops small crystals had grown, for example, as shown in **FIG. 33**.

**Crystal optimization.** Once bonafide crystals were achieved, the next step was to optimize the conditions to obtain larger and more regularly-shaped crystals of IDH-NADP/isocitrate/Ca<sup>2+</sup> in a timely and consistent manner. The optimal screen focused on varying the pH from 5.7 to 6.2, the MES concentration from 50 to 200 mM and the PEG 6000 concentration from 20 to 25%. Also, bigger drops were set up (5-6  $\mu$ l) and the drop ratios were again varied. These attempts failed to produce larger, diffraction-quality crystals but did reproduce the results reported above. Either a dense precipitate, oily phase separation or small crystals were observed.

**Using  $\alpha$ -Ketoglutarate.** Concurrent to the optimization of the isocitrate crystals, other screens were performed to obtain crystals of IDH(R132S) complexed with  $\alpha$ -ketoglutarate instead. The protein solution was consistently 10 mg/mL IDH in 20 mM Tris-HCl (pH 7.4) and 100 mM NaCl. The following were added in this order: 5 mM NADP, 5 mM  $\alpha$ -ketoglutaric acid (free acid, pH balanced with NaOH) and 10 mM calcium chloride. The protein was allowed to incubate with these compounds for at least an hour before the drops were set up. The precipitant was either 100 mM

MES (pH 6.5) and 12% PEG 20000 or 100 mM MES (pH 6.5) and 20% PEG 6000. Again, precipitation or phase separation was primarily seen, but in some drops small crystals did form. At the edge of one of the drops, a single large crystal formed, pictured below. This was the single crystal used in the following structure determination. **FIG. 34** shows crystal obtained from a protein solution contained 5 mM NADP, 5 mM  $\alpha$ -ketoglutarate, 10 mM Ca<sup>2+</sup>. Precipitant contained 100 mM MES (pH 6.5) and 12% PEG 20000.

**Cryo conditions.** In order to ship the crystal to the X-ray source and protect it during cryo-crystallography, a suitable cryo-protectant was needed. Glycerol is quite widely used and was the first choice. A cryo solution was made, basically as a mixture of the protein buffer and precipitant solution plus glycerol: 20 mM Tris-HCl (pH 7.5), 100 mM NaCl, 5 mM NADP, 5 mM  $\alpha$ -ketoglutaric acid, 10 mM calcium chloride, 100 mM MES (pH 6.5), 12% PEG 20000 and either 12.5% glycerol or 25% glycerol. The crystal was transferred to the cryo solution in two steps. First, 5  $\mu$ L of the 12.5% glycerol solution was added directly to the drop and incubated for 10 minutes, watching for possible cracking of the crystal. The liquid was removed from the drop and 10  $\mu$ L of the 25% glycerol solution was added on top of the crystal. Again, this incubated for 10 minutes, harvested into a nylon loop and plunged into liquid nitrogen. The crystal was stored submerged in a liquid nitrogen dewar for transport.

**Data collection and processing.** The frozen crystal was mounted on a Rigaku RAXIS IV X-ray instrument under a stream of nitrogen gas at temperatures near  $-170$  °C. A 200° dataset was collected with the image plate detector using 1.54 Å wavelength radiation from a rotating copper anode home source, 1° oscillations and 10 minute exposures. The presence of 25% glycerol as a cryoprotectant was sufficient for proper freezing, as no signs of crystal cracking (split spots or superimposed lattices) were observed. A diffuse ring was observed at 3.6 Å resolution, most likely caused by icing. The X-ray diffraction pattern showed clear lattice planes and reasonable spot separation, although the spacing along one reciprocal axis was rather small ( $b = 275.3$ ). The data was indexed to 2.7 Å resolution into space group  $P2_12_12$  with HKL2000 (Otwinowski and Minor, 1997). Three structures for HcIDH are known, designated the closed form (1T0L), the open form (1T09 subunit A) and semi-open form (1T09 subunit B). Molecular replacement was performed with the



CCP4 program PHASER (Bailey, 1994) using only the protein atoms from these three forms. Only the closed form yielded a successful molecular replacement result with 6 protein subunits in the asymmetric unit. The unit cell contains approximately 53.8% solvent.

**Model refinement.** Using the CCP4 program REFMAC5, rigid-body refinement was performed to fit each of the 6 IDH subunits in the asymmetric unit. This was followed by rigid-body refinement of the three domains in each protein subunit. Restrained refinement utilizing non-crystallographic symmetry averaging of related pairs of subunits yielded an initial structure with  $R_{cryst}$  of 33% and  $R_{free}$  of 42%. Model building and real-space refinement were performed using the graphics program COOT (Emsley and Cowtan, 2004). A difference map was calculated and this showed strong electron density into which six individual copies of the NADP ligand and calcium ion were manually fit with COOT. Density for the  $\alpha$ -ketoglutarate structure was less defined and was fit after the binding-site protein residues were fit using a  $2F_o - F_c$  composite omit map. Automated Ramachandran-plot optimization coupled with manual real-space density fitting was applied to improve the overall geometry and fit. A final round of restrained refinement with NCS yielded an  $R_{cryst}$  of 30.1% and  $R_{free}$  of 35.2%.

a, Å	b, Å	c, Å	$\alpha$	$\beta$	$\gamma$	Unit cell volume, Å <sup>3</sup>	Z
116.14	275.30	96.28	90°	90°	90°	$3.08 \times 10^6$	24

Reflections in working set / test set	68,755 / 3,608 (5.0%)
$R_{cryst}$	30.1%
$R_{free}$	35.2%

X-ray data and refinement statistics for IDH(R132S)-NADP/ $\alpha$ -ketoglurate/ $\text{Ca}^{2+}$ 

Crystal parameters	
Space group	<i>P2<sub>1</sub>2<sub>1</sub>2</i>
Unit cell dimensions	
a, b, c, Å	116.139, 275.297, 96.283
$\alpha$ , $\beta$ , $\gamma$ , °	90.0, 90.0, 90.0
Volume, Å <sup>3</sup>	3,078,440
No. protein molecules in asymmetric unit	6
No. protein molecules in unit cell, Z	24
Data collection	
Beam line	
Date of collection	Apr 25, 2009
$\lambda$ , Å	1.5418
Detector	Rigaku Raxis IV
Data set ( $\phi$ ), °	200
Resolution, Å	25-2.7 (2.8-2.7)
Unique reflections ( <i>N</i> , <i>F</i> > 0)	73,587
Completeness, %	85.4 (48.4)
$\langle I \rangle / \sigma I$	9.88 (1.83)

R-merge	0.109 (0.33)
Redundancy	4.3 (1.8)
Mosaicity	0.666
Wilson B factor	57.9
Anisotropy B factor, Å <sup>2</sup>	-1.96
Refinement Statistics	
Resolution limit, Å	20.02-2.70
No. of reflections used for R-work <sup>a</sup> / R-free <sup>b</sup>	68,755 / 3608
Protein atoms	19788
Ligand atoms	348
No. of waters	357
Ions etc.	6
Matthews coeff. Å <sup>3</sup> /Dalton	2.68
Solvent, %	53.8
R-work <sup>a</sup> / R-free <sup>b</sup> , (%)	30.1 / 35.2
Figure-of-merit <sup>c</sup>	0.80 (0.74)
Average B factors	31.0
Coordinates error (Luzzati plot), Å	0.484
R.M.S. deviations	

Bond lengths, Å	0.026
Bond angles, °	2.86

Completeness and *R*-merge are given for all data and for data in the highest resolution shell. Highest shell values are in parentheses.

<sup>a</sup>*R* factor =  $\sum_{hkl} |F_o - F_c| / \sum_{hkl} F_o$ , where  $F_o$  and  $F_c$  are the observed and calculated structure factor amplitudes, respectively for all reflections *hkl* used in refinement.

<sup>b</sup>*R*-free is calculated for 5% of the data that were not used in refinement.

<sup>c</sup>Figure of merit =  $\sqrt{x^2 + y^2}$ , where  $x = (\sum_0^{2\pi} P(\alpha) \cos \alpha) / (\sum_0^{2\pi} P(\alpha))$ ,  $y = (\sum_0^{2\pi} P(\alpha) \sin \alpha) / (\sum_0^{2\pi} P(\alpha))$ , and the phase probability  $P(\alpha) = \exp(A \cos \alpha + B \sin \alpha + C \cos(2\alpha) + D \sin(2\alpha))$ , where *A*, *B*, *C*, and *D* are the Hendrickson-Lattman coefficients and  $\alpha$  is the phase.

#### Stereochemistry of IDH(R132S)-NADP/ $\alpha$ -ketoglurate/Ca<sup>2+</sup>

Ramachandran plot statistics	No. of amino acids	% of Residues
Residues in most favored regions [A, B, L]	1824	82.2
Residues in additional allowed regions [a, b, l, p]	341	15.4
Residues in generously allowed regions [-a, -b, -l, -p]	38	1.7
Residues in disallowed regions	17	0.8
Number of non-glycine and non-proline residues	2220	100
Number of end-residues (excl. Gly and Pro)	387	
Number of glycine residues	198	
Number of proline residues	72	
Total number of residues	2877	
Overall <G> -factor <sup>d</sup> score (> -1.0)	-0.65	

Generated by PROCHECK (Laskowski RA, MacArthur MW, Moss DS, Thornton JM (1993) J Appl Crystallogr 26:283-291.)

<sup>d</sup> G-factors for main-chain and side-chain dihedral angles, and main-chain covalent forces (bond lengths and bond angles). Values should be ideally -0.5 or above -1.0.

Radiation wavelength, Å	1.54
Resolution, Å (outer shell)	20-2.70 (2.80-2.70)
Unique reflections	73,587
Completeness (outer shell)	85.4% (48.4%)
Redundancy (outer shell)	4.3 (1.8)
$R_{\text{merge}}$ (outer shell)	10.9% (33%)
$\langle I \rangle / \langle \sigma(I) \rangle$ (outer shell)	9.88 (1.83)

**Clinical Specimens, metabolite extraction and analysis.** Human brain tumors were obtained during surgical resection, snap frozen in isopentane cooled by liquid nitrogen and stored at -80 C. Clinical classification of the tissue was performed using standard clinical pathology categorization and grading as established by the WHO. Genomic sequence analysis was deployed to identify brain tumor samples containing either wild-type isocitrate dehydrogenase (IDH1) or mutations altering amino acid 132. Genomic DNA was isolated from 50-100 mgs of brain tumor tissue using standard methods. A polymerase chain reaction on the isolated genomic DNA was used to amplify a 295 base pair fragment of the genomic DNA that contains both the intron and 2<sup>nd</sup> exon sequences of human IDH1 and mutation status assessed by standard molecular biology techniques.

Metabolite extraction was accomplished by adding a 10x volume (m/v ratio) of -80 °C methanol:water mix (80%:20%) to the brain tissue (approximately 100mgs) followed by 30 s homogenization at 4 C. These chilled, methanol extracted homogenized tissues were then centrifuged at 14,000 rpm for 30 minutes to sediment the cellular and tissue debris and the cleared tissue supernatants were transferred to a screw-cap freezer vial and stored at -80 °C. For analysis, a 2X volume of tributylamine (10 mM) acetic acid (10 mM) pH 5.5 was added to the samples and analyzed by LCMS as follows. Sample extracts were filtered using a Millex-FG 0.20 micron disk and 10 µL were injected onto a reverse-phase HPLC column (Synergi 150mm x 2 mm,

Phenomenex Inc.) and eluted using a linear gradient LCMS-grade methanol (50% with 10 mM tributylamine and 10 mM acetic acid) ramping to 80 % methanol:10 mM tributylamine: 10 mM acetic acid over 6 minutes at 200  $\mu$ L/min. Eluted metabolite ions were detected using a triple-quadrupole mass spectrometer, tuned to detect in negative mode with multiple-reaction-monitoring mode transition set (MRM's) according to the molecular weights and fragmentation patterns for 8 known central metabolites, including 2-hydroxyglutarate as described above. Data was processed using Analyst Software (Applied Biosystems, Inc.) and metabolite signal intensities were obtained by standard peak integration methods.




EXAMPLE 9 COMPOUNDS THAT INHIBIT IDH1 R132H



Assays were conducted in a volume of 76  $\mu$ l assay buffer (150 mM NaCl, 10 mM MgCl<sub>2</sub>, 20 mM Tris pH 7.5, 0.03% bovine serum albumin) as follows in a standard 384-well plate: To 25  $\mu$ l of substrate mix (8  $\mu$ M NADPH, 2 mM aKG), 1  $\mu$ l of test compound was added in DMSO. The plate was centrifuged briefly, and then 25  $\mu$ l of enzyme mix was added (0.2  $\mu$ g/ml ICDH1 R132H) followed by a brief centrifugation and shake at 100 RPM. The reaction was incubated for 50 minutes at room temperature, then 25  $\mu$ l of detection mix (30  $\mu$ M resazurin, 36  $\mu$ g/ml ) was added and the mixture further incubated for 5 minutes at room temperature. The conversion of resazurin to resorufin was detected by fluorescent spectroscopy at Ex544 Em590 c/o 590.

**Table 24a** shows the wild type vs mutant selectivity profile of 5 examples of IDH1R132H inhibitors. The IDH1wt assay was performed at 1x Km of NADPH as opposed to IDHR132H at 10x or 100x Km of NADPH. The second example showed no inhibition, even at 100  $\mu$ M. Also, the first example has IC<sub>50</sub>=5.74  $\mu$ M but is shifted significantly when assayed at 100x Km, indicating direct NADPH-competitive inhibitor. The selectivity between wild type vs mutant could be >20-fold.



**Table 24a**

STRUCTURE	LDHa IC50	LDHb IC50	ICDH IC50 ( $\mu$ M) @ 4 $\mu$ M (10x Km) NADPH	ICDH IC50 ( $\mu$ M) @ 40 $\mu$ M NADPH	IC50 Ratio (40/4)	IDH1wt IC50 @ 1x Km ( $\mu$ M)






	25.43	64.07	5.74	>100	17.42	<b>16.22</b>
	5.92	17.40	12.26	41.40	3.38	<b>NO inhibition</b>
	8.61	>100	12.79	14.70	1.15	<b>19.23</b>






	33.75	>100	14.98	19.17	1.28	<b>46.83</b>
	12.76	>100	23.80	33.16	1.39	<b>69.33</b>






Additional exemplary compounds that inhibit IDH1R132H are provided below in Table 24b.






Compound	No.
	<b>1</b>
	<b>2</b>













Compound	No.
	3
	4
	5
	6
	7






Compound	No.
	8
	9
	10
	11
	12






Compound	No.
	13
	14
	15
	16
	17

Compound	No.
	18
	19
	20
	21
	22






Compound	No.
	23
	24
	25
	26
	27






Compound	No.
	28
	29
	30
	31
	32






Compound	No.
	33
	34
	35
	36
	37






Compound	No.
	38
	39
	40
	41
	42













Compound	No.
	43
	44
	45
	46
	47






Compound	No.
	48
	49
	50
	51
	52






Compound	No.
	53
	54
	55
	56
	57

Compound	No.
	58
	59
	60
	61
	62






Compound	No.
	63
	64
	65
	66
	67






Compound	No.
	68
	69
	70
	71
	72

Compound	No.
	73
	74
	75
	76
	77

Compound	No.
	78
	79
	80
	81
	82



Compound	No.
	83
	84
	85
	86
	87

Compound	No.
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C3)c4ccccc4F)</chem>	88
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C3)c4ccccc4C</chem>	89
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C3)c4ccccc4OC</chem>	90
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C3)c4ccccc4OC</chem>	91
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C3)c4ccccc4OC</chem>	92

EXAMPLE 10. The mutant enzyme IDH2-R172K has elevated NADPH reductive catalysis activity as compared to wildtype IDH2 enzyme.

NADPH reduction activity was measured for the enzymes IDH2-R172K, IDH2-wildtype, IDH1-R132H and IDH1-wildtype. The final reactant concentrations for each reaction were as follows: 20 mM Tris 7.5, 150 mM NaCl, 2 mM MnCl<sub>2</sub>, 10% glycerol, 0.03% BSA, enzyme (1-120 µg/mL), 1 mM NADPH, and 5 mM αKG (alpha ketoglutarate). The resulting specific activities (µmol/min/mg) are presented in the graph in FIG. 35. The results indicate that the mutant IDH2 has elevated reductive activity as compared to wildtype IDH2, even though both the mutant and wildtype IDH2 enzymes were able to make 2HG (2-hydroxyglutarate) at saturating levels of reactants αKG and NADPH.

EXAMPLE 11: 2-HG accumulates in AML with IDH1/2 mutations

Patients and clinical data

Peripheral blood and bone marrow were collected from AML patients at the time of diagnosis and at relapse, following REB approved informed consent. The cells were separated by ficol hypaque centrifugation, and stored at -150° C in 10% DMSO, 40% FCS and 50% alpha-MEM medium. Patient sera were stored at -80° C. Cytogenetics and molecular testing were performed in the diagnostic laboratory of the University Health Network (Toronto, Canada). A subgroup of patients (n=132) was given consistent initial treatment using a standard induction and consolidation chemotherapy regimen consisting of daunorubicin and cytarabine.

IDH1 and IDH2 Genotyping

DNA was extracted from leukemic cells and cell lines using the Qiagen Puregene kit (Valencia CA). For a subset of samples (n=96), RNA was extracted from leukemic cells using a Qiagen RNeasy kit, and reverse transcribed into cDNA for IDH1 and IDH2 genotyping. IDH1 and IDH2 genotype was determined at the Analytical Genetics Technology Centre at the University Health Network (Toronto, Canada) using a Sequenom MassARRAY™ platform (Sequenom, San Diego, CA). Positive results were confirmed by direct sequencing on an ABI PRISM 3130XL genetic analyzer (Applied Biosystems, Foster City, CA).

Cell lines

AML cell lines (OCI/AML-1, OCI/AML-2, OCI/AML-3, OCI/AML-4, OCI/AML-5, HL-60, MV-4-11, THP-1, K562, and KG1A) and 5637 cells were obtained from the laboratory of Mark Minden (Ontario Cancer Institute, Toronto, Canada). Primary AML cells were cultured in alpha-MEM media supplemented with 20% fetal bovine serum, and 10% 5637 cell conditioned media as previously described<sup>13</sup>. Growth curves were generated by counting viable cells as assessed by trypan blue exclusion on a Vi-CELL automated cell counter (Beckman Coulter, Fullerton, CA).

Expression / purification of IDH1 and IDH2 proteins

The human IDH1 cDNA (ref. ID NM\_005896) and IDH2 cDNA (ref. ID NM\_002168) were purchased from OriGene Technologies (Rockville, MD). For expression in *E. coli*, the coding region was amplified by PCR using primers designed to add NDEI and XHO1 restrictions sites at the 5' and 3' ends respectively. The resultant fragments for IDH1 (full length) and IDH2 (residues 40-452) were cloned into vector pET41a (EMD Biosciences, Madison, WI) to enable the *E. coli* expression of C-terminal His8-tagged protein. Site directed mutagenesis was performed on the pET41a-IDH1 and pET41a-IDH2 plasmid using the QuikChange® Lightning Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA) to change C394 to T in the IDH1 cDNA, resulting in the R132C mutation, and to change G515 to A in the IDH2 cDNA, resulting in the R172K mutation. Wild-type and mutant IDH1 proteins were expressed in and purified from the *E. coli* Rosetta™ (DE3) strain according to manufacturer's instructions (Invitrogen, Carlsbad, CA). Overexpression of IDH2 protein was accomplished by co-transfection of expression plasmids encoding respective IDH2 clones and pG-KJE8 expressing chaperone proteins.

IDH1/2 activity assays

Enzymatic activity was assessed by following the change in NADPH absorbance at 340 nm over time in an SFM-400 stopped-flow spectrophotometer (BioLogic, Knoxville, TN) in the presence of isocitrate and NADP<sup>+</sup> (forward reaction), or  $\alpha$ -KG and NADPH (reverse reaction). All reactions were performed in standard enzyme reaction buffer (150 mM NaCl, 20 mM Tris-Cl, pH 7.5, 10mM MgCl<sub>2</sub> and 0.03% (w/v) bovine serum albumin). For determination of kinetic parameters, sufficient enzyme was added to give a linear reaction for 1 to 5 seconds. Enzymatic binding constants were determined using curve fitting algorithms to standard kinetic models with the Sigmaplot software package (Systat Software, San Jose, CA). For determination of kcat, enzyme was incubated with 5X Km of substrate and cofactor;

consumption of NADPH or NADP was determined by a change in the OD<sub>340</sub> over time. In both cases an extinction coefficient of 6200 M<sup>-1</sup> cm<sup>-1</sup> was used for NADPH.

#### 2-HG and metabolite analysis

Metabolites were extracted from cultured cells, primary leukemic cells, and sera using 80% aqueous methanol (-80°C) as previously described. For cell extraction, frozen biopsies were thawed quickly at 37°C, and an aliquot of 2 million cells was spun down at 4 °C. The pellet was resuspended in -80°C 80% methanol. For serum extraction, 1 ml of serum was thawed quickly and mixed with 4 ml -80° C methanol. All extracts were spun at 13000 rpm at 4 °C to remove precipitate, dried at room temperature, and stored at -80° C until analysis by LC-MS. Metabolite levels (2-HG, α-KG, succinate, fumarate, and malate) were determined by ion paired reverse phase LC coupled to negative mode electrospray triple-quadropole MS using multiple reaction monitoring, and integrated elution peaks were compared with metabolite standard curves for absolute quantification as described.

#### Statistical analysis

Fisher's exact test was used to test for differences in categorical variables between IDH1/2 wt and IDH1/2 mutant patients. One way ANOVA followed by a student's t-test with correction for multiple comparisons was used to test for differences in IDH1 activity and metabolite concentrations. Differences with p<0.05 were considered significant.

### **Results**

In order to investigate the role of IDH1 R132 mutations in AML, leukemic cells obtained at initial presentation, from a series of 145 AML patients treated at the Princess Margaret Hospital with the aim of identifying mutant samples in our viable cell tissue bank were genotyped. Heterozygous IDH1 R132 mutations were found in 11 (8%) of these patients (Table 25). The spectrum of IDH1 mutations observed in AML appears to differ from that seen in CNS tumors. In the CNS, the majority of mutations (80-90%) are IDH1 R132H substitutions, whereas 5, 4, and 2 patients with IDH1 R132H, R132C, and R132G mutations, respectively (Table 25), were observed. In four cases, leukemic cells were also available from samples taken at the time of relapse. The IDH1 mutation was retained in 4/4 of these samples (Table 25). One of the patients harboring an IDH1 mutation had progressed to AML from an earlier

myelodysplastic syndrome (MDS). When cells from the prior MDS in this patient were analyzed, IDH1 was found to be wild-type. An additional 14 patients with MDS were genotyped, and all patients were found to be wild-type for IDH1, suggesting that IDH1 mutations are not a common feature of this disease. In samples from a subset of IDH1 mutant patients (n=8), reverse transcribed RNA was used for genotyping in order to assess the relative expression of mutant and wild-type alleles. Sequenom genotyping showed balanced allele peaks for these samples, indicating that both the wild-type and mutant genes are expressed. Ten established AML cell lines were also genotyped (OCI/AML-1, OCI/AML-2, OCI/AML-3, OCI/AML-4, OCI/AML-5, HL-60, MV-4-11, THP-1, K562, and KG1A) and none carried an IDH1 R132 mutation. Table 25: Identification of 13 AML patients bearing an IDH1 R132 or IDH2 R172 mutation\*

**Table 25**

Patient ID	Mutation	Amino acid change	FAB subtype	NPM1 and FLT3 status	Cytogenetic profile	Genotype at relapse	2-HG level (ng/2x10 <sup>6</sup> cells)
<b>IDH1 mutations</b>							
090108	G/A	R132H	M4	na	Normal	na	2090
090356	G/A	R132H	na	na	na	na	1529
0034	C/T	R132C	M5a	Normal	Normal	na	10285
0086	C/G	R132G	M2	Normal	Normal	na	10470
0488	C/T	R132C	M0	Normal	Normal	R132C	13822
8587	G/A	R132H	na	Normal	Normal	na	5742
8665	C/T	R132C	M1	na	Normal	na	7217
8741	G/A	R132H	M4	NPM1	Normal	R132H	6419
9544	C/G	R132G	na	na	Normal	R132G	4962
0174268	G/A	R132H	M1	NPM1	Normal	R132H	8464
090148	C/T	R132C	M1	na	46, xx, i(7)(p10) [20]	na	na
<b>IDH2 mutations</b>							
9382	G/A	R172K	M0	Normal	Normal	na	19247
0831	G/A	R172K	M1	Normal	Normal	na	15877

\* NPM1 denotes nucleophosmin 1, and FLT FMS-related tyrosine kinase 3. na indicates that some data was not available for some patients.

A metabolite screening assay to measure 2-HG in this set of AML samples was set up. Levels of 2-HG were approximately 50-fold higher in samples harboring an IDH1 R132 mutation (Table 25, Figure 36A, Table 26). 2-HG was also elevated in the sera of patients with IDH1 R132 mutant AML (Figure 36B). There was no relationship between the specific amino acid substitution at residue 132 of IDH1 and the level of 2-HG in this group of patients.

Table 26: Metabolite concentrations in individual IDH1/2 mutant and wild-type AML cells\*

Sample	IDH1/2 Genotype	2-HG (ng / 2x10 <sup>6</sup> cells)	$\alpha$ -KG (ng / 2x10 <sup>6</sup> cells)	Malate (ng / 2x10 <sup>6</sup> cells)	Fumarate (ng / 2x10 <sup>6</sup> cells)	Succinate (ng / 2x10 <sup>6</sup> cells)
0034	R132C	10285	125	192	239	2651
0086	R132G	10470	124	258	229	3043
0488	R132C	13822	95	184	193	2671
8587	R132H	5742	108	97	95	1409
8665	R132C	7217	137	118	120	1648
8741	R132H	6419	87	66	61	938
9544	R132G	4962	95	76	72	1199
0174268	R132H	8464	213	323	318	2287
090356	R132H	1529	138	657	366	1462
090108	R132H	2090	Na	246	941	3560
090148†	R132C	na	Na	na	Na	Na
8741‡	R132H	2890	131	113	106	1509
9554‡	R132G	7448	115	208	227	2658
0174268‡	R132H	964	72	134	138	2242
0488‡	R132C	7511	85	289	310	3448
9382	R172K	19247	790	821	766	5481
0831	R172K	15877	350	721	708	5144
157	Wild type	212	121	484	437	3057
202	Wild type	121	57	161	136	1443
205	Wild type	147	39	162	153	1011
209	Wild type	124	111	167	168	1610
239	Wild type	112	106	305	361	1436
277	Wild type	157	61	257	257	2029
291	Wild type	113	118	124	128	1240
313	Wild type	116	75	151	181	1541
090158	Wild type	411	217	658	647	3202
090156	Wild type	407	500	1276	1275	6091

\* IDH1/2 denotes isocitrate dehydrogenase 1 and 2, 2-HG 2-hydroxy glutarate, and  $\alpha$ -KG alpha-ketoglutarate. Metabolite measurements were not available for all patients.

† metabolic measurements were not made due to limited patient sample

‡ indicates samples obtained at relapse.

Two samples harboring wild-type IDH1 also showed high levels of 2-HG (Table 25). The high 2-HG concentration prompted sequencing of the IDH2 gene in these two AML samples, which established the presence of IDH2 R172K mutations in both samples (Table 25).

Evaluation of the clinical characteristics of patients with or without IDH1/2 mutations revealed a significant correlation between IDH1/2 mutations and normal karyotype ( $p=0.05$ ), but no other differences between these two groups (Table 27). Notably, there was no difference in treatment response for a subgroup of patients who received consistent treatment ( $n=136$ ). These findings are consistent with the initial report identifying IDH1 mutations in AML.

Table 27: Characteristics of IDH1/2 mutant and wild-type patients\*

Variable	IDH1/2 Wild-type (N=132)	IDH1/2 Mutant (N=13)	P Value
Age (yr)	58.8 ±16.2	52.6 ±7.0	0.17†
Sex (% male)	53 (70/132)	62 (8/13)	0.77‡
WBC at diagnosis ( $10^9$ cells/L)	40.7 ±50.6	28.7 ±34.1	0.38†
Initial treatment response (% complete remission)	70 (85/122)	62 (8/13)	0.54‡
Cytogenetic profile (% normal)	62 (72/117)	92 (11/12)	0.05‡
Additional mutations			
FLT3 (%)	17 (8/47)	0 (0/8)	0.58‡
NPM1 (%)	30 (14/47)	25(2/8)	1.0‡

\* For plus-minus values, the value indicates the mean, and ± indicates the standard deviation. IDH1/2 denotes isocitrate dehydrogenase 1 and 2, WBC white blood cell count, FLT3 FMS-related tyrosine kinase 3, and NPM1 nucleophosmin 1.

† P-value was calculated using the student's t-test.

‡ P-value was calculated using Fisher's exact test.

Panels of AML cells from wild-type and IDH1 mutant patients were cultured *in vitro*. There was no difference in the growth rates or viability of the IDH1 R132 mutant and wild-type cells, with both groups showing high variability in their ability to proliferate in culture, as is characteristic of primary AML cells (Figure 36C). There was no relationship between 2-HG levels in the IDH1 R132 mutant cells and their growth rate or viability in culture. After 14 days in culture, the mutant AML cells retained their



IDH1 R132 mutations (11/11), and continued to accumulate high levels of 2-HG (Figure 36A), further confirming that IDH1 R132 mutations lead to the production and accumulation of 2-HG in AML cells.

To investigate the effect of IDH1/2 mutations on the concentration of cellular metabolites proximal to the IDH reaction,  $\alpha$ -KG, succinate, malate, and fumarate levels were measured in AML cells with IDH1/2 mutations and in a set of wild-type AML cells matched for AML subtype and cytogenetic profile. None of the metabolites were found to be greatly altered in the IDH1 mutants compared to the IDH1 wild-type cells (Figure 27, Supplementary Table 26). The mean level of  $\alpha$ -KG was not altered in the IDH1/2 mutant AML cells, suggesting that the mutation does not decrease the concentration of this metabolite as has been previously hypothesized. To confirm that the R132C mutation of IDH1, and the R172K mutation of IDH2 confer a novel enzymatic activity that produces 2-HG, recombinant mutant enzymes were assayed for the NADPH-dependent reduction of  $\alpha$ -KG. When samples were analyzed by LC-MS upon completion of the enzyme assay, 2-HG was identified as the end product for both the IDH1 R132C and IDH2 R172K mutant enzymes (Figure 38). No isocitrate was detectable by LC-MS, indicating that 2-HG is the sole product of this reaction (Figure 38). This observation held true even when the reductive reaction was performed in buffer containing  $\text{NaHCO}_3$  saturated with  $\text{CO}_2$ .

A large proportion of IDH1 mutant patients in AML have an IDH1 R132C mutation (Table 25). In order to biochemically characterize mutant IDH1 R132C, the enzymatic properties of recombinant R132C protein were assessed *in vitro*. Kinetic analyses showed that the R132C substitution severely impairs the oxidative decarboxylation of isocitrate to  $\alpha$ -KG, with a significant decrease in  $k_{\text{cat}}$ , even though the affinity for the co-factor  $\text{NADP}^+$  remains essentially unchanged (Table 28). However, unlike the R132H mutant enzyme described previously the R132C mutation leads to a dramatic loss of affinity for isocitrate ( $K_M$ ), and a drop in net isocitrate metabolism efficiency ( $k_{\text{cat}}/K_M$ ) of more than six orders of magnitude (Table 28). This suggests a potential difference in the substrate-level regulation of enzyme activity in the context of AML. While substitution of cysteine at R132 inactivates the canonical conversion of isocitrate to  $\alpha$ -KG, the IDH1 R132C mutant enzyme acquires the ability to catalyze the reduction of  $\alpha$ -KG to 2-HG in an NADPH dependent manner (Figure 39). This reductive reaction of mutant IDH1 R132C is highly efficient ( $k_{\text{cat}}/K_M$ )

compared to the wild-type enzyme, due to the considerable increase in binding affinity of both the NADPH and  $\alpha$ -KG substrates ( $K_M$ ) (Table 28).

Table 28: Kinetic parameters of the IDH1 R132C mutant enzyme

<b>Oxidative (<math>\rightarrow</math> NADPH)</b>	<b>WT</b>	<b>R132C</b>
$K_{M,NADP^+}$ ( $\mu$ M)	49	21
$K_{M,isocitrate}$ ( $\mu$ M)	57	$8.7 \times 10^4$
$K_{M,MgCl_2}$ ( $\mu$ M)	29	$4.5 \times 10^2$
$K_{i,\alpha KG}$ ( $\mu$ M)	$6.1 \times 10^2$	61
$k_{cat}$ ( $s^{-1}$ )	$1.3 \times 10^5$	$7.1 \times 10^2$
$k_{cat} / K_{M,isoc}$ ( $M^{-1} \cdot s^{-1}$ )	$2.3 \times 10^9$	$8.2 \times 10^3$
<b>Reductive (<math>\rightarrow</math> NADP<sup>+</sup>)</b>	<b>WT</b>	<b>R132C</b>
$K_{M,NADPH}$ ( $\mu$ M)	n/a*	0.3
$K_{M,\alpha KG}$ ( $\mu$ M)	n/a	295
$k_{cat}$ ( $s^{-1}$ )	~ 7 (est.)	$5.5 \times 10^2$

\* n/a indicates no measureable activity

## **METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS**

### **CLAIM OF PRIORITY**

This application is a continuation of U.S.S.N. 13/939,519, filed July 11, 2013, which is a continuation of U.S.S.N. 13/256,396, filed November 29, 2011, which is a national stage application under 35 U.S.C. §371 of International Application No. PCT/US2010/027253, filed March 12, 2010, published as International Publication No. WO 2010/105243 on September 16, 2010, which claims priority to U.S.S.N. 61/160,253, filed March 13, 2009; U.S.S.N. 61/160,664, filed March 16, 2009; U.S.S.N. 61/173,518, filed April 28, 2009; U.S.S.N. 61/180,609, filed May 22, 2009; U.S.S.N. 61/220,543, filed June 25, 2009; U.S.S.N. 61/227,649, filed July 22, 2009; U.S.S.N. 61/229,689, filed July 29, 2009; U.S.S.N. 61/253,820, filed October 21, 2009; and U.S.S.N. 61/266,929, filed December 4, 2009, the contents of each of which are incorporated herein by reference.

### **FIELD OF THE INVENTION**

The invention relates to methods and compositions for evaluating and treating cell proliferation-related disorders, *e.g.*, proliferative disorders such as cancer.

### **BACKGROUND**

Isocitrate dehydrogenase, also known as IDH, is an enzyme which participates in the citric acid cycle. It catalyzes the third step of the cycle: the oxidative decarboxylation of isocitrate, producing alpha-ketoglutarate ( $\alpha$ -ketoglutarate or  $\alpha$ -KG) and CO<sub>2</sub> while converting NAD<sup>+</sup> to NADH. This is a two-step process, which involves oxidation of isocitrate (a secondary alcohol) to oxalosuccinate (a ketone), followed by the decarboxylation of the carboxyl group beta to the ketone, forming alpha-ketoglutarate. Another isoform of the enzyme catalyzes the same reaction; however this reaction is unrelated to the citric acid cycle, is carried out in the cytosol as well as the mitochondrion and peroxisome, and uses NADP<sup>+</sup> as a cofactor instead of NAD<sup>+</sup>.

### SUMMARY OF THE INVENTION

Methods and compositions disclosed herein relate to the role played in disease by neoactive products produced by neoactive mutant enzymes, e.g., mutant metabolic pathway enzymes. The inventors have discovered, *inter alia*, a neoactivity associated with IDH mutants and that the product of the neoactivity can be significantly elevated in cancer cells. Disclosed herein are methods and compositions for treating, and methods of evaluating, subjects having or at risk for a disorder, e.g., a cell proliferation-related disorder characterized by a neoactivity in a metabolic pathway enzyme, e.g., IDH neoactivity. Such disorders include e.g., proliferative disorders such as cancer. The inventors have discovered and disclosed herein novel therapeutic agents for the treatment of disorders, e.g., cancers, characterized by, e.g., by a neoactivity, neoactive protein, neoactive mRNA, or neoactive mutations. In embodiments a therapeutic agent reduces levels of neoactivity or neoactive product or ameliorates an effect of a neoactive product. Methods described herein also allow the identification of a subject, or identification of a treatment for the subject, on the basis of neoactivity genotype or phenotype. This evaluation can allow for optimal matching of subject with treatment, e.g., where the selection of subject, treatment, or both, is based on an analysis of neoactivity genotype or phenotype. E.g., methods describe herein can allow selection of a treatment regimen comprising administration of a novel compound, e.g., a novel compound disclosed herein, or a known compound, e.g., a known compound not previously recommended for a selected disorder. In embodiments the known compound reduces levels of neoactivity or neoactive product or ameliorates an effect of a neoactive product. Methods described herein can guide and provide a basis for selection and administration of a novel compound or a known compound, or combination of compounds, not previously recommended for subjects having a disorder characterized by a somatic neoactive mutation in a metabolic pathway enzyme. In embodiments the neoactive genotype or phenotype can act as a biomarker the presence of which indicates that a compound, either novel, or previously known, should be administered, to treat a disorder characterized by a somatic neoactive mutation in a metabolic pathway enzyme. Neoactive mutants of IDH1 having a neoactivity that results in the production of 2-hydroxyglutarate, e.g., R-2-hydroxyglutarate and associated disorders are discussed in detail herein. They are exemplary, but not limiting, examples of embodiments of the invention.

While not wishing to be bound by theory it is believed that the balance between the production and elimination of neoactive product, e.g., 2HG, e.g., R-2HG, is important in disease. Neoactive mutants, to varying degrees for varying mutations, increase the level of neoactive product, while other processes, e.g., in the case of 2HG, e.g., R-2HG, enzymatic degradation of 2HG, e.g., by 2HG dehydrogenase, reduce the level of neoactive product. An incorrect balance is associated with disease. In embodiments, the net result of a neoactive mutation at IDH1 or IDH2 result in increased levels, in affected cells, of neoactive product, 2HG, e.g., R-2HG,

Accordingly, in one aspect, the invention features, a method of treating a subject having a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder. The cell proliferation-related disorder is characterized by a somatic mutation in a metabolic pathway enzyme. The mutation is associated with a neoactivity that results in the production of a neoactivity product. The method comprises: administering to the subject a therapeutically effective amount of a therapeutic agent described herein, e.g., a therapeutic agent that decreases the level of neoactivity product encoded by a selected or mutant somatic allele, e.g., an inhibitor of a neoactivity of the metabolic pathway enzyme (the neoactive enzyme), a therapeutic agent that ameliorates an unwanted affect of the neoactivity product, or a nucleic acid based inhibitor, e.g., a dRNA which targets the neoactive enzyme mRNA, to thereby treat the subject.

In an embodiment the subject is a subject not having, or not diagnosed as having, 2-hydroxyglutaric aciduria.

In an embodiment the subject has a cell proliferation-related disorder, e.g., a cancer, characterized by the neoactivity of the metabolic pathway enzyme encoded by selected or mutant allele.

In an embodiment the subject has a cell proliferation-related disorder, e.g., a cancer, characterized by the product formed by the neoactivity of the metabolic pathway enzyme encoded by selected or mutant allele.

In one embodiment, the metabolic pathway is selected from a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, nucleotide biosynthetic pathways, or the fatty acid biosynthetic pathway.

In an embodiment the therapeutic agent is a therapeutic agent described herein.

In an embodiment the method comprises selecting a subject on the basis of having a cancer characterized by the selected or mutant allele, the neoactivity, or an elevated level of neoactivity product.

In an embodiment the method comprises selecting a subject on the basis of having a cancer characterized by the product formed by the neoactivity of the protein encoded by selected or mutant allele, *e.g.*, by the imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI (magnetic resonance imaging) and/or MRS (magnetic resonance spectroscopy), to determine the presence, distribution or level of the product of the neoactivity, *e.g.*, in the case of an IDH1 allele described herein, 2-hydroxyglutarate (sometimes referred to herein as 2HG), *e.g.*, R-2-hydroxyglutarate (sometimes referred to herein as R-2HG).

In an embodiment the method comprises confirming or determining, *e.g.*, by direct examination or evaluation of the subject, or sample *e.g.*, tissue, product (*e.g.*, feces, sweat, semen, exhalation, hair or nails), or bodily fluid (*e.g.*, blood (*e.g.*, blood plasma), urine, lymph, or cerebrospinal fluid or other sample sourced disclosed herein) therefrom, (*e.g.*, by DNA sequencing, immuno analysis, or assay for enzymatic activity), or receiving such information about the subject, that the cancer is characterized by the selected or mutant allele.

In an embodiment the method comprises confirming or determining, *e.g.*, by direct examination or evaluation of the subject, the level of neoactivity or the level of the product of the neoactivity, or receiving such information about the subject. In an embodiment the presence, distribution or level of the product of the neoactivity, *e.g.*, in the case of an IDH1 allele described herein, 2HG, *e.g.*, R-2HG, is determined non-invasively, *e.g.*, by imaging methods, *e.g.*, by magnetic resonance-based methods.

In an embodiment the method comprises administering a second anti-cancer agent or therapy to the subject, *e.g.*, surgical removal or administration of a chemotherapeutic.

In another aspect, the invention features, a method of treating a subject having a cell proliferation-related disorder, *e.g.*, a precancerous disorder, or cancer. In an embodiment the subject does not have, or has not been diagnosed as having, 2-hydroxyglutaric aciduria. The cell proliferation-related disorder is characterized by a somatic allele, *e.g.*, a preselected allele, or mutant allele, of an IDH, *e.g.*, IDH1 or IDH2, which encodes a mutant IDH, *e.g.*, IDH1 or IDH2, enzyme having a neoactivity.

In embodiments the neoactivity is alpha hydroxy neoactivity. As used herein, alpha hydroxy neoactivity refers to the ability to convert an alpha ketone to an alpha hydroxy. In embodiments alpha hydroxy neoactivity proceeds with a reductive cofactor, e.g., NADPH or NADH. In embodiments the alpha hydroxyl neoactivity is 2HG neoactivity. 2HG neoactivity, as used herein, refers to the ability to convert alpha ketoglutarate to 2-hydroxyglutarate (sometimes referred to herein as 2HG), e.g., R-2-hydroxyglutarate (sometimes referred to herein as R-2HG). In embodiments 2HG neoactivity proceeds with a reductive cofactor, e.g., NADPH or NADH. In an embodiment a neoactive enzyme, e.g., an alpha hydroxyl, e.g., a 2HG, neoactive enzyme, can act on more than one substrate, e.g., more than one alpha hydroxy substrate.

The method comprises administering to the subject an effective amount of a therapeutic agent of type described herein to thereby treat the subject.

In an embodiment the therapeutic agent: results in lowering the level of a neoactivity product, e.g., an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the method comprises administering a therapeutic agent that lowers neoactivity, e.g., 2HG neoactivity. In an embodiment the method comprises administering an inhibitor of a mutant IDH protein, e.g., a mutant IDH1 or mutant IDH2 protein, having a neoactivity, e.g., alpha hydroxy neoactivity, e.g., 2HG neoactivity.

In an embodiment the therapeutic agent comprises a compound from Table 24a or Table 24b or a compound having the structure of Formula (X) or (Formula (XI) described herein.

In an embodiment the therapeutic agent comprises nucleic acid-based therapeutic agent, e.g., a dsRNA, e.g., a dsRNA described herein.

In an embodiment the the therapeutic agent is an inhibitor, e.g., a polypeptide, peptide, or small molecule (e.g., a molecule of less than 1,000 daltons), or aptomer, that binds to an IDH1 mutant or wildtype subunit and inhibits neoactivity, e.g., by inhibiting formation of a dimer, e.g., a homodimer of mutant IDH1 subunits or a heterodimer of a mutant and a wildtype subunit. In an embodiment the inhibitor is a polypeptide. In an embodiment the polypeptide acts as a dominant negative with respect to the neoactivity of the mutant enzyme. The polypeptide can correspond to full length IDH1 or a fragment thereof. The polypeptide need not be identical with

the corresponding residues of wildtype IDH1, but in embodiments has at least 60, 70, 80, 90 or 95 % homology with wildtype IDH1.

In an embodiment the therapeutic agent decreases the affinity of an IDH, *e.g.*, IDH1 or IDH2 neoactive mutant protein for NADH, NADPH or a divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , or decreases the levels or availability of NADH, NADPH or divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , *e.g.*, by competing for binding to the mutant enzyme. In an embodiment the enzyme is inhibited by replacing  $Mg^{2+}$  or  $Mn^{2+}$  with  $Ca^{2+}$ .

In an embodiment the therapeutic agent is an inhibitor that reduces the level a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the level of the product of a mutant having a neoactivity of an IDH, *e.g.*, IDH1 or IDH2 mutant, *e.g.*, it reduces the level of 2HG, *e.g.*, R-2HG.

In an embodiment the therapeutic agent is an inhibitor that:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that is selected on the basis that it:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH1, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of a mutant IDH, *e.g.*, IDH1 or IDH2, protein or mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2, protein.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of neoactive enzyme activity, *e.g.*, by interacting with, *e.g.*, binding to, mutant IDH, *e.g.*, IDH1 or IDH2, protein. In an embodiment the inhibitor is other than an antibody.

In an embodiment the therapeutic agent is an inhibitor that is a small molecule



and interacts with, *e.g.*, binds, the mutant RNA, *e.g.*, mutant IDH1 or IDH2 mRNA (*e.g.*, mutant IDH1 mRNA).

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, binds, either the mutant IDH, *e.g.*, IDH1 or IDH2, protein or interacts directly with, *e.g.*, binds, the mutant IDH mRNA, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the IDH is IDH1 and the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. Mutations in IDH1 associated with 2HG neoactivity include mutations at residue 132, *e.g.*, R132H, R132C, R132S, R132G, R132L, or R132V (*e.g.*, R132H or R132C).

In an embodiment the IDH is IDH2 and the neoactivity of the IDH2 mutant is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. Mutations in IDH2 associated with 2HG neoactivity include mutations at residue 172, *e.g.*, R172K, R172M, R172S, R172G, or R172W.

Treatment methods described herein can comprise evaluating a neoactivity genotype or phenotype. Methods of obtaining and analyzing samples, and the *in vivo* analysis in subjects, described elsewhere herein, *e.g.*, in the section entitled, “Methods of evaluating samples and/or subjects,” can be combined with this method.

In an embodiment, prior to or after treatment, the method includes evaluating the growth, size, weight, invasiveness, stage or other phenotype of the cell proliferation-related disorder.

In an embodiment, prior to or after treatment, the method includes evaluating the IDH, *e.g.*, IDH1 or IDH2, alpha hydroxyl neoactivity genotype, *e.g.*, 2HG, genotype, or alpha hydroxy neoactivity phenotype, *e.g.*, 2HG, *e.g.*, R-2HG, phenotype. Evaluating the alpha hydroxyl, *e.g.*, 2HG, genotype can comprise determining if an IDH1 or IDH2 mutation having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, is present, *e.g.*, a mutation disclosed herein having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. Alpha hydroxy neoactivity phenotype, *e.g.*, 2HG, *e.g.*, R-2HG, phenotype, as used herein, refers to the level of alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, level of alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, or level of mutant enzyme having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity (or corresponding mRNA). The evaluation can be by a method described herein.

In an embodiment the subject can be evaluated, before or after treatment, to determine if the cell proliferation-related disorder is characterized by an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment a cancer, *e.g.*, a glioma or brain tumor in a subject, can be analyzed, *e.g.*, by imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS, *e.g.*, before or after treatment, to determine if it is characterized by presence of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method comprises evaluating, *e.g.*, by direct examination or evaluation of the subject, or a sample from the subject, or receiving such information about the subject, the IDH, *e.g.*, IDH1 or IDH2, genotype, or an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG phenotype of, the subject, *e.g.*, of a cell, *e.g.*, a cancer cell, characterized by the cell proliferation-related disorder. (As described in more detail elsewhere herein the evaluation can be, *e.g.*, by DNA sequencing, immuno analysis, evaluation of the presence, distribution or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, from spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS measurement, sample analysis such as serum or spinal cord fluid analysis, or by analysis of surgical material, *e.g.*, by mass-spectroscopy). In embodiments this information is used to determine or confirm that a proliferation-related disorder, *e.g.*, a cancer, is characterized by an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In embodiments this information is used to determine or confirm that a cell proliferation-related disorder, *e.g.*, a cancer, is characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having a mutation, *e.g.*, a His, Ser, Cys, Gly, Val, Pro or Leu (*e.g.*, His, Ser, Cys, Gly, Val, or Leu at residue 132, more specifically, His or Cys, or an IDH2 allele having a mutation at residue 172, *e.g.*, a K, M, S, G, or W).

In an embodiment, before and/or after treatment has begun, the subject is evaluated or monitored by a method described herein, *e.g.*, the analysis of the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, to select, diagnose or prognose the subject, to select an inhibitor, or to evaluate response to the treatment or progression of disease.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, a leukemia, *e.g.*, AML or ALL, *e.g.*, B-ALL or T-ALL, prostate cancer, fibrosarcoma, paraganglioma, or myelodysplasia or myelodysplastic syndrome (*e.g.*, B-ALL or T-ALL, prostate cancer, or myelodysplasia or myelodysplastic syndrome) and the evaluation is: evaluation of the presence, distribution, or level of an alpha

hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG; or evaluation of the presence, distribution, or level of a neoactivity, *e.g.*, an alpha hydroxy neoactivity, e.g., 2HG neoactivity, of an IDH1 or IDH2, mutant protein.

In an embodiment the disorder is other than a solid tumor. In an embodiment the disorder is a tumor that, at the time of diagnosis or treatment, does not have a necrotic portion. In an embodiment the disorder is a tumor in which at least 30, 40, 50, 60, 70, 80 or 90% of the tumor cells carry an IHD, e.g., IDH1 or IDH2, mutation having 2HG neoactivity, at the time of diagnosis or treatment.

In an embodiment the cell proliferation-related disorder is a cancer, e.g., a cancer described herein, characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by unwanted (i.e., increased) levels of an alpha hydroxy neoactivity, product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, *e.g.*, wherein the tumor is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, *e.g.*, a mutant described herein. Gliomas include astrocytic tumors, oligodendroglial tumors, oligoastrocytic tumors, anaplastic astrocytomas, and glioblastomas. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu (e.g., His, Ser, Cys, Gly, Val, or Leu), or any residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **Fig. 21**). In an embodiment the allele encodes an IDH1 having His at residue 132. In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the IDH1 allele has an A (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394A, a C394G, a C394T,

a G395C, a G395T or a G395A mutation; specifically a C394A or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8), more specifically His, Ser, Cys, Gly, Val, or Leu; or His or Cys.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8), more specifically His, Ser, Cys, Gly, Val, or Leu; or His or Cys.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method comprises selecting a subject having a fibrosarcoma or paraganglioma wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having a fibrosarcoma or paraganglioma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having a fibrosarcoma or paraganglioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is localized or metastatic prostate cancer, *e.g.*, prostate adenocarcinoma, *e.g.*, wherein the cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

*E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al*, 2009, Int. J. Cancer, 125: 353-355 at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (*e.g.*, His, Ser, Cys,

Gly, Val, or Leu). In an embodiment the allele encodes an IDH1 having His or Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394T or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, wherein the cancer is characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a hematological cancer, *e.g.*, a leukemia, *e.g.*, AML, or ALL, wherein the hematological cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

In an embodiment the cell proliferation-related disorder is acute lymphoblastic leukemia (*e.g.*, an adult or pediatric form), *e.g.*, wherein the acute lymphoblastic leukemia (sometimes referred to herein as ALL) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. The ALL can be, *e.g.*, B-ALL or T-ALL. In an embodiment the cancer is characterized by increased levels of 2 an alpha hydroxy neoactivity product, *e.g.*, HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID

NO:8 (see also **FIG. 21**), more specifically His, Ser, Cys, Gly, Val, or Leu. In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, characterized by an IDH1 allele described herein, e.g., an IDH1 allele having Cys at residue 132 according to the sequence of SEQ ID NO:8.

In an embodiment the method comprises selecting a subject ALL, e.g., B-ALL or T-ALL, on the basis of cancer being characterized by having an IDH1 allele described herein, e.g., an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the cell proliferation-related disorder is acute myelogenous leukemia (e.g., an adult or pediatric form), e.g., wherein the acute myelogenous leukemia (sometimes referred to herein as AML) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, e.g., a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type. E.g., in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). E.g., the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**). In an embodiment the allele encodes an IDH1 having Cys, His or Gly at residue 132, more specifically, Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplasmic leukemia (AML) characterized by an IDH1 allele described herein, e.g., an IDH1 allele having Cys, His, or Gly at residue 132 according to the sequence of SEQ ID NO:8, more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML) on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His, or Gly at residue 132 (SEQ ID NO:8), more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML), on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method further comprises evaluating the subject for the presence of a mutation in the NRAS or NPMc gene.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, *e.g.*, wherein the myelodysplasia or myelodysplastic syndrome is characterized by having an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the disorder is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), more specifically His, Ser, Cys, Gly, Val, or Leu. In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His, or Gly at residue 132 according to the sequence of SEQ ID NO:8, more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His, or Gly at residue 132 (SEQ ID NO:8), more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being

characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the cell proliferation-related disorder is a glioma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, e.g., 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH2 allele described herein, e.g., an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), more specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH2 allele described herein, e.g., an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), more specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the cell proliferation-related disorder is a prostate cancer, e.g., prostate adenocarcinoma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, e.g., 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a prostate cancer, e.g., prostate adenocarcinoma, wherein the cancer is characterized by having



an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is ALL, *e.g.*, B-ALL or T-ALL, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**). In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is AML, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys,

Gly, Met, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172. In an embodiment the allele encodes an IDH2 having Gly at residue 172.

In an embodiment the method comprises selecting a subject having AML, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly or Met at residue 172 (SEQ ID NO:10), more specifically Lys or Met.

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, or Met at residue 172 (SEQ ID NO:10), more specifically Lys or Met.

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, characterized by a mutation, or preselected allele, of IDH2. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys, Gly, Met, Trp or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172. In an embodiment the allele encodes an IDH2 having Gly at residue 172.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, or Met at residue 172 (SEQ ID NO:10), in specific embodiments, Lys or Met.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, or Met at residue 172 (SEQ ID NO:10), in specific embodiments, Lys or Met.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being

characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment a product of the neoactivity is 2HG (e.g., R-2HG) which acts as a metabolite. In another embodiment a product of the neoactivity is 2HG (e.g., R-2HG) which acts as a toxin, e.g., a carcinogen.

In some embodiments, the methods described herein can result in reduced side effects relative to other known methods of treating cancer.

Therapeutic agents and methods of subject evaluation described herein can be combined with other therapeutic modalities, e.g., with art-known treatments.

In an embodiment the method comprises providing a second treatment, to the subject, e.g., surgical removal, irradiation or administration of a chemotherapeutic agent, e.g., an administration of an alkylating agent. Administration (or the establishment of therapeutic levels) of the second treatment can: begin prior to the beginning or treatment with (or prior to the establishment of therapeutic levels of) the inhibitor; begin after the beginning or treatment with (or after the establishment of therapeutic levels of) the inhibitor, or can be administered concurrently with the inhibitor, e.g., to achieve therapeutic levels of both concurrently.

In an embodiment the cell proliferation-related disorder is a CNS tumor, e.g., a glioma, and the second therapy comprises administration of one or more of: radiation; an alkylating agent, e.g., temozolomide, e.g., Temoader®, or BCNU; or an inhibitor of HER1/EGFR tyrosine kinase, e.g., erlotinib, e.g., Tarceva®.

The second therapy, e.g., in the case of glioma, can comprise implantation of BCNU or carmustine in the brain, e.g., implantation of a Gliadel® wafer.

The second therapy, e.g., in the case of glioma, can comprise administration of imatinib, e.g., Gleevec®.

In an embodiment the cell proliferation-related disorder is prostate cancer and the second therapy comprises one or more of: androgen ablation; administration of a microtubule stabilizer, e.g., docetaxol, e.g., Taxotere®; or administration of a topoisomerase II inhibitor, e.g., mitoxantrone.

In an embodiment the cell proliferation-related disorder is ALL, e.g., B-ALL or T-ALL, and the second therapy comprises one or more of:

induction phase treatment comprising the administration of one or more of: a steroid; an inhibitor of microtubule assembly, e.g., vincristine; an agent that reduces

the availability of asparagine, e.g., asparaginase; an anthracycline; or an antimetabolite, e.g., methotrexate, e.g., intrathecal methotrexate, or 6-mercaptopurine;

consolidation phase treatment comprising the administration of one or more of: a drug listed above for the induction phase; an antimetabolite, e.g., a guanine analog, e.g., 6-thioguanine; an alkylating agent, e.g., cyclophosphamide; an anti-metabolite, e.g., AraC or cytarabine; or an inhibitor of topoisomerase I, e.g., etoposide; or

maintenance phase treatment comprising the administration of one or more of the drugs listed above for induction or consolidation phase treatment.

In an embodiment the cell proliferation-related disorder is AML and the second therapy comprises administration of one or more of: an inhibitor of topoisomerase II, e.g., daunorubicin, idarubicin, topotecan or mitoxantrone; an inhibitor of topoisomerase I, e.g., etoposide; or an anti-metabolite, e.g., AraC or cytarabine.

In another aspect, the invention features, a method of evaluating, e.g. diagnosing, a subject, e.g., a subject not having, or not diagnosed as having, 2-hydroxyglutaric aciduria. The method comprises analyzing a parameter related to the neoactivity genotype or phenotype of the subject, e.g., analyzing one or more of:

a) the presence, distribution, or level of a neoactive product, e.g., the product of an alpha hydroxy neoactivity, e.g., 2HG, e.g., R-2HG, e.g., an increased level of product, 2HG, e.g., R-2HG (as used herein, an increased level of a product of an alpha hydroxy neoactivity, e.g., 2HG, e.g., R-2HG, or similar term, e.g., an increased level of neoactive product or neoactivity product, means increased as compared with a reference, e.g., the level seen in an otherwise similar cell lacking the IDH mutation, e.g., IDH1 or IDH2 mutation, or in a tissue or product from a subject not having the mutation (the terms increased and elevated as referred to the level of a product of alpha hydroxyl neoactivity as used herein, are used interchangeably);

b) the presence, distribution, or level of a neoactivity, e.g., alpha hydroxy neoactivity, e.g., 2HG neoactivity, of an IDH1 or IDH2, mutant protein;

c) the presence, distribution, or level of a neoactive mutant protein, e.g., an IDH, e.g., an IDH1 or IDH2, mutant protein which has a neoactivity, e.g., alpha hydroxy neoactivity, e.g., 2HG neoactivity, or a corresponding RNA; or

d) the presence of a selected somatic allele or mutation conferring neoactivity, e.g., an IDH, e.g., IDH1 or IDH2, which encodes a protein with a neoactivity, e.g.,

alpha hydroxy neoactivity, e.g., 2HG neoactivity, e.g., an allele disclosed herein, in cells characterized by a cell proliferation-related disorder from the subject, thereby evaluating the subject.

In an embodiment analyzing comprises performing a procedure, e.g., a test, to provide data or information on one or more of a-d, e.g., performing a method which results in a physical change in a sample, in the subject, or in a device or reagent used in the analysis, or which results in the formation of an image representative of the data. Methods of obtaining and analyzing samples, and the in vivo analysis in subjects, described elsewhere herein, e.g., in the section entitled, "Methods of evaluating samples and/or subjects," can be combined with this method. In another embodiment analyzing comprises receiving data or information from such test from another party. In an embodiment the analyzing comprises receiving data or information from such test from another party and, the method comprises, responsive to that data or information, administering a treatment to the subject.

As described herein, the evaluation can be used in a number of applications, e.g., for diagnosis, prognosis, staging, determination of treatment efficacy, patent selection, or drug selection.

Thus, in an embodiment method further comprises, e.g., responsive to the analysis of one or more of a-d:

diagnosing the subject, e.g., diagnosing the subject as having a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder;

staging the subject, e.g., determining the stage of a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder;

providing a prognosis for the subject, e.g., providing a prognosis for a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder;

determining the efficacy of a treatment, e.g., the efficacy of a chemotherapeutic agent, irradiation or surgery;

determining the efficacy of a treatment with a therapeutic agent, e.g., an inhibitor, described herein;

selecting the subject for a treatment for a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a

precancerous disorder. The selection can be based on the need for a reduction in neoactivity or on the need for amelioration of a condition associated with or resulting from neoactivity. For example, if it is determined that the subject has a cell proliferation-related disorder, *e.g.*, *e.g.*, cancer, or a precancerous disorder characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, or by a mutant IDH1 or IDH2, having alpha hydroxyl neoactivity, *e.g.*, 2HG, neoactivity, selecting the subject for treatment with a therapeutic agent described herein, *e.g.*, an inhibitor (*e.g.*, a small molecule or a nucleic acid-based inhibitor) of the neoactivity of that mutant (*e.g.*, conversion of alpha-ketoglutarate to 2HG, *e.g.*, R-2HG);

correlating the analysis with an outcome or a prognosis;

providing a value for an analysis on which the evaluation is based, *e.g.*, the value for a parameter correlated to the presence, distribution, or level of an alpha hydroxyl neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG;

providing a recommendation for treatment of the subject; or

memorializing a result of, or output from, the method, *e.g.*, a measurement made in the course of performing the method, and optionally transmitting the memorialization to a party, *e.g.*, the subject, a healthcare provider, or an entity that pays for the subject's treatment, *e.g.*, a government, insurance company, or other third party payer.

As described herein, the evaluation can provide information on which a number of decisions or treatments can be based.

Thus, in an embodiment the result of the evaluation, *e.g.*, an increased level of an alpha hydroxyl neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, the presence of an IDH, *e.g.*, IDH1 or IDH2, neoactivity, *e.g.*, alpha hydroxyl neoactivity, *e.g.*, 2HG neoactivity, the presence of an IDH, *e.g.*, IDH1 or IDH2, mutant protein (or corresponding RNA) which has alpha hydroxyl neoactivity, *e.g.*, 2HG neoactivity, the presence of a mutant allele of IDH, *e.g.*, IDH1 or IDH2, having alpha hydroxyl neoactivity, 2HG neoactivity, *e.g.*, an allele disclosed herein, is indicative of:

a cell proliferation-related disorder, *e.g.*, cancer, *e.g.*, it is indicative of a primary or metastatic lesion;

the stage of a cell proliferation-related disorder;

a prognosis or outcome for a cell proliferation-related disorder, *e.g.*, it is indicative of a less aggressive form of the disorder, *e.g.*, cancer. *E.g.*, in the case of

glioma, presence of an alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG, can indicate a less aggressive form of the cancer;

the efficacy of a treatment, e.g., the efficacy of a chemotherapeutic agent, irradiation or surgery;

the need of of a therapy disclosed herein, e.g., inhibition a neoactivity of an IDH, e.g., IDH1 or IDH2, neoactive mutant described herein. In an embodiment relatively higher levels (or the presence of the mutant) is correlated with need of inhibition a neoactivity of an IDH, e.g., IDH1 or IDH2, mutant described herein; or

responsiveness to a treatment. The result can be used as a noninvasive biomarker for clinical response. E.g., elevated levels can be predictive on better outcome in glioma patients (e.g., longer life expectancy).

As described herein, the evaluation can provide for the selection of a subject.

Thus, in an embodiment the method comprises, e.g., responsive to the analysis of one or more of a-d, selecting a subject, e.g., for a treatment. The subject can be selected on a basis described herein, e.g., on the basis of:

said subject being at risk for, or having, higher than normal levels of an alpha hydroxy neoactivity product, e.g., 2-hydroxyglutarate (e.g., R-2HG) in cell having a cell proliferation-related disorder, e.g., a leukemia such as AML or ALL, e.g., B-ALL or T-ALL, or a tumor lesion, e.g., a glioma or a prostate tumor;

said subject having a proliferation-related disorder characterized by a selected IDH, e.g., IDH1 or IDH2 allele, e.g., an IDH1 or IDH2 mutation, having alpha hydroxyl neoactivity, e.g., 2HG neoactivity;

said subject having a selected IDH allele, e.g., a selected IDH1 or IDH2 allele; having alpha hydroxyl neoactivity, e.g., 2HG neoactivity;

said subject having a proliferation-related disorder;

said subject being in need of, or being able to benefit from, a therapeutic agent of a type described herein;

said subject being in need of, or being able to benefit from, a compound that inhibits alpha hydroxyl neoactivity, e.g., 2HG neoactivity;

said subject being in need of, or being able to benefit from, a compound that lowers the level of an alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment evaluation comprises selecting the subject, e.g., for treatment with an anti-neoplastic agent, on the establishment of, or determination that, the subject has increased alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG,

or increased alpha hydroxyl neoactivity, e.g., 2HG neoactivity, or that the subject is in need of inhibition of a neoactivity of an IDH, e.g., IDH1 or IDH2, mutant described herein.

As described herein, the evaluations provided for by methods described herein allow the selection of optimal treatment regimens.

Thus, in an embodiment the method comprises, e.g., responsive to the analysis of one or more of a-d, selecting a treatment for the subject, e.g., selecting a treatment on a basis disclosed herein. The treatment can be the administration of a therapeutic agent disclosed herein. The treatment can be selected on the basis that:

it is useful in treating a disorder characterized by one or more of alpha hydroxyl neoactivity, e.g., 2HG neoactivity, an IDH1 or IDH2, mutant protein having alpha hydroxyl neoactivity, e.g., 2HG neoactivity (or a corresponding RNA);

it is useful in treating a disorder characterized by a selected somatic allele or mutation of an IDH, e.g., IDH1 or IDH2, which encodes a protein with alpha hydroxyl neoactivity, e.g., 2HG neoactivity, e.g., an allele disclosed herein, in cells characterized by a cell proliferation-related disorder from the subject;

it reduces the level of an alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG;

it reduces the level of alpha hydroxyl neoactivity, e.g., 2HG neoactivity.

In an embodiment evaluation comprises selecting the subject, e.g., for treatment.

In embodiments the treatment is the administration of a therapeutic agent described herein.

The methods can also include treating a subject, e.g., with a treatment selected in response to, or on the basis of, an evaluation made in the method.

Thus, in an embodiment the method comprises, e.g., responsive to the analysis of one or more of a-d, administering a treatment to the subject, e.g., the administration of a therapeutic agent of a type described herein.

In an embodiment the therapeutic agent comprises a compound from Table 24a or Table 24b or a compound having the structure of Formula (X) or (XI) described below.

In an embodiment the therapeutic agent comprises nucleic acid, e.g., dsRNA, e.g., a dsRNA described herein.



In an embodiment the the therapeutic agent is an inhibitor, *e.g.*, a polypeptide, peptide, or small molecule (*e.g.*, a molecule of less than 1,000 daltons), or aptomer, that binds to an IDH1 or IDH2 mutant (*e.g.*, an aptomer that binds to an IDH1 mutant) or wildtype subunit and inhibits neoactivity, *e.g.*, by inhibiting formation of a dimer, *e.g.*, a homodimer of mutant IDH1 or IDH2 subunits (*e.g.*, a homodimer of mutant IDH1 subunits) or a heterodimer of a mutant and a wildtype subunit. In an embodiment the inhibitor is a polypeptide. In an embodiment the polypeptide acts as a dominant negative with respect to the neoactivity of the mutant enzyme. The polypeptide can correspond to full length IDH1 or IDH2 or a fragment thereof (*e.g.*, the polypeptide correspondes to full length IDH1 or a fragment thereof). The polypeptide need not be indetical with the corresponding residues of wildtype IDH1 or IDH2 (*e.g.*, wildtype IDH1), but in embodiments has at least 60, 70, 80, 90 or 95 % homology with wildtype IDH1 or IDH2 (*e.g.*, wildtype IDH1).

In an embodiment the therapeutic agent decreases the affinity of an IDH, *e.g.*, IDH1 or IDH2 neoactive mutant protein for NADH, NADPH or a divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , or decreases the levels or availability of NADH, NADPH or divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , *e.g.*, by competing for binding to the mutant enzyme. In an embodiment the enzyme is inhibited by replacing  $Mg^{2+}$  or  $Mn^{2+}$  with  $Ca^{2+}$ .

In an embodiment the therapeutic agent is an inhibitor that reduces the level a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the level of the product of a mutant having a neoactivity of an IDH, *e.g.*, IDH1 or IDH2 mutant, *e.g.*, it reduces the level of 2HG, *e.g.*, R-2HG.

In an embodiment the therapeutic agent is an inhibitor that:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that is selected on the basis that it:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH1, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of a mutant IDH, *e.g.*, IDH1 or IDH2, protein or mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2, protein.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of neoactive enzyme activity, *e.g.*, by interacting with, *e.g.*, binding to, mutant IDH, *e.g.*, IDH1 or IDH2, protein. In an embodiment the inhibitor is other than an antibody.

In an embodiment the therapeutic agent is an inhibitor that is a small molecule and interacts with, *e.g.*, binds, the mutant RNA, *e.g.*, mutant IDH1 mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, binds, either the mutant IDH, *e.g.*, IDH1 or IDH2, protein or interacts directly with, *e.g.*, binds, the mutant IDH mRNA, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the therapeutic agent is administered.

In an embodiment the treatment: inhibits, *e.g.*, specifically, a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, a neoactivity described herein; or inhibits both the wildtype and activity and a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, a neoactivity described herein. In an embodiment, the subject is subsequently evaluated or monitored by a method described herein, *e.g.*, the analysis of the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, to evaluate response to the treatment or progression of disease.

In an embodiment the treatment is selected on the basis that it: inhibits, *e.g.*, specifically, a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity; or inhibits both the wildtype and activity and a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, a neoactivity described herein.

In an embodiment, the method comprises determining the possibility of a mutation other than a mutation in IDH1 or in IDH2. In embodiments a relatively high level of 2HG, *e.g.*, R-2HG is indicative of another mutation.

In an embodiment, which embodiment includes selecting or administering a treatment for the subject, the subject:

has not yet been treated for the subject the cell proliferation-related disorder and the selected or administered treatment is the initial or first line treatment;

has already been treated for the the cell proliferation-related and the selected or administered treatment results in an alteration of the existing treatment;

has already been treated for the the cell proliferation-related, and the selected treatment results in continuation of the existing treatment; or

has already been treated for the the cell proliferation-related disorder and the selected or administered treatment is different, *e.g.*, as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment, which embodiment includes selecting or administering a treatment for the subject, the selected or administered treatment can comprise:

a treatment which includes administration of a therapeutic agent at different, *e.g.*, a greater (or lesser) dosage (*e.g.*, different as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG);

a treatment which includes administration of a therapeutic agent at a different frequency, *e.g.*, more or less frequently, or not at all (*e.g.*, different as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG); or

a treatment which includes administration of a therapeutic agent in a different therapeutic setting (*e.g.*, adding or deleting a second treatment from the treatment regimen) (*e.g.*, different as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG).

Methods of evaluating a subject described herein can comprise evaluating a neoactivity genotype or phenotype. Methods of obtaining and analyzing samples, and the in vivo analysis in subjects, described elsewhere herein, *e.g.*, in the section entitled, "Methods of evaluating samples and/or subjects," can be combined with this method.

In an embodiment the method comprises:

subjecting the subject (*e.g.*, a subject not having 2-hydroxyglutaric aciduria) to imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS *e.g.*, imaging analysis, to provide a determination of the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, as associated with a tumor, *e.g.*, a glioma, in the subject;

optionally storing a parameter related to the determination, *e.g.*, the image or a value related to the image from the imaging analysis, in a tangible medium; and

responsive to the determination, performing one or more of: correlating the determination with outcome or with a prognosis; providing an indication of outcome or prognosis; providing a value for an analysis on which the evaluation is based, *e.g.*, the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG; providing a recommendation for treatment of the subject; selecting a course of treatment for the subject, *e.g.*, a course of treatment described herein, *e.g.*, selecting a course of treatment that includes inhibiting a neoactivity of a mutant IDH, *e.g.*, IDH1 or IDH2, allele, *e.g.*, a neoactivity described herein; administering a course of treatment to the subject, *e.g.*, a course of treatment described herein, *e.g.*, a course of treatment that includes inhibiting a neoactivity of a mutant IDH, *e.g.*, IDH1 or IDH2, allele, *e.g.*, a neoactivity described herein; and memorializing memorializing a result of the method or a measurement made in the course of the method, *e.g.*, one or more of the above and/or transmitting memorialization of one or more of the above to a party, *e.g.*, the subject, a healthcare provider, or an entity that pays for the subject's treatment, *e.g.*, a government, insurance company, or other third party payer.

In an embodiment the method comprises confirming or determining, *e.g.*, by direct examination or evaluation of the subject, or sample *e.g.*, tissue or bodily fluid (*e.g.*, blood (*e.g.*, blood plasma), urine, lymph, or cerebrospinal fluid) therefrom, (*e.g.*, by DNA sequencing or immuno analysis or evaluation of the presence, distribution or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG), or receiving such information about the subject, that the subject has a cancer characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8), in specific embodiments, an IDH1 allele having His, Ser, Cys, Gly, Val, or Leu at residue 132 or an IDH1 allele having His or Cys at residue 132; or an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10).

In an embodiment, prior to or after treatment, the method includes evaluating the growth, size, weight, invasiveness, stage or other phenotype of the cell proliferation-related disorder.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, a leukemia, *e.g.*, AML or ALL, *e.g.*, B-ALL or T-ALL, prostate cancer, or myelodysplasia or myelodysplastic syndrome and the evaluation is a or b. In an embodiment the method comprises evaluating a sample, *e.g.*, a sample described herein, *e.g.*, a tissue, *e.g.*, a cancer sample, or a bodily fluid, *e.g.*, serum or blood, for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment, a subject is subjected to MRS and the evaluation comprises evaluating the presence or elevated amount of a peak correlated to or corresponding to 2HG, *e.g.*, R-2HG, as determined by magnetic resonance. For example, a subject can be analyzed for the presence and/or strength of a signal at about 2.5 ppm to determine the presence and/or amount of 2HG, *e.g.*, R-2HG in the subject.

In an embodiment the method comprises obtaining a sample from the subject and analyzing the sample, or analyzing the subject, *e.g.*, by imaging the subject and optionally forming a representation of the image on a computer.

In an embodiment the results of the analysis is compared to a reference.

In an embodiment a value for a parameter correlated to the presence, distribution, or level, *e.g.*, of 2HG, *e.g.*, R-2HG, is determined. It can be compared with a reference value, *e.g.*, the value for a reference subject not having abnormal presence, level, or distribution, *e.g.*, a reference subject cell not having a mutation in IDH, *e.g.*, IDH1 or IDH2, having a neoactivity described herein.

In an embodiment the method comprises determining if an IDH, *e.g.*, IDH1 or IDH2, mutant allele that is associated with 2HG neoactivity is present. *E.g.*, in the case of IDH1, the presence of a mutation at residue 132 associated with 2HG neoactivity can be determined. In the case of IDH2, the presence of a mutation at residue 172 associated with 2HG neoactivity can be determined. The determination can comprise sequencing a nucleic acid, *e.g.*, genomic DNA or cDNA, from an affected cell, which encodes the relevant amino acid(s). The mutation can be a deletion, insertion, rearrangement, or substitution. The mutation can involve a single nucleotide, *e.g.*, a single substitution, or more than one nucleotide, *e.g.*, a deletion of more than one nucleotides.

In an embodiment the method comprises determining the sequence at position 394 or 395 of the IDH1 gene, or determining the identity of amino acid residue 132 (SEQ ID NO:8) in the IDH1 gene in a cell characterized by the cell proliferation related disorder.

In an embodiment the method comprises determining the amino acid sequence, *e.g.*, by DNA sequencing, at position 172 of the IDH2 gene in a cell characterized by the cell proliferation related disorder.

In an embodiment a product of the neoactivity is 2-HG, *e.g.*, R-2HG, which acts as a metabolite. In another embodiment a product of the neoactivity is 2HG, *e.g.*, R-2HG, which acts as a toxin, *e.g.*, a carcinogen.

In an embodiment the disorder is other than a solid tumor. In an embodiment the disorder is a tumor that, at the time of diagnosis or treatment, does not have a necrotic portion. In an embodiment the disorder is a tumor in which at least 30, 40, 50, 60, 70, 80 or 90% of the tumor cells carry an IHD, *e.g.*, IDH1 or IDH2, mutation having 2HG neoactivity, at the time of diagnosis or treatment.

In an embodiment the cell proliferation-related disorder is a cancer, *e.g.*, a cancer described herein, characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, *e.g.*, wherein the tumor is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. Gliomas include astrocytic tumors, oligodendroglial tumors, oligoastrocytic tumors, anaplastic astrocytomas, and glioblastomas. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8

(see also **Fig. 21**). In an embodiment the allele encodes an IDH1 having His at residue 132. In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the IDH1 allele has an A (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394A, a C394G, a C394T, a G395C, a G395T or a G395A mutation, specifically C394A or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8) (*e.g.*, His, Ser, Cys, Gly, Val, or Leu; or His or Cys).

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8) (*e.g.*, His, Ser, Cys, Gly, Val, or Leu; or His or Cys).

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment, the cell proliferation disorder is fibrosarcoma or paraganglioma wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment, the cell proliferation disorder is fibrosarcoma or paraganglioma wherein the cancer is characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment, the cell proliferation disorder is fibrosarcoma or paraganglioma wherein the cancer is characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is localized or metastatic prostate cancer, *e.g.*, prostate adenocarcinoma, *e.g.*, wherein the cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

*E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al*, 2009, Int. J. Cancer, 125: 353-355 at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (*e.g.*, His, Ser, Cys, Gly, Val, or Leu). In an embodiment the allele encodes an IDH1 having His or Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394T or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, wherein the cancer is characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a hematological cancer, *e.g.*, a leukemia, *e.g.*, AML, or ALL, wherein the hematological cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is acute lymphoblastic leukemia (*e.g.*, an adult or pediatric form), *e.g.*, wherein the acute lymphoblastic leukemia (sometimes referred to herein as ALL) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. The ALL can be, *e.g.*, B-ALL or T-ALL. In an embodiment the



cancer is characterized by increased levels of 2 an alpha hydroxy neoactivity product, e.g., HG, e.g., R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (e.g., His, Ser, Cys, Gly, Val, or Leu). In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 according to the sequence of SEQ ID NO:8.

In an embodiment the method comprises selecting a subject ALL, e.g., B-ALL or T-ALL, on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is acute myelogenous leukemia (*e.g.*, an adult or pediatric form), *e.g.*, wherein the acute myelogenous leukemia (sometimes referred to herein as AML) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (e.g., His, Ser, Cys, Gly, Val or Leu). In an embodiment the allele encodes an IDH1 having Cys, His or Gly at residue 132, specifically, Cys.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplasmic leukemia (AML) characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His or Gly at residue 132 according to the sequence of SEQ ID NO:8, specifically, Cys.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplasmic leukemia (AML) on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His or Gly at residue 132 (SEQ ID NO:8), specifically, Cys.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplasmic leukemia (AML), on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method further comprises evaluating the subject for the presence of a mutation in the NRAS or NPMc gene.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, *e.g.*, wherein the myelodysplasia or myelodysplastic syndrome is characterized by having an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the disorder is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), specifically, His, Ser, Cys, Gly, Val, or Leu. In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 according to the sequence of SEQ ID NO:8.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a glioma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically, Lys, Gly, Met, Trp or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a prostate cancer, *e.g.*, prostate adenocarcinoma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in

described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is ALL, *e.g.*, B-ALL or T-ALL, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels

of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is AML, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, e.g., 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having AML, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, characterized by a mutation, or preselected allele, of IDH2. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In another aspect the invention features a pharmaceutical composition of an inhibitor (*e.g.*, a small molecule or a nucleic acid-based inhibitor) described herein.

In an embodiment a mutant protein specific reagent, *e.g.*, an antibody that specifically binds an IDH mutant protein, *e.g.*, an antibody that specifically binds an IDH1-R132H mutant protein, can be used to detect neoactive mutant enzyme see, for example, that described by Y.Kato et al., "A monoclonal antibody IMab-1 specifically recognizes IDH1<sup>R132H</sup>, the most common glioma-derived mutation: (Kato, Biochem. Biophys. Res. Commun. (2009), which is hereby incorporated by reference in its entirety.

In another aspect, the invention features, a method of evaluating a candidate compound, *e.g.*, for the ability to inhibit a neoactivity of a mutant enzyme, *e.g.*, for use as an anti-proliferative or anti-cancer agent. In an embodiment the mutant enzyme is an IDH, *e.g.*, an IDH1 or IDH2 mutant, *e.g.*, a mutant described herein. In an embodiment the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. The method comprises:

optionally supplying the candidate compound;

contacting the candidate compound with a mutant enzyme having a neoactivity, or with another enzyme, a referred to herein as a proxy enzyme, having an activity, referred to herein as a proxy activity, which is the same as the neoactivity (or with a cell or cell lysate comprising the same); and

evaluating the ability of the candidate compound to modulate, *e.g.*, inhibit or promote, the neoactivity or the proxy activity, thereby evaluating the candidate compound.

In an embodiment the mutant enzyme is a mutant IDH1, *e.g.*, an IDH1 mutant described herein, and the neoactivity is an alpha hydroxy neoactivity, *e.g.*, 2HG

neoactivity. Mutations associated with 2HG neoactivity in IDH1 include mutations at residue 132, e.g., R132H, R132C, R132S, R132G, R132L, or R132V, more specifically, R132H or R132C.

In an embodiment the mutant enzyme is a mutant IDH2, e.g., an IDH2 mutant described herein, and the neoactivity is an alpha hydroxy neoactivity, e.g., 2HG neoactivity. Mutations associated with 2HG neoactivity in IDH2 include mutations at residue 172, e.g., R172K, R172M, R172S, R172G, or R172W.

In an embodiment the method includes evaluating the ability of the candidate compound to inhibit the neoactivity or the proxy activity.

In an embodiment the method further comprises evaluating the ability of the candidate compound to inhibit the forward reaction of non-mutant or wild type enzyme activity, e.g., in the case of IDH, e.g., IDH1 or IDH2, the conversion of isocitrate to  $\alpha$ -ketoglutarate (or an intermediate thereof, including the reduced hydroxyl intermediate).

In an embodiment, the contacting step comprises contacting the candidate compound with a cell, or a cell lysate thereof, wherein the cell comprises a mutant enzyme having the neoactivity or an enzyme having the activity.

In an embodiment, the cell comprises a mutation, or preselected allele, of a mutant IDH1 gene. E.g., in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. E.g., the allele can encode His, Ser, Cys, Gly, Val, Pro or Leu, or any other residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), specifically His, Ser, Cys, Gly, Val, or Leu.

In an embodiment the allele encodes an IDH1 having His at residue 132.

In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the allele is an Arg132His mutation, or an Arg132Ser mutation, according to the sequence of SEQ ID NO:8 (see **FIGs. 2** and **21**).

In an embodiment, the cell comprises a mutation, or preselected allele, of a mutant IDH2 gene. E.g., in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. E.g., the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically, Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment, the cell includes a heterologous copy of a mutant IDH gene, *e.g.*, a mutant IDH1 or IDH2 gene. (Heterologous copy refers to a copy introduced or formed by a genetic engineering manipulation.)

In an embodiment, the cell is transfected (*e.g.*, transiently or stably transfected) or transduced (*e.g.*, transiently or stably transduced) with a nucleic acid sequence encoding an IDH, *e.g.*, IDH1 or IDH2, described herein, *e.g.*, an IDH1 having other than an Arg at residue 132. In an embodiment, the IDH, *e.g.*, IDH1 or IDH2, is epitope-tagged, *e.g.*, myc-tagged.

In an embodiment, the cell, *e.g.*, a cancer cell, is non-mutant or wild type for the IDH, *e.g.*, IDH1 or IDH2, allele. The cell can include a heterologous IDH1 or IDH2 mutant.

In an embodiment, the cell is a cultured cell, *e.g.*, a primary cell, a secondary cell, or a cell line. In an embodiment, the cell is a cancer cell, *e.g.*, a glioma cell (*e.g.*, a glioblastoma cell), a prostate cancer cell, a leukemia cell (*e.g.*, an ALL, *e.g.*, B-ALL or T-ALL, cell or AML cell) or a cell characterized by myelodysplasia or myelodysplastic syndrome. In embodiment, the cell is a 293T cell, a U87MG cell, or an LN-18 cell (*e.g.*, ATCC HTB-14 or CRL-2610).

In an embodiment, the cell is from a subject, *e.g.*, a subject having cancer, *e.g.*, a cancer characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8); specifically His or Cys; or an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), specifically Lys, Gly, Met, Trp, or Ser.

In an embodiment, the evaluating step comprises evaluating the presence and/or amount of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, in the cell lysate or culture medium, *e.g.*, by LC-MS.

In an embodiment, the evaluating step comprises evaluating the presence and/or amount of an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, in the cell lysate or culture medium.

In an embodiment, the method further comprises evaluating the presence/amount one or more of TCA metabolite(s), *e.g.*, citrate,  $\alpha$ -KG, succinate, fumarate, and/or malate, *e.g.*, by LC-MS, *e.g.*, as a control.

In an embodiment, the method further comprises evaluating the oxidation state of NADPH, *e.g.*, the absorbance at 340 nm, *e.g.*, by spectrophotometer.



In an embodiment, the method further comprises evaluating the ability of the candidate compound to inhibit a second enzymatic activity, *e.g.*, the forward reaction of non-mutant or wild type enzyme activity, *e.g.*, in the case of IDH1 or IDH2 (*e.g.*, IDH1), the conversion of isocitrate to  $\alpha$ -ketoglutarate (or an intermediate thereof, including the reduced hydroxyl intermediate).

In an embodiment, the candidate compound is a small molecule, a polypeptide, peptide, a carbohydrate based molecule, or an aptamer (*e.g.*, a nucleic acid aptamer, or a peptide aptamer). The method can be used broadly and can, *e.g.*, be used as one or more of a primary screen, to confirm candidates produced by this or other methods or screens, or generally to guide drug discovery or drug candidate optimization.

In an embodiment, the method comprises evaluating, *e.g.*, confirming, the ability of a candidate compound (*e.g.*, a candidate compound which meets a predetermined level of inhibition in the evaluating step) to inhibit the neoactivity or proxy activity in a second assay.

In an embodiment, the second assay comprises repeating one or more of the contacting and/or evaluating step(s) of the basic method.

In another embodiment, the second assay is different from the first. *E.g.*, where the first assay can use a cell or cell lysate or other non-whole animal model the second assay can use an animal model, *e.g.*, a tumor transplant model, *e.g.*, a mouse having an IDH, *e.g.*, IDH1 or IDH2, mutant cell or tumor transplanted in it. *E.g.*, a U87 cell, or glioma, *e.g.*, glioblastoma, cell, harboring a transfected IDH, *e.g.*, IDH1 or IDH2, neoactive mutant can be implanted as a xenograft and used in an assay. Primary human glioma or AML tumor cells can be grafted into mice to allow propagation of the tumor and used in an assay. A genetically engineered mouse model (GEMM) harboring an IDH1 or IDH2 mutation and/or other mutation, *e.g.*, a p53 null mutation, can also be used in an assay.

In an embodiment the method comprises:

optionally supplying the candidate compound;

contacting the candidate compound with a cell comprising a nucleic acid sequence, *e.g.*, a heterologous sequence, encoding an IDH1 having other than an Arg at residue 132 (*e.g.*, IDH1R132H) or an IDH2 having other than an Arg at residue 172 (specifically an IDH1 having other than an Arg at residue 132); and

evaluating the presence and/or amount of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, in the cell lysate or culture medium, by LC-MS,

thereby evaluating the compound.

In an embodiment the result of the evaluation is compared with a reference, *e.g.*, the level of product, *e.g.*, an alpha hydroxy neoactivity product, *e.g.*, 2HG. *e.g.*, R-2HG, in a control cell, *e.g.*, a cell having inserted therein a wild type or non-mutant copy of IDH1 or IDH2 (*e.g.*, IDH1).

In another aspect, the invention features, a method of evaluating a candidate compound, *e.g.*, for the ability to inhibit an RNA encoding a mutant enzyme having a neoactivity, *e.g.*, for use as an anti-proliferative or anti-cancer agent. In an embodiment the mutant enzyme is an IDH, *e.g.*, an IDH1 or IDH2 mutant, *e.g.*, a mutant described herein. In an embodiment the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. The method comprises:

optionally supplying the candidate compound, *e.g.*, a nucleic acid based inhibitor (*e.g.*, a dsRNA (*e.g.*, siRNA or shRNA), an antisense, or a microRNA);

contacting the candidate compound with an RNA, *e.g.*, an mRNA, which encodes IDH, *e.g.*, an IDH1 or IDH2, *e.g.*, an RNA that encode mutant enzyme having a neoactivity (or with a cell or cell lysate comprising the same); and

evaluating the ability of the candidate compound to inhibit the RNA, thereby evaluating the candidate compound. By inhibit the RNA means, *e.g.*, to cleave or otherwise inactivate the RNA.

In an embodiment the RNA encodes a fusion of all or part of the IDH, *e.g.*, IDH1 or IDH2, wildtype or mutant protein to a second protein, *e.g.*, a reporter protein, *e.g.*, a fluorescent protein, *e.g.*, a green or red fluorescent protein.

In an embodiment the mutant enzyme is a mutant IDH1, *e.g.*, an IDH1 mutant described herein, and the neoactivity is an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity.

In an embodiment the mutant enzyme is a mutant IDH2, *e.g.*, an IDH2 mutant described herein, and the neoactivity is an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity.

In an embodiment, the contacting step comprises contacting the candidate compound with a cell, or a cell lysate thereof, wherein the cell comprises RNA encoding IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a mutant IDH, *e.g.*, IDH1 or IDH2, enzyme having the neoactivity.

In an embodiment, the cell comprises a mutation, or preselected allele, of a mutant IDH1 gene. *E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having

other than an Arg at residue 132. *E.g.*, the allele can encode His, Ser, Cys, Gly, Val, Pro or Leu, or any other residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), specifically His, Ser, Cys, Gly, Val, or Leu.

In an embodiment the allele encodes an IDH1 having His at residue 132.

In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the allele is an Arg132His mutation, or an Arg132Ser mutation, according to the sequence of SEQ ID NO:8 (see **FIGs. 2 and 21**).

In an embodiment, the cell comprises a mutation, or preselected allele, of a mutant IDH2 gene. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment, the cell includes a heterologous copy of a wildtype or mutant IDH gene, *e.g.*, a wildtype or mutant IDH1 or IDH2 gene. (Heterologous copy refers to a copy introduced or formed by a genetic engineering manipulation.) In an embodiment the heterologous gene comprises a fusion to a reporter protein, *e.g.*, a fluorescent protein, *e.g.*, a green or red fluorescent protein.

In an embodiment, the cell is transfected (*e.g.*, transiently or stably transfected) or transduced (*e.g.*, transiently or stably transduced) with a nucleic acid sequence encoding an IDH, *e.g.*, IDH1 or IDH2, described herein, *e.g.*, an IDH1 having other than an Arg at residue 132 or an IDH2 having other than an Arg at residue 172 (*e.g.*, an IDH1 having other than an Arg at residue 132). In an embodiment, the IDH, *e.g.*, IDH1 or IDH2, is epitope-tagged, *e.g.*, myc-tagged.

In an embodiment, the cell, *e.g.*, a cancer cell, is non-mutant or wild type for the IDH, *e.g.*, IDH1 or IDH2, allele. The cell can include a heterologous IDH1 or IDH2 mutant.

In an embodiment, the cell is a cultured cell, *e.g.*, a primary cell, a secondary cell, or a cell line. In an embodiment, the cell is a cancer cell, *e.g.*, a glioma cell (*e.g.*, a glioblastoma cell), a prostate cancer cell, a leukemia cell (*e.g.*, an ALL, *e.g.*, B-ALL or T-ALL cell or AML cell) or a cell characterized by myelodysplasia or

myelodysplastic syndrome. In embodiment, the cell is a 293T cell, a U87MG cell, or an LN-18 cell (*e.g.*, ATCC HTB-14 or CRL-2610).

In an embodiment, the cell is from a subject, *e.g.*, a subject having cancer, *e.g.*, a cancer characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8); specifically His or Cys. In an embodiment, the cancer is characterized by an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), specifically Lys, Gly, Met, Trp, or Ser.

In an embodiment, the method comprises a second assay and the second assay comprises repeating one or more of the contacting and/or evaluating step(s) of the basic method.

In another embodiment, the second assay is different from the first. *E.g.*, where the first assay can use a cell or cell lysate or other non-whole animal model the second assay can use an animal model

In an embodiment the efficacy of the candidate is evaluated by its effect on reporter protein activity.

In another aspect, the invention features, a method of evaluating a candidate compound, *e.g.*, for the ability to inhibit transcription of an RNA encoding a mutant enzyme having a neoactivity, *e.g.*, for use as an anti-proliferative or anti-cancer agent. In an embodiment the mutant enzyme is an IDH, *e.g.*, an IDH1 or IDH2 mutant, *e.g.*, a mutant described herein. In an embodiment the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. The method comprises:

optionally supplying the candidate compound, *e.g.*, a small molecule, polypeptide, peptide, aptomer, a carbohydrate-based molecule or nucleic acid based molecule;

contacting the candidate compound with a system comprising a cell or cell lysate; and

evaluating the ability of the candidate compound to inhibit the translation of IDH, *e.g.*, IDH1 or IDH2, RNA, *e.g.*, thereby evaluating the candidate compound.

In an embodiment the the system comprises a fusion gene encoding of all or part of the IDH, *e.g.*, IDH1 or IDH2, wildtype or mutant protein to a second protein, *e.g.*, a reporter protein, *e.g.*, a fluorescent protein, *e.g.*, a green or red fluorescent protein.

In an embodiment the mutant enzyme is a mutant IDH1, *e.g.*, an IDH1 mutant described herein, and the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity.

In an embodiment the mutant enzyme is a mutant IDH2, *e.g.*, an IDH2 mutant described herein, and the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity.

In an embodiment, the system includes a heterologous copy of a wildtype or mutant IDH gene, *e.g.*, a wildtype or mutant IDH1 or IDH2 gene. (Heterologous copy refers to a copy introduced or formed by a genetic engineering manipulation.) In an embodiment the heterologous gene comprises a fusion to a reporter protein, *e.g.*, a fluorescent protein, *e.g.*, a green or red fluorescent protein.

In an embodiment the cell, *e.g.*, a cancer cell, is non-mutant or wild type for the IDH, *e.g.*, IDH1 or IDH2, allele. The cell can include a heterologous IDH1 or IDH2 mutant.

In an embodiment, the cell is a cultured cell, *e.g.*, a primary cell, a secondary cell, or a cell line. In an embodiment, the cell is a cancer cell, *e.g.*, a glioma cell (*e.g.*, a glioblastoma cell), a prostate cancer cell, a leukemia cell (*e.g.*, an ALL, *e.g.*, B-ALL or T-ALL, cell or AML cell) or a cell characterized by myelodysplasia or myelodysplastic syndrome. In embodiment, the cell is a 293T cell, a U87MG cell, or an LN-18 cell (*e.g.*, ATCC HTB-14 or CRL-2610).

In an embodiment, the cell is from a subject, *e.g.*, a subject having cancer, *e.g.*, a cancer characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8); specifically His, Ser, Cys, Gly, Val, or Leu. In an embodiment, the cancer is characterized an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10).

In an embodiment, the method comprises a second assay and the second assay comprises repeating the method.

In another embodiment, the second assay is different from the first. *E.g.*, where the first assay can use a cell or cell lysate or other non-whole animal model the second assay can use an animal model.

In an embodiment the efficacy of the candidate is evaluated by its effect on reporter protein activity.

In another aspect, the invention features, a method of evaluating a candidate compound, e.g., a therapeutic agent, or inhibitor, described herein in an animal model. The candidate compound can be, e.g., a small molecule, polypeptide, peptide, aptomer, a carbohydrate-based molecule or nucleic acid based molecule. The method comprises, contacting the candidate with the animal model and evaluating the animal model.

In an embodiment evaluating comprises;

- determining an effect of the compound on the general health of the animal;
- determining an effect of the compound on the weight of the animal;
- determining an effect of the compound on liver function, e.g, on a liver enzyme;
- determining an effect of the compound on the cardiovascular system of the animal;
- determining an effect of the compound on neurofunction, e.g., on neuromuscular control or response;
- determining an effect of the compound on eating or drinking;
- determining the distribution of the compound in the animal;
- determining the persistence of the compound in the animal or in a tissue or organ of the animal, e.g., determining plasma half-life; or
- determining an effect of the compound on a selected cell in the animal;
- determining an effect of the compound on the growth, size, weight, invasiveness or other phenotype of a tumor, e.g., an endogenous tumor or a tumor arising from introduction of cells from the same or a different species.

In an embodiment the animal is a non-human primate, e.g., a cynomolgus monkey or chimpanzee.

In an embodiment the animal is a rodent, e.g., a rat or mouse.

In an embodiment the animal is a large animal, e.g., a dog or pig, other than a non-human primate.

In an embodiment the evaluation is memorialized and optionally transmitted to another party.

In one aspect, the invention provides, a method of evaluating or processing a therapeutic agent, e.g., a therapeutic agent referred to herein, e.g., a therapeutic agent that results in a lowering of the level of a product of an IDH, e.g., IDH1 or IDH2, mutant having a neoactivity. In an embodiment the neoactivity is an alpha hydroxy

neoactivity, e.g., 2HG neoactivity, and the level of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, is lowered.

The method includes:

providing, e.g., by testing a sample, a value (e.g., a test value) for a parameter related to a property of the therapeutic agent, e.g., the ability to inhibit the conversion of alpha ketoglutarate to 2 hydroxyglutarate (i.e., 2HG), e.g., R-2 hydroxyglutarate (i.e., R-2HG), and,

optionally, providing a determination of whether the value determined for the parameter meets a preselected criterion, e.g., is present, or is present within a preselected range,

thereby evaluating or processing the therapeutic agent.

In an embodiment the therapeutic agent is approved for use in humans by a government agency, e.g., the FDA.

In an embodiment the parameter is correlated to the ability to inhibit 2HG neoactivity, and, e.g., the therapeutic agent is an inhibitor which binds to IDH1 or IDH2 protein and reduces an alpha hydroxy neoactivity, e.g., 2HG neoactivity.

In an embodiment the parameter is correlated to the level of mutant IDH, e.g., IDH1 or IDH2, protein, and, e.g., the therapeutic agent is an inhibitor which reduces the level of IDH1 or IDH2 mutant protein.

In an embodiment the parameter is correlated to the level of an RNA that encodes a mutant IDH, e.g., IDH1 or IDH2, protein, and, e.g., the therapeutic agent reduces the level of RNA, e.g., mRNA, that encodes IDH1 or IDH2 mutant protein.

In an embodiment the method includes contacting the therapeutic agent with a mutant IDH, e.g., IDH1 or IDH2, protein (or corresponding RNA).

In an embodiment, the method includes providing a comparison of the value determined for a parameter with a reference value or values, to thereby evaluate the therapeutic agent. In an embodiment, the comparison includes determining if a test value determined for the therapeutic agent has a preselected relationship with the reference value, e.g., determining if it meets the reference value. The value need not be a numerical value but, e.g., can be merely an indication of whether an activity is present.

In an embodiment the method includes determining if a test value is equal to or greater than a reference value, if it is less than or equal to a reference value, or if it falls within a range (either inclusive or exclusive of one or both endpoints). In an

embodiment, the test value, or an indication of whether the preselected criterion is met, can be memorialized, *e.g.*, in a computer readable record.

In an embodiment, a decision or step is taken, *e.g.*, a sample containing the therapeutic agent, or a batch of the therapeutic agent, is classified, selected, accepted or discarded, released or withheld, processed into a drug product, shipped, moved to a different location, formulated, labeled, packaged, contacted with, or put into, a container, *e.g.*, a gas or liquid tight container, released into commerce, or sold or offered for sale, or a record made or altered to reflect the determination, depending on whether the preselected criterion is met. *E.g.*, based on the result of the determination or whether an activity is present, or upon comparison to a reference standard, the batch from which the sample is taken can be processed, *e.g.*, as just described.

The evaluation of the presence or level of activity can show if the therapeutic agent meets a reference standard.

In an embodiment, methods and compositions disclosed herein are useful from a process standpoint, *e.g.*, to monitor or ensure batch-to-batch consistency or quality, or to evaluate a sample with regard to a reference, *e.g.*, a preselected value.

In an embodiment, the method can be used to determine if a test batch of a therapeutic agent can be expected to have one or more of the properties. Such properties can include a property listed on the product insert of a therapeutic agent, a property appearing in a compendium, *e.g.*, the US Pharmacopea, or a property required by a regulatory agency, *e.g.*, the FDA, for commercial use.

In an embodiment the method includes testing the therapeutic agent for its effect on the wildtype activity of an IDH, *e.g.*, IDH1 or IDH2, protein, and providing a determination of whether the value determined meets a preselected criterion, *e.g.*, is present, or is present within a preselected range.

In an embodiment the method includes:

contacting a therapeutic agent that is an inhibitor of IDH1 an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, with an IDH1 mutant having an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity,

determining a value related to the inhibition of an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, and

comparing the value determined with a reference value, *e.g.*, a range of values, for the inhibition of an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. In an embodiment the reference value is an FDA required value, *e.g.*, a release criteria.



In an embodiment the method includes:

contacting a therapeutic agent that is an inhibitor of mRNA which encodes a mutant IDH1 having an alpha hydroxy neoactivity, e.g., 2HG neoactivity, with an mRNA that encodes an IDH1 mutant having an alpha hydroxy neoactivity, e.g., 2HG neoactivity,

determining a value related to the inhibition of the mRNA, and,

comparing the value determined with a reference value, e.g., a range of values for inhibition of the mRNA. In an embodiment the reference value is an FDA required value, e.g., a release criteria.

In one aspect, the invention features a method of evaluating a sample of a therapeutic agent, e.g., a therapeutic agent referred to herein, that includes receiving data with regard to an activity of the therapeutic agent; providing a record which includes said data and optionally includes an identifier for a batch of therapeutic agent; submitting said record to a decision-maker, e.g., a government agency, e.g., the FDA; optionally, receiving a communication from said decision maker; optionally, deciding whether to release market the batch of therapeutic agent based on the communication from the decision maker. In one embodiment, the method further includes releasing, or other wise processing, e.g., as described herein, the sample.

In another aspect, the invention features, a method of selecting a payment class for treatment with a therapeutic agent described herein, e.g., an inhibitor of IDH, e.g., IDH1 or IDH2, neoactivity, for a subject having a cell proliferation-related disorder. The method includes:

providing (e.g., receiving) an evaluation of whether the subject is positive for increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, or neoactivity, e.g., an alpha hydroxy neoactivity, e.g., 2HG neoactivity, a mutant IDH1 or IDH2 having neoactivity, e.g., an alpha hydroxy neoactivity, e.g., 2HG neoactivity, (or a corresponding RNA), or a mutant IDH, e.g., IDH1 or IDH2, somatic gene, e.g., a mutant described herein, and

performing at least one of (1) if the subject is positive selecting a first payment class, and (2) if the subject is a not positive selecting a second payment class.

In an embodiment the selection is memorialized, e.g., in a medical records system.

In an embodiment the method includes evaluation of whether the subject is positive for increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, or neoactivity, *e.g.*, an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity.

In an embodiment the method includes requesting the evaluation.

In an embodiment the evaluation is performed on the subject by a method described herein.

In an embodiment, the method comprises communicating the selection to another party, *e.g.*, by computer, compact disc, telephone, facsimile, email, or letter.

In an embodiment, the method comprises making or authorizing payment for said treatment.

In an embodiment, payment is by a first party to a second party. In some embodiments, the first party is other than the subject. In some embodiments, the first party is selected from a third party payor, an insurance company, employer, employer sponsored health plan, HMO, or governmental entity. In some embodiments, the second party is selected from the subject, a healthcare provider, a treating physician, an HMO, a hospital, a governmental entity, or an entity which sells or supplies the drug. In some embodiments, the first party is an insurance company and the second party is selected from the subject, a healthcare provider, a treating physician, an HMO, a hospital, a governmental entity, or an entity which sells or supplies the drug. In some embodiments, the first party is a governmental entity and the second party is selected from the subject, a healthcare provider, a treating physician, an HMO, a hospital, an insurance company, or an entity which sells or supplies the drug.

As used herein, a cell proliferation-related disorder is a disorder characterized by unwanted cell proliferation or by a predisposition to lead to unwanted cell proliferation (sometimes referred to as a precancerous disorder). Examples of disorders characterized by unwanted cell proliferation include cancers, *e.g.*, tumors of the CNS, *e.g.*, a glioma. Gliomas include astrocytic tumors, oligodendroglial tumors, oligoastrocytic tumors, anaplastic astrocytomas, and glioblastomas. Other examples include hematological cancers, *e.g.*, a leukemia, *e.g.*, AML (*e.g.*, an adult or pediatric form) or ALL, *e.g.*, B-ALL or T-ALL (*e.g.*, an adult or pediatric form), localized or metastatic prostate cancer, *e.g.*, prostate adenocarcinoma, fibrosarcoma, and paraganglioma; specifically leukemia, *e.g.*, AML (*e.g.*, an adult or pediatric form) or ALL, *e.g.*, B-ALL or T-ALL (*e.g.*, an adult or pediatric form), localized or metastatic prostate cancer, *e.g.*, prostate adenocarcinoma. Examples of disorders characterized

by a predisposition to lead to unwanted cell proliferation include myelodysplasia or myelodysplastic syndrome, which are a diverse collection of hematological conditions marked by ineffective production (or dysplasia) of myeloid blood cells and risk of transformation to AML.

As used herein, specifically inhibits a neoactivity (and similar language), means the neoactivity of the mutant enzyme is inhibited to a significantly greater degree than is the wildtype enzyme activity. By way of example, “specifically inhibits the 2HG neoactivity of mutant IDH1 (or IDH2)” means the 2HG neoactivity is inhibited to a significantly greater degree than is the forward reaction (the conversion of isocitrate to alpha ketoglutarate) of wildtype IDH1 (or IDH2) activity. In embodiments the neoactivity is inhibited at least 2, 5, 10, or 100 fold more than the wildtype activity. In embodiments an inhibitor that is specific for the 2HG neoactivity of IDH, e.g., IDH1 or IDH2, will also inhibit another dehydrogenase, e.g., malate dehydrogenase. In other embodiments the specific inhibitor does inhibit other dehydrogenases, e.g., malate dehydrogenase.

As used herein, a cell proliferation-related disorder, e.g., a cancer, characterized by a mutation or allele, means a cell proliferation-related disorder having a substantial number of cells which carry that mutation or allele. In an embodiment at least 10, 25, 50, 75, 90, 95 or 99% of the cell proliferation-related disorder cells, e.g., the cells of a cancer, or a representative, average or typical sample of cancer cells, e.g., from a tumor or from affected blood cells, carry at least one copy of the mutation or allele. A cell proliferation-related disorder, characterized by a mutant IDH, e.g., a mutant IDH1 or mutant IDH2, having 2HG neoactivity is exemplary. In an embodiment the mutation or allele is present as a heterozygote at the indicated frequencies.

As used herein, a “SNP” is a DNA sequence variation occurring when a single nucleotide (A, T, C, or G) in the genome (or other shared sequence) differs between members of a species (or between paired chromosomes in an individual).

As used herein, a subject can be a human or non-human subject. Non-human subjects include non-human primates, rodents, e.g., mice or rats, or other non-human animals.

The details of one or more embodiments of the invention are set forth in the description below. Other features, objects, and advantages of the invention will be apparent from the description and the drawings, and from the claims.

### BRIEF DESCRIPTION OF THE DRAWINGS

**FIG. 1** depicts DNA sequence verification of pET41a-IDH1 and alignment against published IDH1 CDS. The sequence of IDH1 (CDS) corresponds to SEQ ID NO:5. The sequence of pET41a-IDH1 corresponds to SEQ ID NO:6, and the “consensus” sequence corresponds to SEQ ID NO:7.

**FIG. 2** depicts DNA sequence verification of R132S and R132H mutants according to the SEQ ID NO:8. The amino acid sequence of IDH1 (SEQ ID NO:8) is provided in FIG. 21.

**FIG. 3** depicts separation of wild type IDH1 protein on Ni-Sepharose column.

**FIG. 4** depicts protein analysis of wild type IDH1 on SDS gel pre and post Ni column fractionation. T: total protein; I: insoluble fractions; S: soluble fraction; L: sample for loading on Ni-column. The numbers in the figure indicates the fraction numbers. Fractions #17 ~ #27 were collected for further purification.

**FIG. 5A** depicts separation of wild type IDH1 protein through SEC column S-200.

**FIG. 5B** depicts protein analysis of wild type IDH1 on SDS gel pre and post S-200 column fractionation. M: molecular weight marker; Ni: nickel column fraction prior to S-200; S200: fraction from SEC column.

**FIG. 6** depicts separation of mutant R132S protein on Ni-Sepharose column.

**FIG. 7** depicts protein analysis of mutant R132S on SDS gel pre and post Ni column fractionation. M: protein marker (KDa): 116, 66.2, 45, 35, 25, 18.4, 14.4; T: total cell protein; So: soluble fraction; In: insoluble fraction; Ft: flow through. #3-#7 indicate the corresponding eluted fraction numbers.

**FIG. 8A** depicts separation of mutant R132S protein through SEC column S-200.

**FIG. 8B** depicts protein analysis of mutant R132S on SDS gel post S-200 column fractionation. M: molecular weight marker; R132S: fraction from SEC column.

**FIG. 9** depicts separation of mutant R132H protein on Ni-Sepharose column.

**FIG. 10** depicts protein analysis of mutant R132H on SDS gel pre and post Ni column fractionation. M: protein marker (KDa): 116, 66.2, 45, 35, 25, 18.4, 14.4; T: total cell protein; So: soluble fraction; In: insoluble fraction; Ft: flow through; #5-#10 indicate the corresponding eluted fraction numbers; Ni: sample from Ni-Sepharose column, pool #5-#10 together.

**FIG. 11A** depicts separation of mutant R132H protein through SEC column S-200.

**FIG. 11B** depicts protein analysis of mutant R132H on SDS gel post S-200 column fractionation. M: molecular weight marker; R132H: fraction from SEC column.

**FIG. 12A** depicts Michaelis-Menten plot of IDH1 wild-type in the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate.

**FIG. 12B** depicts Michaelis-Menten plot of R132H mutant enzyme in the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate.

**FIG. 12C** depicts Michaelis-Menten plot of R132S mutant enzyme in the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate.

**FIG. 13A** depicts  $\alpha$ -KG inhibition of IDH1 wild-type.

**FIG. 13B** depicts  $\alpha$ -KG inhibition of R132H mutant enzyme.

**FIG. 13C** depicts  $\alpha$ -KG inhibition of R132S mutant enzyme.

**FIG. 14** depicts IDH1 wt, R132H, and R132S in the conversion  $\alpha$ -ketoglutarate to 2-hydroxyglutarate.

**FIG. 15A** depicts Substrate-Concentration velocity plot for R132H mutant enzyme.

**FIG. 15B** depicts Substrate-Concentration velocity plot for R132S mutant enzyme.

**FIG. 16** depicts IDH1 wt, R132H, and R132S in the conversion  $\alpha$ -ketoglutarate to 2-hydroxyglutarate with NADH.

**FIG. 17A** depicts oxalomalate inhibition to IDH1 wt.

**FIG. 17B** depicts oxalomalate inhibition to R132H.

**FIG. 17C** depicts oxalomalate inhibition to R132S.

**FIG. 18A** depicts LC-MS/MS analysis of the control reaction.

**FIG. 18B** depicts LC-MS/MS analysis of the reaction containing enzyme.

**FIG. 18C** depicts LC-MS/MS analysis of the spiked control reaction.

**FIG. 19** depicts LC-MS/MS analysis of alpha-hydroxyglutarate.

**FIG. 20** depicts LC-MS/MS analysis showing that R132H consumes  $\alpha$ -KG to produce 2-hydroxyglutaric acid.

**FIG. 21** depicts the amino acid sequence of IDH1 (SEQ ID NO:13) as described in GenBank Accession No. NP\_005887.2 (GI No. 28178825) (record dated May 10, 2009).

**FIG. 21A** is the cDNA sequence of IDH1 as presented at GenBank Accession No. NM\_005896.2 (Record dated May 10, 2009; GI No. 28178824) (SEQ ID NO:8).

**FIG. 21B** depicts the mRNA sequence of IDH1 as described in GenBank Accession No. NM\_005896.2 (Record dated May 10, 2009; GI No. 28178824) (SEQ ID NO:9).

**FIG. 22** is the amino acid sequence of IDH2 as presented at GenBank Accession No. NM\_002168.2 (Record dated August 16, 2009; GI28178831) (SEQ ID NO:10).

**FIG. 22A** is the cDNA sequence of IDH2 as presented at GenBank Accession No. NM\_002168 (Record dated August 16, 2009; GI28178831) (SEQ ID NO:11).

**FIG. 22B** is the mRNA sequence of IDH2 as presented at GenBank Accession No. NM\_002168.2 (Record dated August 16, 2009; GI28178831) (SEQ ID NO:12).

**FIG. 23** depicts the progress of forward reactions (isocitrate to  $\alpha$ -KG) for the mutant enzyme R132H and R132S.

**FIG. 24A** depicts LC-MS/MS analysis of derivitized 2-HG racemic mixture.

**FIG. 24B** depicts LC-MS/MS analysis of derivitized R-2HG standard.

**FIG. 24C** depicts LC-MS/MS analysis of a coinjection of derivitized 2-HG racemate and R-2-HG standard.

**FIG. 24D** depicts LC-MS/MS analysis of the derivitized neoactivity reaction product.

**FIG. 24E** depicts LC-MS/MS analysis of a coinjection of the neoactivity enzyme reaction product and the R-2-HG standard.

**FIG. 24F** depicts LC-MS/MS analysis of a coinjection of the neoactivity enzyme reaction product and the 2-HG racemic mixture.

**FIG. 25** depicts the inhibitory effect of 2-HG derived from the reduction of  $\alpha$ -KG by ICDH1 R132H on the wild-type ICDH1 catalytic oxidative decarboxylation of isocitrate to  $\alpha$ -KG.

**FIG. 26A** depicts levels of 2-HG in CRL-2610 cell lines expressing wildtype or IDH-1 R132H mutant protein.

**FIG. 26B** depicts levels of 2-HG in HTB-14 cell lines expressing wildtype or IDH-1 R132H mutant protein.

**FIG. 27** depicts human IDH1 genomic DNA: intron/2<sup>nd</sup> exon sequence.

**FIG. 28** depicts concentrations of 2HG in human malignant gliomas containing R132 mutations in IDH1. Human glioma samples obtained by surgical resection were snap frozen, genotyped to stratify as wild-type (WT) (N=10) or carrying an R132 mutant allele (Mutant) (n=12) and metabolites extracted for LC-MS analysis. Among the 12 mutant tumors, 10 carried a R132H mutation, one an R132S mutation, and one an R132G mutation. Each symbol represents the amount of the listed metabolite found in each tumor sample. Red lines indicate the group sample means. The difference in 2HG observed between WT and R132 mutant IDH1 mutant tumors was statistically significant by Student's t-test ( $p < 0.0001$ ). There were no statistically significant

differences in  $\alpha$ KG, malate, fumarate, succinate, or isocitrate levels between the WT and R132 mutant IDH1 tumors.

**FIG. 29A** depicts the structural analysis of R132H mutant IDH1. On left is shown an overlay structure of R132H mutant IDH1 and WT IDH1 in the ‘closed’ conformation. On the right is shown an overlay structure of WT IDH1 in the ‘open’ conformation with mutant IDH1 for comparison.

**FIG. 29B** depicts the close-up structural comparison of the R132H IDH1 (left) and wild-type (WT) IDH1 (right) active-site containing both  $\alpha$ KG and NADPH. In addition to changes at residue 132, the position of the catalytic residues Tyr 139 and Lys 212 are different and  $\alpha$ KG is oriented differently relative to NADPH for catalytic hydride transfer in the WT versus R132H mutant enzymes.

**FIG. 30A** depicts the enzymatic properties of IDH1 R132H mutants when recombinant human wild-type (WT) and R132H mutant (R132H) IDH1 enzymes were assessed for oxidative decarboxylation of isocitrate to  $\alpha$ KG with NADP<sup>+</sup> as cofactor. Different concentrations of enzyme were used to generate the curves.

**FIG. 30B** depicts the enzymatic properties of IDH R132 mutants when WT and R132H mutant IDH1 enzymes were assessed for reduction of  $\alpha$ KG with NADPH as cofactor. Different concentrations of enzyme were used to generate the curves.

**FIG. 30C** depicts kinetic parameters of oxidative and reductive reactions as measured for WT and R132H IDH1 enzymes are shown.  $K_m$  and  $k_{cat}$  values for the reductive activity of the WT enzyme were unable to be determined as no measurable enzyme activity was detectable at any substrate concentration.

**FIG. 31A** depicts the LC-MS/MS analysis identifying 2HG as the reductive reaction product of recombinant human R132H mutant IDH1.

**FIG. 31B** depicts the diacetyl-L-tartaric anhydride derivatization and LC-MS/MS analysis of the chirality of 2HG produced by R132H mutant IDH1. Normalized LC-MS/MS signal for the reductive reaction (rxn) product alone, an R(-)-2HG standard alone, and the two together (Rxn + R(-)-2HG) are shown as is the signal for a racemic mixture of R(-) and S(+) forms (2HG Racemate) alone or with the reaction products (Rxn + Racemate).

**FIG. 32A** depicts SDS-PAGE and Western blot analyses of C-terminal affinity-purification tagged IDH1 R132S protein used for crystallization.

**FIG. 32B** depicts the chromatogram of FPLC analysis of the IDH1 R132S protein sample.

**FIG. 33** depicts crystals obtained from a protein solution contained 5 mM NADP, 5 mM isocitrate, 10 mM Ca<sup>2+</sup>. Precipitant solution contained 100 mM MES (pH 6.0) and 20% PEG 6000 using a hanging drop method of crystallization.

**FIG. 34** depicts crystal obtained from a protein solution contained 5 mM NADP, 5 mM  $\alpha$ -ketoglutarate, 10 mM Ca<sup>2+</sup>. Precipitant contained 100 mM MES (pH 6.5) and 12% PEG 20000.

**FIG. 35** is a bar graph depicting elevated NADPH reductive catalysis activity in IDH2-R172K mutant enzyme as compared to wildtype IDH2.

**FIGs. 36A-C** are graphs depicting the following: **(A)** Extracts from IDH1/2 wt (n=10), and IDH1/2 mutant (n=16) patient leukemia cells obtained at presentation and relapse, and IDH1 R132 mutant leukemia cells grown in culture for 14 days (n=14) analyzed by LC-MS to measure levels of 2-HG; and **(B)** 2-HG measured in serum of patients with IDH1 wt or IDH1 R132 mutant leukemia. In **(A)** and **(B)**, each point represents an individual patient sample. Diamonds represent wildtype, circles represent IDH1 mutants, and triangles represent IDH2 mutants. Horizontal bars indicate the mean. (\*) indicates a statistically significant difference relative to wild-type patient cells (p<0.05). **(C)** depicts *In vitro* growth curves of IDH1 R132 mutant and IDH1 wild-type AML cells.

**FIG. 37** is a graph depicting the results of extracts from leukemia cells of AML patients carrying an IDH1/2 mutant (n=16) or wild-type (n=10) allele obtained at initial presentation and relapse assayed by LC-MS for levels of  $\alpha$ -KG, succinate, malate, and fumarate. Each point represents an individual patient sample. Open circles represent wild-types, closed circles represent IDH1 mutants, and triangles represent IDH2 mutants. Horizontal bars represent the mean. There were no statistically significant differences between the wild-type and IDH1/2 mutant AML samples.

**FIG. 38** depicts graphical representations of LC-MS analysis of *in vitro* reactions using recombinant IDH1 R132C and IDH2 R172K confirming that 2-HG and not isocitrate is the end product of the mutant enzyme reactions.

**FIGs. 39A and B** depict **(A)** the wild-type IDH1 enzyme catalysis of the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate with the concomitant reduction of NADP to NADPH; and **(B)** the IDH1 R132C mutant reduction of  $\alpha$ -ketoglutarate



to 2-hydroxyglutarate while oxidizing NADPH to NADP. These are referred to as the “forward” and “partial reverse” reactions, respectively.

#### **DETAILED DESCRIPTION**

The inventors have discovered that certain mutated forms of an enzyme (*e.g.*, IDH1 or IDH2) have a gain of function, referred to herein as a neoactivity, which can be targeted in the treatment of a cell proliferation-related disorder, *e.g.*, a proliferative disorder such as cancer. For example, in the case of a metabolic pathway enzyme, a gain of function or neoactivity can serve as a target for treatment of cancer.

Described herein are methods and compositions for the treatment of a cell proliferation-related disorder, *e.g.*, a proliferative disorder such as cancer. The methods include, *e.g.*, treating a subject having a glioma or brain tumor characterized by a preselected IDH1 allele, *e.g.*, an allele having A at position 394, such as a C394A, a C394G, a C394T, a G395C, a G395T or a G395A mutation, (*e.g.*, a C394A mutant) or an A at position 395 (*e.g.*, a G395A mutant) according to the sequence of SEQ ID NO:5, that encodes an IDH1 having His, Ser, Cys, Gly, Val, Pro or Leu at position 132 (*e.g.*, His); or a preselected IDH2 allele that encodes an IDH2 having Lys, Gly, Met, Trp, Thr, or Ser at position 172 and having a neoactivity disclosed herein, by administering to the subject a therapeutically effective amount of an inhibitor of IDH1 or IDH2 (*e.g.*, IDH1), *e.g.*, a small molecule or nucleic acid. The nucleic acid based inhibitor is, for example, a dsRNA, *e.g.*, a dsRNA that comprises the primary sequences of the sense strand and antisense strands of **Tables 7-14**. The dsRNA is composed of two separate strands, or a single strand folded to form a hairpin structure (*e.g.*, a short hairpin RNA (shRNA)). In some embodiments, the nucleic acid based inhibitor is an antisense nucleic acid, such as an antisense having a sequence that overlaps, or includes, an antisense sequence provided in **Tables 7-14**.

#### Neoactivity of an enzyme

Neoactivity, as used herein, means an activity that arises as a result of a mutation, *e.g.*, a point mutation, *e.g.*, a substitution, *e.g.*, in the active site of an enzyme. In an embodiment the neoactivity is substantially absent from wild type or non-mutant enzyme. This is sometimes referred to herein as a first degree neoactivity. An example of a first degree neoactivity is a “gain of function” wherein the mutant enzyme gains a new catalytic activity. In an embodiment the neoactivity is present in wild type or non-mutant enzyme but at a level which is less than 10, 5, 1, 0.1, 0.01 or

0.001 % of what is seen in the mutant enzyme. This is sometimes referred to herein as a second degree neoactivity. An example of a second degree neoactivity is a “gain of function” wherein the mutant enzyme has an increase, for example, a 5 fold increase in the rate of a catalytic activity possessed by the enzyme when lacking the mutation.

In some embodiments, a non-mutant form the enzyme, *e.g.*, a wild type form, converts substance A (*e.g.*, isocitrate) to substance B (*e.g.*,  $\alpha$ -ketoglutarate), and the neoactivity converts substance B (*e.g.*,  $\alpha$ -ketoglutarate) to substance C, sometimes referred to as the neoactivity product (*e.g.*, 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate). In some embodiments, the enzyme is in a metabolic pathway, *e.g.*, a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway, *e.g.*, IDH1 or IDH2.

In some embodiments, a non-mutant form the enzyme, *e.g.*, a wild type form, converts substance A to substance B, and the neoactivity converts substance B to substance A. In some embodiments, the enzyme is in a metabolic pathway, *e.g.*, a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway.

#### Isocitrate Dehydrogenases

Isocitrate dehydrogenases (IDHs) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate (*i.e.*,  $\alpha$ -ketoglutarate). These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer.

IDH1 (isocitrate dehydrogenase 1 (NADP+), cytosolic) is also known as IDH; IDP; IDCD; IDPC or PICD. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for

intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production.


The human IDH1 gene encodes a protein of 414 amino acids. The nucleotide and amino acid sequences for human IDH1 can be found as GenBank entries NM\_005896.2 and NP\_005887.2 respectively. The nucleotide and amino acid sequences for IDH1 are also described in, *e.g.*, Nekrutenko *et al.*, *Mol. Biol. Evol.* 15:1674-1684(1998); Geisbrecht *et al.*, *J. Biol. Chem.* 274:30527-30533(1999); Wiemann *et al.*, *Genome Res.* 11:422-435(2001); The MGC Project Team, *Genome Res.* 14:2121-2127(2004); Lubec *et al.*, Submitted (DEC-2008) to UniProtKB; Kullmann *et al.*, Submitted (JUN-1996) to the EMBL/GenBank/DDBJ databases; and Sjoebloom *et al.*, *Science* 314:268-274(2006).

IDH2 (isocitrate dehydrogenase 2 (NADP+), mitochondrial) is also known as IDH; IDP; IDHM; IDPM; ICD-M; or mNADP-IDH. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the mitochondria. It plays a role in intermediary metabolism and energy production. This protein may tightly associate or interact with the pyruvate dehydrogenase complex. Human IDH2 gene encodes a protein of 452 amino acids. The nucleotide and amino acid sequences for IDH2 can be found as GenBank entries NM\_002168.2 and NP\_002159.2 respectively. The nucleotide and amino acid sequence for human IDH2 are also described in, *e.g.*, Huh *et al.*, Submitted (NOV-1992) to the EMBL/GenBank/DDBJ databases; and The MGC Project Team, *Genome Res.* 14:2121-2127(2004).

Non-mutant, *e.g.*, wild type, IDH1 catalyzes the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate thereby reducing  $\text{NAD}^+$  ( $\text{NADP}^+$ ) to NADP (NADPH), *e.g.*, in the forward reaction:



In some embodiments, the neoactivity of a mutant IDH1 can have the ability to convert  $\alpha$ -ketoglutarate to 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate:



In some embodiments, the neoactivity can be the reduction of pyruvate or malate to the corresponding  $\alpha$ -hydroxyl compounds.

In some embodiments, the neoactivity of a mutant IDH1 can arise from a mutant IDH1 having a His, Ser, Cys, Gly, Val, Pro or Leu, or any other mutations described in Yan *et al.*, at residue 132 (e.g., His, Ser, Cys, Gly, Val or Leu; or His, Ser, Cys or Lys). In some embodiments, the neoactivity of a mutant IDH2 can arise from a mutant IDH2 having a Lys, Gly, Met, Trp, Thr, or Ser (e.g., Lys, Gly, Met, Trp, or Ser; or Gly, Met or Lys), or any other mutations described in Yan H *et al.*, at residue 172. Exemplary mutations include the following: R132H, R132C, R132S, R132G, R132L, and R132V.

In some embodiments, the mutant IDH1 and/or IDH2 (e.g., a mutant IDH1 and/or IDH2 having a neoactivity described herein) could lead to an increased level of 2-hydroxyglutarate, e.g., R-2-hydroxyglutarate in a subject. The accumulation of 2-hydroxyglutarate, e.g., R-2-hydroxyglutarate in a subject, e.g., in the brain of a subject, can be harmful. For example, in some embodiments, elevated levels of 2-hydroxyglutarate, e.g., R-2-hydroxyglutarate can lead to and/or be predictive of cancer in a subject such as a cancer of the central nervous system, e.g., brain tumor, e.g., glioma, e.g., glioblastoma multiforme (GBM). Accordingly, in some embodiments, a method described herein includes administering to a subject an inhibitor of the neoactivity.


#### Detection of 2-hydroxyglutarate

2-hydroxyglutarate can be detected, e.g., by LC/MS. To detect secreted 2-hydroxyglutarate in culture media, 500  $\mu$ L aliquots of conditioned media can be collected, mixed 80:20 with methanol, and centrifuged at 3,000 rpm for 20 minutes at 4 degrees Celsius. The resulting supernatant can be collected and stored at -80 degrees Celsius prior to LC-MS/MS to assess 2-hydroxyglutarate levels. To measure whole-cell associated metabolites, media can be aspirated and cells can be harvested, e.g., at a non-confluent density. A variety of different liquid chromatography (LC) separation methods can be used. Each method can be coupled by negative electrospray ionization (ESI, -3.0 kV) to triple-quadrupole mass spectrometers operating in multiple reaction monitoring (MRM) mode, with MS parameters optimized on infused metabolite standard solutions. Metabolites can be separated by reversed phase chromatography using 10 mM tributyl-amine as an ion pairing agent in the aqueous mobile phase, according to a variant of a previously reported method (Luo *et al. J Chromatogr A* 1147, 153-64, 2007). One method allows resolution of

TCA metabolites:  $t = 0$ , 50% B;  $t = 5$ , 95% B;  $t = 7$ , 95% B;  $t = 8$ , 0% B, where B refers to an organic mobile phase of 100% methanol. Another method is specific for 2-hydroxyglutarate, running a fast linear gradient from 50% -95% B (buffers as defined above) over 5 minutes. A Synergi Hydro-RP, 100mm  $\times$  2 mm, 2.1  $\mu$ m particle size (Phenomenex) can be used as the column, as described above.

Metabolites can be quantified by comparison of peak areas with pure metabolite standards at known concentration. Metabolite flux studies from  $^{13}\text{C}$ -glutamine can be performed as described, *e.g.*, in Munger *et al.* Nat Biotechnol 26, 1179-86, 2008.

In an embodiment 2HG, *e.g.*, R-2HG, is evaluated and the analyte on which the determination is based is 2HG, *e.g.*, R-2HG. In an embodiment the analyte on which the determination is based is a derivative of 2HG, *e.g.*, R-2HG, formed in process of performing the analytic method. By way of example such a derivative can be a derivative formed in MS analysis. Derivatives can include a salt adduct, *e.g.*, a Na adduct, a hydration variant, or a hydration variant which is also a salt adduct, *e.g.*, a Na adduct, *e.g.*, as formed in MS analysis. Exemplary 2HG derivatives include dehydrated derivatives such as the compounds provided below or a salt adduct thereof:



#### Methods of evaluating samples and/or subjects

This section provides methods of obtaining and analyzing samples and of analyzing subjects.

Embodiments of the method comprise evaluation of one or more parameters related to IDH, *e.g.*, IDH1 or IDH2, an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, to evaluate the IDH1 or IDH2 2HG neoactivity genotype or phenotype. The evaluation can be performed, *e.g.*, to select, diagnose or prognose the subject, to select a therapeutic agent, *e.g.*, an inhibitor, or to evaluate response to the treatment or progression of disease. In an embodiment the evaluation, which can be performed before and/or after treatment has begun, is based, at least in part, on analysis of a tumor sample, cancer cell sample, or precancerous cell sample, from the subject. *E.g.*, a sample from the patient can be analyzed for the presence or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, by evaluating a parameter correlated to the presence or level of an alpha hydroxy neoactivity product,

e.g., 2HG, *e.g.*, R-2HG. An alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG, in the sample can be determined by a chromatographic method, *e.g.*, by LC-MS analysis. It can also be determined by contact with a specific binding agent, *e.g.*, an antibody, which binds the alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG, and allows detection. In an embodiment the sample is analyzed for the level of neoactivity, *e.g.*, an alpha hydroxy neoactivity, e.g., 2HG neoactivity. In an embodiment the sample is analysed for the presence of a mutant IDH, *e.g.*, IDH1 or IDH2, protein having an alpha hydroxy neoactivity, e.g., 2HG neoactivity (or a corresponding RNA). *E.g.*, a mutant protein specific reagent, e.g., an antibody that specifically binds an IDH mutant protein, e.g., an antibody that specifically binds an IDH1-R132H mutant protein or an IDH2-R172 mutant protein (e.g., an IDH1-R132H mutant protein), can be used to detect neoactive mutant enzyme. In an embodiment a nucleic acid from the sample is sequenced to determine if a selected allele or mutation of IDH1 or IDH2 disclosed herein is present. In an embodiment the analysis is other than directly determining the presence of a mutant IDH, *e.g.*, IDH1 or IDH2, protein (or corresponding RNA) or sequencing of an IDH, *e.g.*, IDH1 or IDH2 gene. In an embodiment the analysis is other than directly determining, *e.g.*, it is other than sequencing genomic DNA or cDNA, the presence of a mutation at residue 132 of IDH1 and/or a mutation at residue 172 of IDH2. *E.g.*, the analysis can be the detection of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG, or the measurement of the mutation's an alpha hydroxy neoactivity, e.g., 2HG neoactivity. In an embodiment the sample is removed from the patient and analyzed. In an embodiment the evaluation can include one or more of performing the analysis of the sample, requesting analysis of the sample, requesting results from analysis of the sample, or receiving the results from analysis of the sample. (Generally herein, analysis can include one or both of performing the underlying method or receiving data from another who has performed the underlying method.)

In an embodiment the evaluation, which can be performed before and/or after treatment has begun, is based, at least in part, on analysis of a tissue (e.g., a tissue other than a tumor sample), or bodily fluid, or bodily product. Exemplary tissues include lymph node, skin, hair follicles and nails. Exemplary bodily fluids include blood, plasma, urine, lymph, tears, sweat, saliva, semen, and cerebrospinal fluid. Exemplary bodily products include exhaled breath. *E.g.*, the tissue, fluid or product can be analyzed for the presence or level of an alpha hydroxy neoactivity product, e.g.,

2HG, *e.g.*, R-2HG, by evaluating a parameter correlated to the presence or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. An alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, in the sample can be determined by a chromatographic method, *e.g.*, by LC-MS analysis. It can also be determined by contact with a specific binding agent, *e.g.*, an antibody, which binds the alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, and allows detection. In embodiments where sufficient levels are present, the tissue, fluid or product can be analyzed for the level of neoactivity, *e.g.*, an alpha hydroxy neoactivity, *e.g.*, the 2HG neoactivity. In an embodiment the sample is analysed for the presence of a mutant IDH, *e.g.*, IDH1 or IDH2, protein having an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity (or a corresponding RNA). *E.g.*, a mutant protein specific reagent, *e.g.*, an antibody that specifically binds an IDH mutant protein, *e.g.*, an antibody that specifically binds an IDH1-R132H mutant protein or an IDH2-R172 mutant protein (*e.g.*, an IDH1-R132H mutant protein), can be used to detect neoactive mutant enzyme. In an embodiment a nucleic acid from the sample is sequenced to determine if a selected allele or mutation of IDH1 or IDH2 disclosed herein is present. In an embodiment the analysis is other than directly determining the presence of a mutant IDH, *e.g.*, IDH1 or IDH2, protein (or corresponding RNA) or sequencing of an IDH, *e.g.*, IDH1 or IDH2 gene. *E.g.*, the analysis can be the detection of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, or the measurement of 2HG neoactivity. In an embodiment the tissue, fluid or product is removed from the patient and analyzed. In an embodiment the evaluation can include one or more of performing the analysis of the tissue, fluid or product, requesting analysis of the tissue, fluid or product, requesting results from analysis of the tissue, fluid or product, or receiving the results from analysis of the tissue, fluid or product.

In an embodiment the evaluation, which can be performed before and/or after treatment has begun, is based, at least in part, on alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, imaging of the subject. In embodiments magnetic resonance methods are used to evaluate the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, in the subject. In an embodiment the subject is subjected to imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS *e.g.*, analysis, and optionally an image corresponding to the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, or of the tumor, is formed. Optionally

the image or a value related to the image is stored in a tangible medium and/or transmitted to a second site. In an embodiment the evaluation can include one or more of performing imaging analysis, requesting imaging analysis, requesting results from imaging analysis, or receiving the results from imaging analysis.

#### Methods of treating a proliferative disorder

Described herein are methods of treating a cell proliferation-related disorder, *e.g.*, a cancer, *e.g.*, a glioma, *e.g.*, by inhibiting a neoactivity of a mutant enzyme, *e.g.*, an enzyme in a metabolic pathway, *e.g.*, a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway, *e.g.*, IDH1 or IDH2. The cancer can be characterized by the presence of a neoactivity, such as a gain of function in one or more mutant enzymes (*e.g.*, an enzyme in the metabolic pathway, *e.g.*, a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway *e.g.*, IDH1 or IDH2). In some embodiments, the gain of function is the conversion of  $\alpha$ -ketoglutarate to 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate.

#### Compounds for the treatment of cancer

A candidate compound can be evaluated for modulation (*e.g.*, inhibition) of neoactivity, for example, using an assay described herein. A candidate compound can also be evaluated for modulation (*e.g.*, inhibition) of wild type or non-mutant activity. For example, the formation of a product or by-product of any activity (*e.g.*, enzymatic activity) can be assayed, thus evaluating a candidate compound. In some embodiments, the activity (*e.g.*, wild type/non-mutant or neoactivity) can be evaluated by measuring one or more readouts from an enzymatic assay. For example, the change in nature and/or amount of substrate and/or product can be measured, *e.g.*, using methods such as fluorescent or radiolabeled substrates. Exemplary substrates and/or products include  $\alpha$ -ketoglutarate, CO<sub>2</sub>, NADP, NADPH, NAD, NADH, and 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate. In some embodiments, the rate of reaction of the enzyme can also be evaluated as can the nature and/or amount of a product of the enzymatic reaction. In addition to the measurement of potential enzymatic activities, activity (*e.g.*, wild type/non-mutant or neoactivity) can be



detected by the quenching of protein fluorescence upon binding of a potential substrate, cofactor, or enzymatic activity modulator to the enzyme.

In one embodiment, assay progress can be monitored by changes in the OD340 or fluorescence of the NAD or NADP cofactor. In another embodiment, the reaction progress can be coupled to a secondary enzyme assay system in continuous mode or endpoint mode for increasing the dynamic range of the assay. For example, an endpoint assay can be performed by adding to the reaction an excess of diaphorase and rezasarin. Diaphorase consumes the remaining NADPH or NADH while producing resorufin from rezasarin. Resorufin is a highly fluorescent product which can be measured by fluorescence at Ex544 Em590. This not only terminates the reaction but also generates an easily detectable signal with greater quantum yield than the fluorescence of the cofactor.

A continuous assay can be implemented through coupling a product of the primary reaction to a secondary enzyme reaction that yields detectable results of greater dynamic range or more convenient detection mode. For example, inclusion in the reaction mix of aldehyde dehydrogenase (ALDH), which is an NADP<sup>+</sup> dependent enzyme, and 6-methoxy-2-napthaldehyde, a chromogenic substrate for ALDH, will result in the production of the fluorescent product 6-methoxy-2-napthoate (Ex310 Em 360) at a rate dependent on the production of NADP<sup>+</sup> by isocitrate dehydrogenase. The inclusion of a coupling enzyme such as aldehyde dehydrogenase has the additional benefit of allowing screening of neoactivity irrespective of whether NADP<sup>+</sup> or NAD<sup>+</sup> is produced, since this enzyme is capable of utilizing both. Additionally, since the NADPH or NADH cofactor required for the “reverse” assay is regenerated, a coupled enzyme system which cycles the cofactor back to the IDH enzyme has the further advantage of permitting continuous assays to be conducted at cofactor concentrations much below K<sub>m</sub> for the purpose of enhancing the detection of competitive inhibitors of cofactor binding.

In yet a third embodiment of an activity (*e.g.*, wild type/non-mutant or neoactivity) screen, one or a number of IDH substrates, cofactors, or products can be isotopically labeled with radioactive or “heavy” elements at defined atoms for the purpose of following specific substrates or atoms of substrates through the chemical reaction. For example, the alpha carbon of α-KG, isocitrate, or 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate may be <sup>14</sup>C or <sup>13</sup>C. Amount, rate, identity and structure of


products formed can be analyzed by means known to those of skill in the art, for example mass spectroscopy or radiometric HPLC.

Compounds that inhibit a neoactivity, *e.g.*, a neoactivity described herein, can include, *e.g.*, small molecule, nucleic acid, protein and antibody.

Exemplary small molecules include, *e.g.*, small molecules that bind to enzymes and decrease their activity, *e.g.*, a neoactivity described herein. The binding of an inhibitor can stop a substrate from entering the enzyme's active site and/or hinder the enzyme from catalyzing its reaction. Inhibitor binding is either reversible or irreversible. Irreversible inhibitors usually react with the enzyme and change it chemically. These inhibitors can modify key amino acid residues needed for enzymatic activity. In contrast, reversible inhibitors bind non-covalently and different types of inhibition are produced depending on whether these inhibitors bind the enzyme, the enzyme-substrate complex, or both.

In some embodiments, the small molecule is oxalomalate, oxalofumarate, or oxalosuccinate.

In some embodiments, the small molecule is a compound of formula (X), or a compound as listed in **Table 24a**. The compound of formula (X) is provided below:



Formula (X)


wherein X is C<sub>1</sub>-C<sub>6</sub> alkylene (*e.g.*, methylene), C(O), or C(O)C<sub>1</sub>-C<sub>6</sub> alkylene;

wherein X is optionally substituted;

R<sup>1</sup> is halo (*e.g.*, fluoro), C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, hydroxyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, cyano, nitro, amino, alkylamino, dialkylamino, amido, -C(O)OH, or C(O)OC<sub>1</sub>-C<sub>6</sub>alkyl; and

m is 0, 1, 2, or 3.

In some embodiments, the compound is a compound of formula (XI) or a pharmaceutically acceptable salt thereof or a compound listed in Table 24b



wherein:

W, X, Y and Z are each independently selected from CH or N;  
B and B<sup>1</sup> are independently selected from hydrogen, alkyl or when taken together with the carbon to which they are attached form a carbonyl group;  
Q is C=O or SO<sub>2</sub>;  
D and D<sup>1</sup> are independently selected from a bond, oxygen or NR<sup>c</sup>;  
A is optionally substituted aryl or optionally substituted heteroaryl;  
R<sup>1</sup> is independently selected from alkyl, acyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, heterocyclylalkyl, cycloalkylalkyl, aralkyl, and heteroaralkyl; each of which may be optionally substituted with 0-3 occurrences of R<sup>d</sup>;  
each R<sup>3</sup> is independently selected from halo, haloalkyl, alkyl and -OR<sup>a</sup>;  
each R<sup>a</sup> is independently selected from alkyl, and haloalkyl;  
each R<sup>b</sup> is independently alkyl;  
each R<sup>c</sup> is independently selected from hydrogen, alkyl and alkenyl;  
each R<sup>d</sup> is independently selected from halo, haloalkyl, alkyl, nitro, cyano, and -OR<sup>a</sup>, or two R<sup>d</sup> taken together with the carbon atoms to which they are attached form an optionally substituted heterocyclyl;  
n is 0, 1, or 2;  
h is 0, 1, 2; and  
g is 0, 1 or 2.

In some embodiments, the small molecule is a selective inhibitor of the neoactivity (*e.g.*, relative to the wild type activity).

Nucleic acids can be used to inhibit a neoactivity, *e.g.*, a neoactivity described herein, *e.g.*, by decreasing the expression of the enzyme. Exemplary nucleic acids include, *e.g.*, siRNA, shRNA, antisense RNA, aptamer and ribozyme. Art-known methods can be used to select inhibitory molecules, *e.g.*, siRNA molecules, for a particular gene sequence.

Proteins can also be used to inhibit a neoactivity, *e.g.*, a neoactivity described herein, by directly or indirectly binding to the enzyme and/or substrate, or competing binding to the enzyme and/or substrate. Exemplary proteins include, *e.g.*, soluble receptors, peptides and antibodies. Exemplary antibodies include, *e.g.*, whole antibody or a fragment thereof that retains its ability to bind to the enzyme or substrate.

Exemplary candidate compounds, which can be tested for inhibition of a neoactivity described herein (*e.g.*, a neoactivity associated with mutant IDH1), are

described in the following references, each of which are incorporated herein by reference: Bioorganic & Medicinal Chemistry (2008), 16(7), 3580-3586; Free Radical Biology & Medicine (2007), 42(1), 44-51; KR 2005036293 A; Applied and Environmental Microbiology (2005), 71(9), 5465-5475; KR 2002095553 A; U.S. Pat. Appl. US 2004067234 A1; PCT Int. Appl. (2002), WO 2002033063 A1; Journal of Organic Chemistry (1996), 61(14), 4527-4531; Biochimica et Biophysica Acta, Enzymology (1976), 452(2), 302-9; Journal of Biological Chemistry (1975), 250(16), 6351-4; Bollettino - Societa Italiana di Biologia Sperimentale (1972), 48(23), 1031-5; Journal of Biological Chemistry (1969), 244(20), 5709-12.

#### Isomers

Certain compounds may exist in one or more particular geometric, optical, enantiomeric, diastereomeric, epimeric, atropic, stereoisomer, tautomeric, conformational, or anomeric forms, including but not limited to, cis- and trans-forms; E- and Z-forms; c-, t-, and r- forms; endo- and exo-forms; R-, S-, and meso-forms; D- and L-forms; d- and l-forms; (+) and (-) forms; keto-, enol-, and enolate-forms; syn- and anti-forms; synclinal- and anticlinal-forms;  $\alpha$ - and  $\beta$ -forms; axial and equatorial forms; boat-, chair-, twist-, envelope-, and halfchair-forms; and combinations thereof, hereinafter collectively referred to as "isomers" (or "isomeric forms").

In one embodiment, a compound described herein, *e.g.*, an inhibitor of a neoactivity or 2-HG is an enantiomerically enriched isomer of a stereoisomer described herein. For example, the compound has an enantiomeric excess of at least about 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%. Enantiomer, when used herein, refers to either of a pair of chemical compounds whose molecular structures have a mirror-image relationship to each other.


In one embodiment, a preparation of a compound disclosed herein is enriched for an isomer of the compound having a selected stereochemistry, *e.g.*, R or S, corresponding to a selected stereocenter, *e.g.*, the 2-position of 2-hydroxyglutaric acid. 2HG can be purchased from commercial sources or can be prepared using methods known in the art, for example, as described in Org. Syn. Coll vol., 7, P-99, 1990. For example, the compound has a purity corresponding to a compound having a selected stereochemistry of a selected stereocenter of at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%.

In one embodiment, a composition described herein includes a preparation of a compound disclosed herein that is enriched for a structure or structures having a selected stereochemistry, *e.g.*, R or S, at a selected stereocenter, *e.g.*, the 2-position of 2-hydroxyglutaric acid. Exemplary R/S configurations can be those provided in an example described herein.

An "enriched preparation," as used herein, is enriched for a selected stereoconfiguration of one, two, three or more selected stereocenters within the subject compound. Exemplary selected stereocenters and exemplary stereoconfigurations thereof can be selected from those provided herein, *e.g.*, in an example described herein. By enriched is meant at least 60%, *e.g.*, of the molecules of compound in the preparation have a selected stereochemistry of a selected stereocenter. In an embodiment it is at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%. Enriched refers to the level of a subject molecule(s) and does not connote a process limitation unless specified.

Note that, except as discussed below for tautomeric forms, specifically excluded from the term "isomers," as used herein, are structural (or constitutional) isomers (*i.e.*, isomers which differ in the connections between atoms rather than merely by the position of atoms in space). For example, a reference to a methoxy group, -OCH<sub>3</sub>, is not to be construed as a reference to its structural isomer, a hydroxymethyl group, -CH<sub>2</sub>OH. Similarly, a reference to ortho-chlorophenyl is not to be construed as a reference to its structural isomer, meta-chlorophenyl. However, a reference to a class of structures may well include structurally isomeric forms falling within that class (*e.g.*, C1-7alkyl includes n-propyl and iso-propyl; butyl includes n-, iso-, sec-, and tert-butyl; methoxyphenyl includes ortho-, meta-, and para-methoxyphenyl).

The above exclusion does not pertain to tautomeric forms, for example, keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, N-nitroso/hydroxyazo, and nitro/aci-nitro.



Note that specifically included in the term "isomer" are compounds with one or more isotopic substitutions. For example, H may be in any isotopic form, including  $1\text{H}$ ,  $2\text{H}$  (D), and  $3\text{H}$  (T); C may be in any isotopic form, including  $12\text{C}$ ,  $13\text{C}$ , and  $14\text{C}$ ; O may be in any isotopic form, including  $16\text{O}$  and  $18\text{O}$ ; and the like. Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including (wholly or partially) racemic and other mixtures thereof. Methods for the preparation (*e.g.*, asymmetric synthesis) and separation (*e.g.*, fractional crystallisation and chromatographic means) of such isomeric forms are either known in the art or are readily obtained by adapting the methods taught herein, or known methods, in a known manner.

### Salts

It may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the active compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge *et al.*, 1977, "Pharmaceutically Acceptable Salts." J. Pharm. Sci. Vol. 66, pp. 1-19.

For example, if the compound is anionic, or has a functional group which may be anionic (*e.g.*,  $-\text{COOH}$  may be  $-\text{COO}^-$ ), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as  $\text{Na}^+$  and  $\text{K}^+$ , alkaline earth cations such as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , and other cations such as  $\text{Al}^{3+}$ . Examples of suitable organic cations include, but are not limited to, ammonium ion (*i.e.*,  $\text{NH}_4^+$ ) and substituted ammonium ions (*e.g.*,  $\text{NH}_3\text{R}^+$ ,  $\text{NH}_2\text{R}_2^+$ ,  $\text{NHR}_3^+$ ,  $\text{NR}_4^+$ ). Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is  $\text{N}(\text{CH}_3)_4^+$ .

If the compound is cationic, or has a functional group that may be cationic (*e.g.*,  $-\text{NH}_2$  may be  $-\text{NH}_3^+$ ), then a salt may be formed with a suitable anion. Examples of suitable inorganic anions include, but are not limited to, those derived from the following inorganic acids: hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfurous, nitric, nitrous, phosphoric, and phosphorous.

Examples of suitable organic anions include, but are not limited to, those derived from the following organic acids: 2-acetyoxybenzoic, acetic, ascorbic, aspartic, benzoic, camphorsulfonic, cinnamic, citric, edetic, ethanedisulfonic, ethanesulfonic, fumaric, gluceptonic, gluconic, glutamic, glycolic, hydroxymaleic, hydroxynaphthalene carboxylic, isethionic, lactic, lactobionic, lauric, maleic, malic, methanesulfonic, mucic, oleic, oxalic, palmitic, pamoic, pantothenic, phenylacetic, phenylsulfonic, propionic, pyruvic, salicylic, stearic, succinic, sulfanilic, tartaric, toluenesulfonic, and valeric. Examples of suitable polymeric organic anions include, but are not limited to, those derived from the following polymeric acids: tannic acid, carboxymethyl cellulose.

Unless otherwise specified, a reference to a particular compound also includes salt forms thereof.

#### Chemically Protected Forms

It may be convenient or desirable to prepare, purify, and/or handle the active compound in a chemically protected form. The term "chemically protected form" is used herein in the conventional chemical sense and pertains to a compound in which one or more reactive functional groups are protected from undesirable chemical reactions under specified conditions (*e.g.*, pH, temperature, radiation, solvent, and the like). In practice, well known chemical methods are employed to reversibly render unreactive a functional group, which otherwise would be reactive, under specified conditions. In a chemically protected form, one or more reactive functional groups are in the form of a protected or protecting group (also known as a masked or masking group or a blocked or blocking group). By protecting a reactive functional group, reactions involving other unprotected reactive functional groups can be performed, without affecting the protected group; the protecting group may be removed, usually in a subsequent step, without substantially affecting the remainder of the molecule. See, for example, *Protective Groups in Organic Synthesis* (T. Green and P. Wuts; 3rd Edition; John Wiley and Sons, 1999). Unless otherwise specified, a reference to a particular compound also includes chemically protected forms thereof.

A wide variety of such "protecting," "blocking," or "masking" methods are widely used and well known in organic synthesis. For example, a compound which has two nonequivalent reactive functional groups, both of which would be reactive under specified conditions, may be derivatized to render one of the functional groups

"protected," and therefore unreactive, under the specified conditions; so protected, the compound may be used as a reactant which has effectively only one reactive functional group. After the desired reaction (involving the other functional group) is complete, the protected group may be "deprotected" to return it to its original functionality.

For example, a hydroxy group may be protected as an ether (-OR) or an ester (-OC(=O)R), for example, as: a t-butyl ether; a benzyl, benzhydryl (diphenylmethyl), or trityl (triphenylmethyl) ether; a trimethylsilyl or t-butyldimethylsilyl ether; or an acetyl ester (-OC(=O)CH<sub>3</sub>, -OAc).

For example, an aldehyde or ketone group may be protected as an acetal (R-CH(OR)<sub>2</sub>) or ketal (R<sub>2</sub>C(OR)<sub>2</sub>), respectively, in which the carbonyl group (>C=O) is converted to a diether (>C(OR)<sub>2</sub>), by reaction with, for example, a primary alcohol. The aldehyde or ketone group is readily regenerated by hydrolysis using a large excess of water in the presence of acid.

For example, an amine group may be protected, for example, as an amide (-NRCO-R) or a urethane (-NRCO-OR), for example, as: a methyl amide (-NHCO-CH<sub>3</sub>); a benzyloxy amide (-NHCO-OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH-Cbz); as a t-butoxy amide (-NHCO-OC(CH<sub>3</sub>)<sub>3</sub>, -NH-Boc); a 2-biphenyl-2-propoxy amide (-NHCO-OC(CH<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>C<sub>6</sub>H<sub>5</sub>, -NH-Bpoc), as a 9-fluorenylmethoxy amide (-NH-Fmoc), as a 6-nitroveratryloxy amide (-NH-Nvoc), as a 2-trimethylsilylethyloxy amide (-NH-Teoc), as a 2,2,2-trichloroethyloxy amide (-NH-Troc), as an allyloxy amide (-NH-Alloc), as a 2-(phenylsulphonyl)ethyloxy amide (-NH-Psec); or, in suitable cases (*e.g.*, cyclic amines), as a nitroxide radical (>N-O<).

For example, a carboxylic acid group may be protected as an ester for example, as: an C<sup>α</sup>alkyl ester (*e.g.*, a methyl ester; a t-butyl ester); a C<sup>γ</sup>haloalkyl ester (*e.g.*, a C<sub>1-7</sub>trihaloalkyl ester); a triC<sub>1-7</sub>alkylsilyl-Ci<sub>7</sub>alkyl ester; or a C<sub>5-20</sub>aryl-C<sub>1-7</sub>alkyl ester (*e.g.*, a benzyl ester; a nitrobenzyl ester); or as an amide, for example, as a methyl amide.

For example, a thiol group may be protected as a thioether (-SR), for example, as: a benzyl thioether; an acetamidomethyl ether (-S-CH<sub>2</sub>NHC(=O)CH<sub>3</sub>).

#### Nucleic acid based inhibitors

Nucleic acid-based inhibitors for inhibition IDH, *e.g.*, IDH1, can be, *e.g.*, double stranded RNA (dsRNA) that function, *e.g.*, by an RNA interference (RNAi



mechanism), an antisense RNA, or a microRNA (miRNA). In an embodiment the nucleic-acid based inhibitor binds to the target mRNA and inhibits the production of protein therefrom, *e.g.*, by cleavage of the target mRNA.

*Double stranded RNA (dsRNA)*

A nucleic acid based inhibitor useful for decreasing IDH1 or IDH2 mutant function is, *e.g.*, a dsRNA, such as a dsRNA that acts by an RNAi mechanism. RNAi refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs). dsRNAs as used herein are understood to include siRNAs. Typically, inhibition of IDH, *e.g.*, IDH1, by dsRNAs does not trigger the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2',5'-oligoadenylate synthetase resulting in non-specific cleavage of mRNA by ribonuclease L.

dsRNAs targeting an IDH, *e.g.*, IDH1, enzyme, *e.g.*, a wildtype or mutant IDH1, can be unmodified or chemically modified. The dsRNA can be chemically synthesized, expressed from a vector or enzymatically synthesized. The invention also features various chemically modified synthetic dsRNA molecules capable of modulating IDH1 gene expression or activity in cells by RNA interference (RNAi). The use of chemically modified dsRNA improves various properties of native dsRNA molecules, such as through increased resistance to nuclease degradation *in vivo* and/or through improved cellular uptake.

The dsRNAs targeting nucleic acid can be composed of two separate RNAs, or of one RNA strand, which is folded to form a hairpin structure. Hairpin dsRNAs are typically referred to as shRNAs.

An shRNA that targets IDH, *e.g.*, a mutant or wildtype IDH1 gene can be expressed from a vector, *e.g.*, viral vector, such as a lentiviral or adenoviral vector. In certain embodiments, a suitable dsRNA for inhibiting expression of an IDH1 gene will be identified by screening an siRNA library, such as an adenoviral or lentiviral siRNA library.

In an embodiment, a dsRNA that targets IDH, *e.g.*, IDH1, is about 15 to about 30 base pairs in length (*e.g.*, about 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, or 29) basepairs in length. In another embodiment, the dsRNA includes overhanging ends of about 1 to about 3 (*e.g.*, about 1, 2, or 3) nucleotides. By "overhang" is meant that 3'-end of one strand of the dsRNA extends beyond the 5'-end of the other strand,

or vice versa. The dsRNA can have an overhang on one or both ends of the dsRNA molecule. In some embodiments, the single-stranded overhang is located at the 3'-terminal end of the antisense strand, or, alternatively, at the 3'-terminal end of the sense strand. In some embodiments, the overhang is a TT or UU dinucleotide overhang, *e.g.*, a TT or UU dinucleotide overhang. For example, in an embodiment, the dsRNA includes a 21-nucleotide antisense strand, a 19 base pair duplex region, and a 3'-terminal dinucleotide. In yet another embodiment, a dsRNA includes a duplex nucleic acid where both ends are blunt, or alternatively, where one of the ends is blunt.

In an embodiment, the dsRNA includes a first and a second strand, each strand is about 18 to about 28 nucleotides in length, *e.g.*, about 19 to about 23 nucleotides in length, the first strand of the dsRNA includes a nucleotide sequence having sufficient complementarity to the IDH, *e.g.*, IDH1, RNA for the dsRNA to direct cleavage of the IDH, *e.g.*, IDH1, mRNA via RNA interference, and the second strand of the dsRNA includes a nucleotide sequence that is complementary to the first strand.

In an embodiment, a dsRNA targeting an IDH, *e.g.*, IDH1, gene can target wildtype and mutant forms of the gene, or can target different allelic isoforms of the same gene. For example, the dsRNA will target a sequence that is identical in two or more of the different isoforms. In an embodiment, the dsRNA targets an IDH1 having G at position 395 or C at position 394 (*e.g.*, a wildtype IDH1 RNA) and an IDH1 having A at position 395 or A at position 394, such as a C394A, a C394G, a C394T, a G395C, a G395T or a G395A mutation, (*e.g.*, an IDH1 RNA carrying a G395A and/or a C394A mutation) (**FIG. 2**).

In an embodiment, a dsRNA will preferentially or specifically target a mutant IDH RNA, or a particular IDH polymorphism. In some embodiments, the IDH has a mutation at position 394 or 395 such as a C394A, a C394G, a C394T, a G395C, a G395T or a G395A mutation. For example, in an embodiment, the dsRNA targets an IDH1 RNA carrying an A at position 395, *e.g.*, G395A, and in another embodiment, the dsRNA targets an IDH1 RNA carrying an A at position 394, *e.g.*, C394A mutation.

In an embodiment, a dsRNA targeting an IDH RNA includes one or more chemical modifications. Non-limiting examples of such chemical modifications include without limitation phosphorothioate internucleotide linkages, 2'-deoxyribonucleotides, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl nucleotides, and

terminal glyceryl and/or inverted deoxy abasic residue incorporation. Such chemical modifications have been shown to preserve RNAi activity in cells while at the same time, dramatically increasing the serum stability of these compounds. Furthermore, one or more phosphorothioate substitutions are well-tolerated and have been shown to confer substantial increases in serum stability for modified dsRNA constructs.

In an embodiment, a dsRNA targeting an IDH, *e.g.*, IDH1, RNA includes modified nucleotides while maintaining the ability to mediate RNAi. The modified nucleotides can be used to improve *in vitro* or *in vivo* characteristics such as stability, activity, and/or bioavailability. For example, the dsRNA can include modified nucleotides as a percentage of the total number of nucleotides present in the molecule. As such, the dsRNA can generally include about 5% to about 100% modified nucleotides (*e.g.*, about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% modified nucleotides).

In some embodiments, the dsRNA targeting IDH, *e.g.*, IDH1, is about 21 nucleotides long. In another embodiment, the dsRNA does not contain any ribonucleotides, and in another embodiment, the dsRNA includes one or more ribonucleotides. In an embodiment, each strand of the dsRNA molecule independently includes about 15 to about 30 (*e.g.*, about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30) nucleotides, wherein each strand includes about 15 to about 30 (*e.g.*, about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30) nucleotides that are complementary to the nucleotides of the other strand. In an embodiment, one of the strands of the dsRNA includes a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of the IDH1 or IDH2 gene, and the second strand of the dsRNA includes a nucleotide sequence substantially similar to the nucleotide sequence of the IDH1 or IDH2 gene or a portion thereof.

In an embodiment, the dsRNA targeting IDH1 or IDH2 includes an antisense region having a nucleotide sequence that is complementary to a nucleotide sequence of the IDH1 or IDH2 gene or a portion thereof, and a sense region having a nucleotide sequence substantially similar to the nucleotide sequence of the IDH1 or IDH2 gene or a portion thereof. In an embodiment, the antisense region and the sense region independently include about 15 to about 30 (*e.g.*, about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30) nucleotides, where the antisense region includes

about 15 to about 30 (*e.g.*, about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30) nucleotides that are complementary to nucleotides of the sense region.

As used herein, the term “dsRNA” is meant to include nucleic acid molecules that are capable of mediating sequence specific RNAi, such as short interfering RNA (siRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. In addition, as used herein, the term “RNAi” is meant to include sequence specific RNA interference, such as post transcriptional gene silencing, translational inhibition, or epigenetics.

Nucleic acid-based IDH inhibitors

In an embodiment the inhibitor is a nucleic acid-based inhibitor, such as a double stranded RNA (dsRNA) or antisense RNA that targets a mutant IDH, *e.g.*, mutant IDH1 or IDH2.

In one embodiment, the nucleic acid based inhibitor, *e.g.*, a dsRNA or antisense molecule, decreases or inhibits expression of an IDH1 having other than an Arg, *e.g.*, having a His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Yan *et al.*, N. Eng. J. Med. 360:765-73, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**). In one embodiment, the nucleic acid based inhibitor decreases or inhibits expression of an IDH1 enzyme having His at residue 132

In an embodiment the nucleic acid-based inhibitor is a dsRNA that targets an mRNA that encodes an IDH1 allele described herein, *e.g.*, an IDH1 allele having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **Fig. 21**).

In an embodiment the allele encodes an IDH1 having His at residue 132.

In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment, the nucleic acid-based inhibitor is a dsRNA that targets IDH1, *e.g.*, an IDH1 having an A or a T (or a nucleotide other than C) at nucleotide position 394 or an A (or a nucleotide other than G) at nucleotide position 395, *e.g.*, a mutant allele carrying a C394T mutation or a G395A mutation according to the IDH1 sequence of SEQ ID NO:8 (see also Fig 21A).

In an embodiment, the dsRNA targets an IDH1 having other than C, *e.g.*, a T or an A, at nucleotide position 394 or and other than G, *e.g.*, an A, at 395 (*e.g.*, a mutant) and an IDH1 having a C at nucleotide position 394 or a G at nucleotide position 395 (*e.g.*, a wildtype), *e.g.*, by targeting a region of the IDH1 mRNA that is identical between the wildtype and mutant transcripts. In yet another embodiment, the dsRNA targets a particular mutant or polymorphism (such as a single nucleotide polymorphism (SNP)), but not a wildtype allele. In this case, the nucleic acid based inhibitor, *e.g.*, a dsRNA, targets the region of the IDH1 containing the mutation.

In some embodiments, the nucleic acid based inhibitor, *e.g.*, a dsRNA preferentially or specifically inhibits the product of a mutant IDH1 as compared to the product of a wildtype IDH1. In some embodiments, the IDH has a mutation at position 394 or 395 such as a C394A, a C394G, a C394T, a G395C, a G395T or a G395A mutation. For example, in one embodiment, a dsRNA targets a region of an IDH1 mRNA that carries the mutation (*e.g.*, a C394A of C394T or a G395A mutation according to SEQ ID NO:5).

In one embodiment, the nucleic acid-based inhibitor is a dsRNA including a sense strand and an antisense strand having a primary sequence presented in **Tables 7- 14**. In another embodiment, the nucleic acid based inhibitor is an antisense oligonucleotide that includes all or a part of an antisense primary sequence presented in **Tables 7- 14** or which targets the same or substantially the same region as does a dsRNA from **Tables 7- 14**.

In one embodiment, the nucleic acid based inhibitor decreases or inhibits expression of an IDH2 having Lys, Gly, Met, Trp, Thr, Ser, or any residue described in Yan *et al.*, at residue 172, according to the amino acid sequence of SEQ ID NO:10 (see also **FIG. 22**). In one embodiment, the nucleic acid based inhibitor decreases or inhibits expression of an IDH2 enzyme having Lys at residue 172.

In an embodiment the nucleic acid-based inhibitor is a dsRNA that targets an mRNA that encodes an IDH2 allele described herein, *e.g.*, an IDH2 allele having other than an Arg at residue 172. *E.g.*, the allele can have Lys, Gly, Met, Trp, Thr, Ser, or any residue described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10 (see also **Fig. 22**).

In an embodiment the allele encodes an IDH2 having Lys at residue 172.

In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment, the nucleic acid-based inhibitor is a dsRNA that targets IDH2, *e.g.*, an IDH2 having a G or a T (or a nucleotide other than A or C) at nucleotide position 514 or an A or T or C (or a nucleotide other than G) at nucleotide position 515, *e.g.*, a mutant allele carrying a A514G mutation or a G515T or a G515A mutation according to the IDH2 sequence of SEQ ID NO:10 (**Fig. 22A**). In one embodiment, the nucleic acid-based inhibitor is a dsRNA that targets IDH2, *e.g.*, an IDH2 having a C or a T (or a nucleotide other than G or A) at nucleotide position 516 according to the IDH2 sequence of SEQ ID NO:10.

In an embodiment, the nucleic acid-based inhibitor is a dsRNA that targets IDH2, *e.g.*, an IDH2 having a G at nucleotide position 514 or a T at nucleotide position 515 or an A at position 515, according to the IDH2 sequence of SEQ ID NO:10.

In an embodiment, the dsRNA targets an IDH2 having other than A, *e.g.*, a G or a T, at nucleotide position 514, or other than G, *e.g.*, an A or C or T at position 515 (*e.g.*, a mutant), or other than G, *e.g.*, C or T, and an IDH2 having an A at nucleotide position 514 or a G at nucleotide position 515 or a G at position 516 (*e.g.*, a wildtype), *e.g.*, by targeting a region of the IDH2 mRNA that is identical between the wildtype and mutant transcripts. In yet another embodiment, the dsRNA targets a particular mutant or polymorphism (such as a single nucleotide polymorphism (SNP)), but not a wildtype allele. In this case, the nucleic acid based inhibitor, *e.g.*, a dsRNA, targets the region of the IDH2 containing the mutation.

In some embodiments, the nucleic acid based inhibitor, *e.g.*, a dsRNA, preferentially or specifically inhibits the product of a mutant IDH2 as compared to the product of a wildtype IDH2. For example, in one embodiment, a dsRNA targets a region of an IDH2 mRNA that carries the mutation (*e.g.*, an A514G or G515T or a G515U mutation according to SEQ ID NO:10).

In one embodiment, the nucleic acid-based inhibitor is a dsRNA including a sense strand and an antisense strand having a primary sequence presented in **Tables 15-23**. In another embodiment, the nucleic acid based inhibitor is an antisense oligonucleotide that includes all or a part of an antisense primary sequence presented in **Tables 15-23** or which targets the same or substantially the same region as does a dsRNA from **Tables 15-23**.

In an embodiment, the nucleic acid based inhibitor is delivered to the brain, *e.g.*, directly to the brain, *e.g.*, by intrathecal or intraventricular delivery. The nucleic

acid based inhibitor can also be delivered from an implantable device. In an embodiment, the nucleic acid-based inhibitor is delivered by infusion using, *e.g.*, a catheter, and optionally, a pump.

*Antisense*

Suitable nucleic acid based inhibitors include antisense nucleic acids. While not being bound by theory it is believed that antisense inhibition is typically based upon hydrogen bonding-based hybridization of oligonucleotide strands or segments such that at least one strand or segment is cleaved, degraded, or otherwise rendered inoperable.

An antisense agent can bind IDH1 or IDH2 DNA. In embodiments it inhibits replication and transcription. While not being bound by theory it is believed that an antisense agent can also function to inhibit target RNA translocation, *e.g.*, to a site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more RNA species, and catalytic activity or complex formation involving the RNA.

An antisense agents can have a chemical modification described above as being suitable for dsRNA.

Antisense agents can include, for example, from about 8 to about 80 nucleobases (*i.e.*, from about 8 to about 80 nucleotides), *e.g.*, about 8 to about 50 nucleobases, or about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression. Anti-sense compounds can include a stretch of at least eight consecutive nucleobases that are complementary to a sequence in the target gene. An oligonucleotide need not be 100% complementary to its target nucleic acid sequence to be specifically hybridizable. An oligonucleotide is specifically hybridizable when binding of the oligonucleotide to the target interferes with the normal function of the target molecule to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the oligonucleotide to non-target sequences under conditions in which specific binding is desired, *i.e.*, under physiological conditions in the case of *in vivo* assays or therapeutic treatment or, in the case of *in vitro* assays, under conditions in which the assays are conducted.

Hybridization of antisense oligonucleotides with mRNA (*e.g.*, an mRNA encoding IDH1 or IDH2) can interfere with one or more of the normal functions of mRNA. While not being bound by theory it is believed that the functions of mRNA to be interfered with include all key functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in by the RNA. Binding of specific protein(s) to the RNA may also be interfered with by antisense oligonucleotide hybridization to the RNA.

Exemplary antisense compounds include DNA or RNA sequences that specifically hybridize to the target nucleic acid, *e.g.*, the mRNA encoding IDH1 or IDH2. The complementary region can extend for between about 8 to about 80 nucleobases. The compounds can include one or more modified nucleobases. Modified nucleobases may include, *e.g.*, 5-substituted pyrimidines such as 5-iodouracil, 5-iodocytosine, and C5-propynyl pyrimidines such as C5-propynylcytosine and C5-propynyluracil. Other suitable modified nucleobases include  $N^4$ -(C<sub>1</sub>-C<sub>12</sub>) alkylaminocytosines and  $N^4,N^4$ -(C<sub>1</sub>-C<sub>12</sub>) dialkylaminocytosines. Modified nucleobases may also include 7-substituted-5-aza-7-deazapurines and 7-substituted-7-deazapurines such as, for example, 7-iodo-7-deazapurines, 7-cyano-7-deazapurines, 7-aminocarbonyl-7-deazapurines. Examples of these include 6-amino-7-iodo-7-deazapurines, 6-amino-7-cyano-7-deazapurines, 6-amino-7-aminocarbonyl-7-deazapurines, 2-amino-6-hydroxy-7-iodo-7-deazapurines, 2-amino-6-hydroxy-7-cyano-7-deazapurines, and 2-amino-6-hydroxy-7-aminocarbonyl-7-deazapurines. Furthermore,  $N^6$ -(C<sub>1</sub>-C<sub>12</sub>) alkylaminopurines and  $N^6,N^6$ -(C<sub>1</sub>-C<sub>12</sub>) dialkylaminopurines, including  $N^6$ -methylaminoadenine and  $N^6,N^6$ -dimethylaminoadenine, are also suitable modified nucleobases. Similarly, other 6-substituted purines including, for example, 6-thioguanine may constitute appropriate modified nucleobases. Other suitable nucleobases include 2-thiouracil, 8-bromoadenine, 8-bromoguanine, 2-fluoroadenine, and 2-fluoroguanine. Derivatives of any of the aforementioned modified nucleobases are also appropriate. Substituents of any of the preceding compounds may include C<sub>1</sub>-C<sub>30</sub> alkyl, C<sub>2</sub>-C<sub>30</sub> alkenyl, C<sub>2</sub>-C<sub>30</sub> alkynyl, aryl, aralkyl, heteroaryl, halo, amino, amido, nitro, thio, sulfonyl, carboxyl, alkoxy, alkylcarbonyl, alkoxy carbonyl, and the like.

### *MicroRNA*



In some embodiments, the nucleic acid-based inhibitor suitable for targeting IDH, *e.g.*, IDH1, is a microRNA (miRNA). A miRNA is a single stranded RNA that regulates the expression of target mRNAs either by mRNA cleavage, translational repression/inhibition or heterochromatic silencing. The miRNA is 18 to 25 nucleotides, typically 21 to 23 nucleotides in length. In some embodiments, the miRNA includes chemical modifications, such as one or more modifications described herein.

In some embodiments, a nucleic acid based inhibitor targeting IDH has partial complementarity (*i.e.*, less than 100% complementarity) with the target IDH, *e.g.*, IDH1 or IDH2, mRNA. For example, partial complementarity can include various mismatches or non-base paired nucleotides (*e.g.*, 1, 2, 3, 4, 5 or more mismatches or non-based paired nucleotides, such as nucleotide bulges), which can result in bulges, loops, or overhangs that result between the antisense strand or antisense region of the nucleic acid-based inhibitor and the corresponding target nucleic acid molecule.

The nucleic acid-based inhibitors described herein, *e.g.*, antisense nucleic acid described herein, can be incorporated into a gene construct to be used as a part of a gene therapy protocol to deliver nucleic acids that can be used to express and produce agents within cells. Expression constructs of such components may be administered in any biologically-effective carrier, *e.g.*, any formulation or composition capable of effectively delivering the component gene to cells *in vivo*. Approaches include insertion of the subject gene in viral vectors including recombinant retroviruses, adenovirus, adeno-associated virus, lentivirus, and herpes simplex virus-1, or recombinant bacterial or eukaryotic plasmids. Viral vectors transfect cells directly; plasmid DNA can be delivered with the help of, for example, cationic liposomes (lipofectin) or derivatized (*e.g.*, antibody conjugated) polylysine conjugates, gramacidin S, artificial viral envelopes or other such intracellular earners, as well as direct injection of the gene construct or CaPO<sub>4</sub> precipitation carried out *in vivo*.

In an embodiment, *in vivo* introduction of nucleic acid into a cell includes use of a viral vector containing nucleic acid, *e.g.*, a cDNA. Infection of cells with a viral vector has the advantage that a large proportion of the targeted cells can receive the nucleic acid. Additionally, molecules encoded within the viral vector, *e.g.*, by a cDNA contained in the viral vector, are expressed efficiently in cells which have taken up viral vector nucleic acid.

Retroviral vectors and adeno-associated virus vectors can be used as a recombinant gene delivery system for the transfer of exogenous genes *in vivo* particularly into humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology, Ausubel, F. M. *et al.* (eds.) Greene Publishing Associates (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE, and pEM which are known to those skilled in the art. Examples of suitable packaging virus lines for preparing both ecotropic and amphotropic retroviral systems include Crip, Cre, 2, and Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells, *in vitro* and/or *in vivo* (see, for example, Eglitis *et al.* (1985) *Science* 230:1395-1398; Danos and Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:6460-6464; Wilson *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:6141-6145; Huber *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:8039-8043; Ferry *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:8377-8381; Chowdhury *et al.* (1991) *Science* 254:1802-1805; van Beusechem *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:7640-7644; Kay *et al.* (1992) *Human Gene Therapy* 3:641-647; Dai *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Hwu *et al.* (1993) *J. Immunol.* 150:4104-4115; U.S. Pat. Nos. 4,868,116 and 4,980,286; PCT Pub. Nos. WO 89/07136, WO 89/02468, WO 89/05345, and WO 92/07573).

Another viral gene delivery system utilizes adenovirus-derived vectors. See, for example, Berkner *et al.* (1988) *BioTechniques* 6:616; Rosenfeld *et al.* (1991) *Science* 252:431-434; and Rosenfeld *et al.* (1992) *Cell* 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 d1324 or other strains of adenovirus (*e.g.*, Ad2, Ad3, Ad7 etc.) are known to those skilled in the art.

Yet another viral vector system useful for delivery of the subject gene is the adeno-associated virus (AAV). See, for example, Flotte *et al.* (1992) *Am. J. Respir. Cell. Mol. Biol.* 7:349-356; Samulski *et al.* (1989) *J. Virol.* 63:3822-3828; and McLaughlin *et al.* (1989) *J. Virol.* 62:1963-1973.

Pharmaceutical compositions

The compositions delineated herein include the compounds delineated herein, as well as additional therapeutic agents if present, in amounts effective for achieving a modulation of disease or disease symptoms, including those described herein.

The term “pharmaceutically acceptable carrier or adjuvant” refers to a carrier or adjuvant that may be administered to a patient, together with a compound of this invention, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a therapeutic amount of the compound.

Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as d- $\alpha$ -tocopherol polyethyleneglycol 1000 succinate, surfactants used in pharmaceutical dosage forms such as Tweens or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Cyclodextrins such as  $\alpha$ -,  $\beta$ -, and  $\gamma$ -cyclodextrin, or chemically modified derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- $\beta$ -cyclodextrins, or other solubilized derivatives may also be advantageously used to enhance delivery of compounds of the formulae described herein.

The pharmaceutical compositions containing inhibitors of IDH, *e.g.*, IDH1, may be administered directly to the central nervous system, such as into the cerebrospinal fluid or into the brain. Delivery can be, for example, in a bolus or by continuous pump infusion. In certain embodiments, delivery is by intrathecal delivery or by intraventricular injection directly into the brain. A catheter and, optionally, a pump can be used for delivery. The inhibitors can be delivered in and released from an implantable device, *e.g.*, a device that is implanted in association with surgical

removal of tumor tissue. *E.g.*, for delivery to the brain, the delivery can be analogous to that with Gliadel, a biopolymer wafer designed to deliver carmustine directly into the surgical cavity created when a brain tumor is resected. The Gliadel wafer slowly dissolves and delivers carmustine.

The therapeutics disclosed herein, *e.g.*, nucleic acid based inhibitors, *e.g.*, siRNAs can be administered directly to the CNS, *e.g.*, the brain, *e.g.*, using a pump and/or catheter system. In one embodiment, the pump is implanted under the skin. In an embodiment and a catheter attached to a pump is inserted into the CNS, *e.g.*, into the brain or spine. In one embodiment, the pump (such as the IsoMed Drug Pump from Medtronic) delivers dosing, *e.g.*, constant dosing, of a nucleic acid based inhibitor. In an embodiment, the pump is programmable to administer variable or constant doses at predetermined time intervals. For example, the IsoMed Drug pump from Medtronic (or a similar device) can be used to administer a constant supply of the inhibitor, or the SynchroMedII Drug Pump (or a similar device) can be used to administer a variable dosing regime.

Methods and devices described in US patents 7,044,932, 6,620,151, 6,283,949, and 6,685,452 can be used in methods described herein.

The pharmaceutical compositions of this invention may be administered orally, parenterally, by inhalation, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir, preferably by oral administration or administration by injection. The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability of the formulated compound or its delivery form. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example, as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents

that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms such as emulsions and or suspensions. Other commonly used surfactants such as Tweens or Spans and/or other similar emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, emulsions and aqueous suspensions, dispersions and solutions. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions and/or emulsions are administered orally, the active ingredient may be suspended or dissolved in an oily phase is combined with emulsifying and/or suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

The pharmaceutical compositions of this invention may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a compound of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active components. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

Topical administration of the pharmaceutical compositions of this invention is useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition should be formulated with a suitable ointment containing the active

components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active compound suspended or dissolved in a carrier with suitable emulsifying agents. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-transdermal patches are also included in this invention.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

When the compositions of this invention comprise a combination of a compound of the formulae described herein and one or more additional therapeutic or prophylactic agents, both the compound and the additional agent should be present at dosage levels of between about 1 to 100%, and more preferably between about 5 to 95% of the dosage normally administered in a monotherapy regimen. The additional agents may be administered separately, as part of a multiple dose regimen, from the compounds of this invention. Alternatively, those agents may be part of a single dosage form, mixed together with the compounds of this invention in a single composition.

The compounds described herein can, for example, be administered by injection, intravenously, intraarterially, subdermally, intraperitoneally, intramuscularly, or subcutaneously; or orally, buccally, nasally, transmucosally, topically, in an ophthalmic preparation, or by inhalation, with a dosage ranging from about 0.02 to about 100 mg/kg of body weight, alternatively dosages between 1 mg and 1000 mg/dose, every 4 to 120 hours, or according to the requirements of the particular drug. The methods herein contemplate administration of an effective amount of compound or compound composition to achieve the desired or stated

effect. Typically, the pharmaceutical compositions of this invention will be administered from about 1 to about 6 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5% to about 95% active compound (w/w). Alternatively, such preparations contain from about 20% to about 80% active compound.

Lower or higher doses than those recited above may be required. Specific dosage and treatment regimens for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the disease, condition or symptoms, the patient's disposition to the disease, condition or symptoms, and the judgment of the treating physician.

Upon improvement of a patient's condition, a maintenance dose of a compound, composition or combination of this invention may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained when the symptoms have been alleviated to the desired level. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

#### Kits

A compound described herein can be provided in a kit.

In an embodiment the kit includes (a) a compound described herein, *e.g.*, a composition that includes a compound described herein (wherein, *e.g.*, the compound can be an inhibitor described herein), and, optionally (b) informational material. The informational material can be descriptive, instructional, marketing or other material that relates to the methods described herein and/or the use of a compound described herein for the methods described herein.

In an embodiment the kit provides materials for evaluating a subject. The evaluation can be, *e.g.*, for: identifying a subject having unwanted levels (*e.g.*, higher than present in normal or wildtype cells) of any of 2HG, 2HG neoactivity, or mutant

IDH1 or IDH2 protien having 2HG neoactivity (or corresponding RNA), or having a somatic mutation in IDH1 or IDH2 characterized by 2HG neoactivity; diagnosing, prognosing, or staging, a subject, *e.g.*, on the basis of having increased levels of 2HG, 2HG neoactivity, or mutant IDH1 or IDH2 protien having 2HG neoactivity (or corresponding RNA), or having a somatic mutation in IDH1 or IDH2 characterized by 2HG neoactivity; selecting a treatment for, or evaluating the efficacy of, a treatment, *e.g.*, on the basis of the subject having increased levels of 2HG, 2HG neoactivity, or mutant IDH1 or IDH2 protien having 2HG neoactivity (or corresponding RNA), or having a somatic mutation in IDH1 or IDH2 characterized by 2HG neoactivity. The kit can include one or more reagent useful in the evaluation, *e.g.*, reagents mentioned elsewhere herein. A detection reagent, *e.g.*, an antibody or other specific binding reagent can be included. Standards or reference samples, *e.g.*, a positive or negative control standard can be included. *E.g.*, if the evaluation is based on the presence of 2HG the kit can include a reagent, *e.g.*, a positive or negative control standards for an assay, *e.g.*, a LC-MS assay.

If the evaluation is based on the presence of 2HG neoactivity, the kit can include a reagent, *e.g.*, one or more of those mentioned elsewhere herein, for assaying 2HG neoactivity. If the evaluation is based on sequencing, the kit can include primers or other matierials useful for sequencing the relevant nucleic acids for identifying an IHD, *e.g.*, IDH1 or IDH2, neoactive mutant. *E.g.*, the kit can contain a reagent that provides for interrogation of the indentity, *i.e.*, sequencing of, residue 132 of IDH1 to determine if a neoactive mutant is present. The kit can include nucleic acids, *e.g.*, an oligomer, *e.g.*, primers, which allow sequencing of of the nucleotides that encode residue 132 of IDH1. In an embodiment the kit includes a nucleic acid whose hybridization, or ability to be amplified, is dependent on the indentity of residue 132 of IDH1. In other embodiments the kit includes a reagent, *e.g.*, an antibody or other specific binding molecule that can identify the presence of a neoactive mutant, *e.g.*, a protein encoded by a neoactive mutant at 132 of IDH1. As described below, a kit can also include buffers, solvents, and information related to the evaluation.

In one embodiment, the informational material can include information about production of the compound, molecular weight of the compound, concentration, date of expiration, batch or production site information, and so forth. In one embodiment, the informational material relates to methods for administering the compound.



In one embodiment, the informational material can include instructions to administer a compound described herein in a suitable manner to perform the methods described herein, *e.g.*, in a suitable dose, dosage form, or mode of administration (*e.g.*, a dose, dosage form, or mode of administration described herein). In another embodiment, the informational material can include instructions to administer a compound described herein to a suitable subject, *e.g.*, a human, *e.g.*, a human having or at risk for a disorder described herein.

The informational material of the kits is not limited in its form. In many cases, the informational material, *e.g.*, instructions, is provided in printed matter, *e.g.*, a printed text, drawing, and/or photograph, *e.g.*, a label or printed sheet. However, the informational material can also be provided in other formats, such as Braille, computer readable material, video recording, or audio recording. In another embodiment, the informational material of the kit is contact information, *e.g.*, a physical address, email address, website, or telephone number, where a user of the kit can obtain substantive information about a compound described herein and/or its use in the methods described herein. Of course, the informational material can also be provided in any combination of formats.

In addition to a compound described herein, the composition of the kit can include other ingredients, such as a solvent or buffer, a stabilizer, a preservative, a flavoring agent (*e.g.*, a bitter antagonist or a sweetener), a fragrance or other cosmetic ingredient, and/or a second agent for treating a condition or disorder described herein. Alternatively, the other ingredients can be included in the kit, but in different compositions or containers than a compound described herein. In such embodiments, the kit can include instructions for admixing a compound described herein and the other ingredients, or for using a compound described herein together with the other ingredients.

A compound described herein can be provided in any form, *e.g.*, liquid, dried or lyophilized form. It is preferred that a compound described herein be substantially pure and/or sterile. When a compound described herein is provided in a liquid solution, the liquid solution preferably is an aqueous solution, with a sterile aqueous solution being preferred. When a compound described herein is provided as a dried form, reconstitution generally is by the addition of a suitable solvent. The solvent, *e.g.*, sterile water or buffer, can optionally be provided in the kit.

The kit can include one or more containers for the composition containing a compound described herein. In some embodiments, the kit contains separate containers, dividers or compartments for the composition and informational material. For example, the composition can be contained in a bottle, vial, or syringe, and the informational material can be contained in a plastic sleeve or packet. In other embodiments, the separate elements of the kit are contained within a single, undivided container. For example, the composition is contained in a bottle, vial or syringe that has attached thereto the informational material in the form of a label. In some embodiments, the kit includes a plurality (*e.g.*, a pack) of individual containers, each containing one or more unit dosage forms (*e.g.*, a dosage form described herein) of a compound described herein. For example, the kit includes a plurality of syringes, ampules, foil packets, or blister packs, each containing a single unit dose of a compound described herein. The containers of the kits can be air tight, waterproof (*e.g.*, impermeable to changes in moisture or evaporation), and/or light-tight.

The kit optionally includes a device suitable for administration of the composition, *e.g.*, a syringe, inhalant, pipette, forceps, measured spoon, dropper (*e.g.*, eye dropper), swab (*e.g.*, a cotton swab or wooden swab), or any such delivery device. In an embodiment, the device is a medical implant device, *e.g.*, packaged for surgical insertion.

#### Combination therapies

In some embodiments, a compound or composition described herein, is administered together with an additional cancer treatment. Exemplary cancer treatments include, for example: surgery, chemotherapy, targeted therapies such as antibody therapies, immunotherapy, and hormonal therapy. Examples of each of these treatments are provided below.

##### *Chemotherapy*

In some embodiments, a compound or composition described herein, is administered with a chemotherapy. Chemotherapy is the treatment of cancer with drugs that can destroy cancer cells. “Chemotherapy” usually refers to cytotoxic drugs which affect rapidly dividing cells in general, in contrast with targeted therapy. Chemotherapy drugs interfere with cell division in various possible ways, *e.g.*, with the duplication of DNA or the separation of newly formed chromosomes. Most forms of chemotherapy *target all* rapidly dividing cells and are not specific for cancer cells,

although some degree of specificity may come from the inability of many cancer cells to repair DNA damage, while normal cells generally can.

Examples of chemotherapeutic agents used in cancer therapy include, for example, antimetabolites (*e.g.*, folic acid, purine, and pyrimidine derivatives) and alkylating agents (*e.g.*, nitrogen mustards, nitrosoureas, platinum, alkyl sulfonates, hydrazines, triazines, aziridines, spindle poison, cytotoxic agents, topoisomerase inhibitors and others). Exemplary agents include Aclarubicin, Actinomycin, Alitretinon, Altretamine, Aminopterin, Aminolevulinic acid, Amrubicin, Amsacrine, Anagrelide, Arsenic trioxide, Asparaginase, Atrasentan, Belotecan, Bexarotene, endamustine, Bleomycin, Bortezomib, Busulfan, Camptothecin, Capecitabine, Carboplatin, Carboquone, Carmofur, Carmustine, Celecoxib, Chlorambucil, Chlormethine, Cisplatin, Cladribine, Clofarabine, Crisantaspase, Cyclophosphamide, Cytarabine, Dacarbazine, Dactinomycin, Daunorubicin, Decitabine, Demecolcine, Docetaxel, Doxorubicin, Efaproxiral, Elesclomol, Elsamitrucin, Enocitabine, Epirubicin, Estramustine, Etoposide, Floxuridine, Fludarabine, Fluorouracil (5FU), Fotemustine, Gemcitabine, Gliadel implants, Hydroxycarbamide, Hydroxyurea, Idarubicin, Ifosfamide, Irinotecan, Irofulven, Ixabepilone, Larotaxel, Leucovorin, Liposomal doxorubicin, Liposomal daunorubicin, Lonidamine, Lomustine, Lucanthone, Mannosulfan, Masoprocol, Melphalan, Mercaptopurine, Mesna, Methotrexate, Methyl aminolevulinate, Mitobronitol, Mitoguazone, Mitotane, Mitomycin, Mitoxantrone, Nedaplatin, Nimustine, Oblimersen, Omacetaxine, Ortaxel, Oxaliplatin, Paclitaxel, Pegaspargase, Pemetrexed, Pentostatin, Pirarubicin, Pixantrone, Plicamycin, Porfimer sodium, Prednimustine, Procarbazine, Raltitrexed, Ranimustine, Rubitecan, Sapacitabine, Semustine, Sitimagene ceradenovec, Strataplatin, Streptozocin, Talaporfin, Tegafur-uracil, Temoporfin, Temozolomide, Teniposide, Tesetaxel, Testolactone, Tetranitrate, Thiotepa, Tiazofurine, Tioguanine, Tipifarnib, Topotecan, Trabectedin, Triaziquone, Triethylenemelamine, Triplatin, Tretinoin, Treosulfan, Trofosfamide, Uramustine, Valrubicin, Verteporfin, Vinblastine, Vincristine, Vindesine, Vinflunine, Vinorelbine, Vorinostat, Zorubicin, and other cytostatic or cytotoxic agents described herein.

Because some drugs work better together than alone, two or more drugs are often given at the same time. Often, two or more chemotherapy agents are used as combination chemotherapy. In some embodiments, the chemotherapy agents

(including combination chemotherapy) can be used in combination with a compound described herein, *e.g.*, phenformin.

*Targeted therapy*

In some embodiments, a compound or composition described herein, is administered with a targeted therapy. Targeted therapy constitutes the use of agents specific for the deregulated proteins of cancer cells. Small molecule targeted therapy drugs are generally inhibitors of enzymatic domains on mutated, overexpressed, or otherwise critical proteins within the cancer cell. Prominent examples are the tyrosine kinase inhibitors such as Axitinib, Bosutinib, Cediranib, dasatinib, erlotinib, imatinib, gefitinib, lapatinib, Lestaurtinib, Nilotinib, Semaxanib, Sorafenib, Sunitinib, and Vandetanib, and also cyclin-dependent kinase inhibitors such as Alvocidib and Seliciclib. Monoclonal antibody therapy is another strategy in which the therapeutic agent is an antibody which specifically binds to a protein on the surface of the cancer cells. Examples include the anti-HER2/neu antibody trastuzumab (HERCEPTIN®) typically used in breast cancer, and the anti-CD20 antibody rituximab and Tositumomab typically used in a variety of B-cell malignancies. Other exemplary antibodies include Cetuximab, Panitumumab, Trastuzumab, Alemtuzumab, Bevacizumab, Edrecolomab, and Gemtuzumab. Exemplary fusion proteins include Aflibercept and Denileukin diftitox. In some embodiments, the targeted therapy can be used in combination with a compound described herein, *e.g.*, a biguanide such as metformin or phenformin, preferably phenformin.

Targeted therapy can also involve small peptides as “homing devices” which can bind to cell surface receptors or affected extracellular matrix surrounding the tumor. Radionuclides which are attached to these peptides (*e.g.*, RGDs) eventually kill the cancer cell if the nuclide decays in the vicinity of the cell. An example of such therapy includes BEXXAR®.

*Immunotherapy*

In some embodiments, a compound or composition described herein, is administered with an immunotherapy. Cancer immunotherapy refers to a diverse set of therapeutic strategies designed to induce the patient's own immune system to fight the tumor. Contemporary methods for generating an immune response against tumors include intravesicular BCG immunotherapy for superficial bladder cancer, and use of interferons and other cytokines to induce an immune response in renal cell carcinoma and melanoma patients.

Allogeneic hematopoietic stem cell transplantation can be considered a form of immunotherapy, since the donor's immune cells will often attack the tumor in a graft-versus-tumor effect. In some embodiments, the immunotherapy agents can be used in combination with a compound or composition described herein.

*Hormonal therapy*

In some embodiments, a compound or composition described herein, is administered with a hormonal therapy. The growth of some cancers can be inhibited by providing or blocking certain hormones. Common examples of hormone-sensitive tumors include certain types of breast and prostate cancers. Removing or blocking estrogen or testosterone is often an important additional treatment. In certain cancers, administration of hormone agonists, such as progestogens may be therapeutically beneficial. In some embodiments, the hormonal therapy agents can be used in combination with a compound or a composition described herein.

In some embodiments, a compound or composition described herein, is administered together with an additional cancer treatment (*e.g.*, surgical removal), in treating cancer in nervous system, *e.g.*, cancer in central nervous system, *e.g.*, brain tumor, *e.g.*, glioma, *e.g.*, glioblastoma multiforme (GBM).

Several studies have suggested that more than 25% of glioblastoma patients obtain a significant survival benefit from adjuvant chemotherapy. Meta-analyses have suggested that adjuvant chemotherapy results in a 6-10% increase in 1-year survival rate.

Temozolomide is an orally active alkylating agent that is used for persons newly diagnosed with glioblastoma multiforme. It was approved by the United States Food and Drug Administration (FDA) in March 2005. Studies have shown that the drug was well tolerated and provided a survival benefit. Adjuvant and concomitant temozolomide with radiation was associated with significant improvements in median progression-free survival over radiation alone (6.9 vs 5 mo), overall survival (14.6 vs 12.1 mo), and the likelihood of being alive in 2 years (26% vs 10%).

Nitrosoureas: BCNU (carmustine)-polymer wafers (Gliadel) were approved by the FDA in 2002. Though Gliadel wafers are used by some for initial treatment, they have shown only a modest increase in median survival over placebo (13.8 vs. 11.6 months) in the largest such phase III trial, and are associated with increased rates of CSF leak and increased intracranial pressure secondary to edema and mass effect.

MGMT is a DNA repair enzyme that contributes to temozolomide resistance. Methylation of the MGMT promoter, found in approximately 45% of glioblastoma multiformes, results in an epigenetic silencing of the gene, decreasing the tumor cell's capacity for DNA repair and increasing susceptibility to temozolomide.

When patients with and without MGMT promoter methylation were treated with temozolomide, the groups had median survivals of 21.7 versus 12.7 months, and 2-year survival rates of 46% versus 13.8%, respectively.

Though temozolomide is currently a first-line agent in the treatment of glioblastoma multiforme, unfavorable MGMT methylation status could help select patients appropriate for future therapeutic investigations.

O6-benzylguanine and other inhibitors of MGMT as well as RNA interference-mediated silencing of MGMT offer promising avenues to increase the effectiveness of temozolomide and other alkylating antineoplastics, and such agents are under active study.

Carmustine (BCNU) and cis -platinum (cisplatin) have been the primary chemotherapeutic agents used against malignant gliomas. All agents in use have no greater than a 30-40% response rate, and most fall into the range of 10-20%.

Data from the University of California at San Francisco indicate that, for the treatment of glioblastomas, surgery followed by radiation therapy leads to 1-, 3-, and 5-year survival rates of 44%, 6%, and 0%, respectively. By comparison, surgery followed by radiation and chemotherapy using nitrosourea-based regimens resulted in 1-, 3-, and 5-year survival rates of 46%, 18%, and 18%, respectively.

A major hindrance to the use of chemotherapeutic agents for brain tumors is the fact that the blood-brain barrier (BBB) effectively excludes many agents from the CNS. For this reason, novel methods of intracranial drug delivery are being developed to deliver higher concentrations of chemotherapeutic agents to the tumor cells while avoiding the adverse systemic effects of these medications.

Pressure-driven infusion of chemotherapeutic agents through an intracranial catheter, also known as convection-enhanced delivery (CED), has the advantage of delivering drugs along a pressure gradient rather than by simple diffusion. CED has shown promising results in animal models with agents including BCNU and topotecan.

Initial attempts investigated the delivery of chemotherapeutic agents via an intraarterial route rather than intravenously. Unfortunately, no survival advantage was observed.

Chemotherapy for recurrent glioblastoma multiforme provides modest, if any, benefit, and several classes of agents are used. Carmustine wafers increased 6-month survival from 36% to 56% over placebo in one randomized study of 222 patients, though there was a significant association between the treatment group and serious intracranial infections.

Genotyping of brain tumors may have applications in stratifying patients for clinical trials of various novel therapies.

The anti-angiogenic agent bevacizumab, when used with irinotecan improved 6-month survival in recurrent glioma patients to 46% compared with 21% in patients treated with temozolomide. This bevacizumab and irinotecan combination for recurrent glioblastoma multiforme has been shown to improve survival over bevacizumab alone. Anti-angiogenic agents also decrease peritumoral edema, potentially reducing the necessary corticosteroid dose.

Some glioblastomas responds to gefitinib or erlotinib (tyrosine kinase inhibitors). The simultaneous presence in glioblastoma cells of mutant EGFR (EGFRviii) and PTEN was associated with responsiveness to tyrosine kinase inhibitors, whereas increased p-akt predicts a decreased effect. Other targets include PDGFR, VEGFR, mTOR, farnesyltransferase, and PI3K.

Other possible therapy modalities include imatinib, gene therapy, peptide and dendritic cell vaccines, synthetic chlorotoxins, and radiolabeled drugs and antibodies.

#### Patient selection/monitoring

Described herein are methods of treating a cell proliferation-related disorder, *e.g.*, cancer, in a subject and methods of identifying a subject for a treatment described herein. Also described herein are methods of predicting a subject who is at risk of developing cancer (*e.g.*, a cancer associate with a mutation in an enzyme (*e.g.*, an enzyme in the metabolic pathway such as IDH1 and/or IDH2)). The cancer is generally characterized by the presence of a neoactivity, such as a gain of function in one or more mutant enzymes (*e.g.*, an enzyme in the metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway, *e.g.*, IDH1 or

IDH2). The subject can be selected on the basis of the subject having a mutant gene having a neoactivity, *e.g.*, a neoactivity described herein. As used herein, “select” means selecting in whole or part on said basis.

In some embodiments, a subject is selected for treatment with a compound described herein based on a determination that the subject has a mutant enzyme described herein (*e.g.*, an enzyme in the metabolic pathway, *e.g.*, a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway, *e.g.*, IDH1 or IDH2). In some embodiments, the mutant enzyme has a neoactivity and the patient is selected on that basis. The neoactivity of the enzyme can be identified, for example, by evaluating the subject or sample (*e.g.*, tissue or bodily fluid) therefrom, for the presence or amount of a substrate, cofactor and/or product of the enzyme. The presence and/or amount of substrate, cofactor and/or product can correspond to the wild-type/non-mutant activity or can correspond to the neoactivity of the enzyme. Exemplary bodily fluid that can be used to identify (*e.g.*, evaluate) the neoactivity of the enzyme include amniotic fluid surrounding a fetus, aqueous humour, blood (*e.g.*, blood plasma), Cerebrospinal fluid, cerumen, chyme, Cowper's fluid, female ejaculate, interstitial fluid, lymph, breast milk, mucus (*e.g.*, nasal drainage or phlegm), pleural fluid, pus, saliva, sebum, semen, serum, sweat, tears, urine, vaginal secretion, or vomit.

In some embodiments, a subject can be evaluated for neoactivity of an enzyme using magnetic resonance. For example, where the mutant enzyme is IDH1 or IDH2 and the neoactivity is conversion of  $\alpha$ -ketoglutarate to 2-hydroxyglutarate, the subject can be evaluated for the presence of and/or an elevated amount of 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate relative to the amount of 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate present in a subject who does not have a mutation in IDH1 or IDH2 having the above neoactivity. In some embodiments, neoactivity of IDH1 or IDH2 can be determined by the presence or elevated amount of a peak corresponding to 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate as determined by magnetic resonance. For example, a subject can be evaluated for the presence and/or strength of a signal at about 2.5 ppm to determine the presence and/or amount of 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate in the subject. This can be correlated to and/or predictive of a neoactivity described herein for the mutant enzyme IDH. Similarly, the presence,



strength and/or absence of a signal at about 2.5 ppm could be predictive of a response to treatment and thereby used as a noninvasive biomarker for clinical response.

Neoactivity of a mutant enzyme such as IDH can also be evaluated using other techniques known to one skilled in the art. For example, the presence or amount of a labeled substrate, cofactor, and/or reaction product can be measured such as a <sup>13</sup>C or <sup>14</sup>C labeled substrate, cofactor, and/or reaction product. The neoactivity can be evaluated by evaluating the forward reaction of the wild-type/non mutant enzyme (such as the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate in a mutant IDH1 or IDH2 enzyme, specifically a mutant IDH1 enzyme) and/or the reaction corresponding to the neoactivity (*e.g.*, the conversion of  $\alpha$ -ketoglutarate to 2-hydroxyglutarate, *e.g.*, R-2-hydroxyglutarate in a mutant IDH1 or IDH2 enzyme, specifically a mutant IDH1 enzyme).

#### Disorders

The IDH-related methods disclosed herein, *e.g.*, methods of evaluating or treating subjects, are directed to subjects having a cell proliferation-related disorder characterized by an IDH mutant, *e.g.*, an IDH1 or IDH2, mutant having neoactivity, *e.g.*, 2HG neoactivity. Examples of some of the disorders below have been shown to be characterized by an IDH1 or IDH2 mutation. Others can be analyzed, *e.g.*, by sequencing cell samples to determine the presence of a somatic mutation at amino acid 132 of IDH1 or at amino acid 172 of IDH2. Without being bound by theory it is expected that a portion of the tumors of given type of cancer will have an IDH, *e.g.*, IDH1 or IDH2, mutant having 2HG neoactivity.

The disclosed methods are useful in evaluating or treating proliferative disorders, *e.g.* evaluating or treating solid tumors, soft tissue tumors, and metastases thereof wherein the solid tumor, soft tissue tumor or metastases thereof is a cancer described herein. Exemplary solid tumors include malignancies (*e.g.*, sarcomas, adenocarcinomas, and carcinomas) of the various organ systems, such as those of brain, lung, breast, lymphoid, gastrointestinal (*e.g.*, colon), and genitourinary (*e.g.*, renal, urothelial, or testicular tumors) tracts, pharynx, prostate, and ovary. Exemplary adenocarcinomas include colorectal cancers, renal-cell carcinoma, liver cancer, non-small cell carcinoma of the lung, and cancer of the small intestine. The disclosed methods are also useful in evaluating or treating non-solid cancers.

The methods described herein can be used with any cancer, for example those described by the National Cancer Institute. A cancer can be evaluated to determine whether it is using a method described herein. Exemplary cancers described by the National Cancer Institute include: Acute Lymphoblastic Leukemia, Adult; Acute Lymphoblastic Leukemia, Childhood; Acute Myeloid Leukemia, Adult; Adrenocortical Carcinoma; Adrenocortical Carcinoma, Childhood; AIDS-Related Lymphoma; AIDS-Related Malignancies; Anal Cancer; Astrocytoma, Childhood Cerebellar; Astrocytoma, Childhood Cerebral; Bile Duct Cancer, Extrahepatic; Bladder Cancer; Bladder Cancer, Childhood; Bone Cancer, Osteosarcoma/Malignant Fibrous Histiocytoma; Brain Stem Glioma, Childhood; Brain Tumor, Adult; Brain Tumor, Brain Stem Glioma, Childhood; Brain Tumor, Cerebellar Astrocytoma, Childhood; Brain Tumor, Cerebral Astrocytoma/Malignant Glioma, Childhood; Brain Tumor, Ependymoma, Childhood; Brain Tumor, Medulloblastoma, Childhood; Brain Tumor, Supratentorial Primitive Neuroectodermal Tumors, Childhood; Brain Tumor, Visual Pathway and Hypothalamic Glioma, Childhood; Brain Tumor, Childhood (Other); Breast Cancer; Breast Cancer and Pregnancy; Breast Cancer, Childhood; Breast Cancer, Male; Bronchial Adenomas/Carcinoids, Childhood; Carcinoid Tumor, Childhood; Carcinoid Tumor, Gastrointestinal; Carcinoma, Adrenocortical; Carcinoma, Islet Cell; Carcinoma of Unknown Primary; Central Nervous System Lymphoma, Primary; Cerebellar Astrocytoma, Childhood; Cerebral Astrocytoma/Malignant Glioma, Childhood; Cervical Cancer; Childhood Cancers; Chronic Lymphocytic Leukemia; Chronic Myelogenous Leukemia; Chronic Myeloproliferative Disorders; Clear Cell Sarcoma of Tendon Sheaths; Colon Cancer; Colorectal Cancer, Childhood; Cutaneous T-Cell Lymphoma; Endometrial Cancer; Ependymoma, Childhood; Epithelial Cancer, Ovarian; Esophageal Cancer; Esophageal Cancer, Childhood; Ewing's Family of Tumors; Extracranial Germ Cell Tumor, Childhood; Extragonadal Germ Cell Tumor; Extrahepatic Bile Duct Cancer; Eye Cancer, Intraocular Melanoma; Eye Cancer, Retinoblastoma; Gallbladder Cancer; Gastric (Stomach) Cancer; Gastric (Stomach) Cancer, Childhood; Gastrointestinal Carcinoid Tumor; Germ Cell Tumor, Extracranial, Childhood; Germ Cell Tumor, Extragonadal; Germ Cell Tumor, Ovarian; Gestational Trophoblastic Tumor; Glioma, Childhood Brain Stem; Glioma, Childhood Visual Pathway and Hypothalamic; Hairy Cell Leukemia; Head and Neck Cancer; Hepatocellular (Liver) Cancer, Adult (Primary); Hepatocellular (Liver) Cancer, Childhood (Primary); Hodgkin's

Lymphoma, Adult; Hodgkin's Lymphoma, Childhood; Hodgkin's Lymphoma During Pregnancy; Hypopharyngeal Cancer; Hypothalamic and Visual Pathway Glioma, Childhood; Intraocular Melanoma; Islet Cell Carcinoma (Endocrine Pancreas); Kaposi's Sarcoma; Kidney Cancer; Laryngeal Cancer; Laryngeal Cancer, Childhood; Leukemia, Acute Lymphoblastic, Adult; Leukemia, Acute Lymphoblastic, Childhood; Leukemia, Acute Myeloid, Adult; Leukemia, Acute Myeloid, Childhood; Leukemia, Chronic Lymphocytic; Leukemia, Chronic Myelogenous; Leukemia, Hairy Cell; Lip and Oral Cavity Cancer; Liver Cancer, Adult (Primary); Liver Cancer, Childhood (Primary); Lung Cancer, Non-Small Cell; Lung Cancer, Small Cell; Lymphoblastic Leukemia, Adult Acute; Lymphoblastic Leukemia, Childhood Acute; Lymphocytic Leukemia, Chronic; Lymphoma, AIDS- Related; Lymphoma, Central Nervous System (Primary); Lymphoma, Cutaneous T-Cell; Lymphoma, Hodgkin's, Adult; Lymphoma, Hodgkin's, Childhood; Lymphoma, Hodgkin's During Pregnancy; Lymphoma, Non-Hodgkin's, Adult; Lymphoma, Non- Hodgkin's, Childhood; Lymphoma, Non-Hodgkin's During Pregnancy; Lymphoma, Primary Central Nervous System; Macroglobulinemia, Waldenstrom's; Male Breast Cancer; Malignant Mesothelioma, Adult; Malignant Mesothelioma, Childhood; Malignant Thymoma; Medulloblastoma, Childhood; Melanoma; Melanoma, Intraocular; Merkel Cell Carcinoma; Mesothelioma, Malignant; Metastatic Squamous Neck Cancer with Occult Primary; Multiple Endocrine Neoplasia Syndrome, Childhood; Multiple Myeloma/Plasma Cell Neoplasm; Mycosis Fungoides; Myelodysplastic Syndromes; Myelogenous Leukemia, Chronic; Myeloid Leukemia, Childhood Acute; Myeloma, Multiple; Myeloproliferative Disorders, Chronic; Nasal Cavity and Paranasal Sinus Cancer; Nasopharyngeal Cancer; Nasopharyngeal Cancer, Childhood; Neuroblastoma; Non-Hodgkin's Lymphoma, Adult; Non-Hodgkin's Lymphoma, Childhood; Non-Hodgkin's Lymphoma During Pregnancy; Non-Small Cell Lung Cancer; Oral Cancer, Childhood; Oral Cavity and Lip Cancer; Oropharyngeal Cancer; Osteosarcoma/Malignant Fibrous Histiocytoma of Bone; Ovarian Cancer, Childhood; Ovarian Epithelial Cancer; Ovarian Germ Cell Tumor; Ovarian Low Malignant Potential Tumor; Pancreatic Cancer; Pancreatic Cancer, Childhood; Pancreatic Cancer, Islet Cell; Paranasal Sinus and Nasal Cavity Cancer; Parathyroid Cancer; Penile Cancer; Pheochromocytoma; Pineal and Supratentorial Primitive Neuroectodermal Tumors, Childhood; Pituitary Tumor; Plasma Cell Neoplasm/Multiple Myeloma; Pleuropulmonary Blastoma; Pregnancy and Breast Cancer; Pregnancy and Hodgkin's

Lymphoma; Pregnancy and Non-Hodgkin's Lymphoma; Primary Central Nervous System Lymphoma; Primary Liver Cancer, Adult; Primary Liver Cancer, Childhood; Prostate Cancer; Rectal Cancer; Renal Cell (Kidney) Cancer; Renal Cell Cancer, Childhood; Renal Pelvis and Ureter, Transitional Cell Cancer; Retinoblastoma; Rhabdomyosarcoma, Childhood; Salivary Gland Cancer; Salivary Gland Cancer, Childhood; Sarcoma, Ewing's Family of Tumors; Sarcoma, Kaposi's; Sarcoma (Osteosarcoma)/Malignant Fibrous Histiocytoma of Bone; Sarcoma, Rhabdomyosarcoma, Childhood; Sarcoma, Soft Tissue, Adult; Sarcoma, Soft Tissue, Childhood; Sezary Syndrome; Skin Cancer; Skin Cancer, Childhood; Skin Cancer (Melanoma); Skin Carcinoma, Merkel Cell; Small Cell Lung Cancer; Small Intestine Cancer; Soft Tissue Sarcoma, Adult; Soft Tissue Sarcoma, Childhood; Squamous Neck Cancer with Occult Primary, Metastatic; Stomach (Gastric) Cancer; Stomach (Gastric) Cancer, Childhood; Supratentorial Primitive Neuroectodermal Tumors, Childhood; T- Cell Lymphoma, Cutaneous; Testicular Cancer; Thymoma, Childhood; Thymoma, Malignant; Thyroid Cancer; Thyroid Cancer, Childhood; Transitional Cell Cancer of the Renal Pelvis and Ureter; Trophoblastic Tumor, Gestational; Unknown Primary Site, Cancer of, Childhood; Unusual Cancers of Childhood; Ureter and Renal Pelvis, Transitional Cell Cancer; Urethral Cancer; Uterine Sarcoma; Vaginal Cancer; Visual Pathway and Hypothalamic Glioma, Childhood; Vulvar Cancer; Waldenstrom's Macro globulinemia; and Wilms' Tumor. Metastases of the aforementioned cancers can also be treated or prevented in accordance with the methods described herein.

The methods described herein are useful in treating cancer in nervous system, *e.g.*, brain tumor, *e.g.*, glioma, *e.g.*, glioblastoma multiforme (GBM), *e.g.*, by inhibiting a neoactivity of a mutant enzyme, *e.g.*, an enzyme in a metabolic pathway, *e.g.*, a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, the nucleotide biosynthetic pathway, or the fatty acid biosynthetic pathway, *e.g.*, IDH1 or IDH2.

Gliomas, a type of brain tumors, can be classified as grade I to grade IV on the basis of histopathological and clinical criteria established by the World Health Organization (WHO). WHO grade I gliomas are often considered benign. Gliomas of WHO grade II or III are invasive, progress to higher-grade lesions. WHO grade IV tumors (glioblastomas) are the most invasive form. Exemplary brain tumors include, *e.g.*, astrocytic tumor (*e.g.*, pilocytic astrocytoma, subependymal giant-cell

astrocytoma, diffuse astrocytoma, pleomorphic xanthoastrocytoma, anaplastic astrocytoma, astrocytoma, giant cell glioblastoma, glioblastoma, secondary glioblastoma, primary adult glioblastoma, and primary pediatric glioblastoma); oligodendroglial tumor (*e.g.*, oligodendroglioma, and anaplastic oligodendroglioma); oligoastrocytic tumor (*e.g.*, oligoastrocytoma, and anaplastic oligoastrocytoma); ependymoma (*e.g.*, myxopapillary ependymoma, and anaplastic ependymoma); medulloblastoma; primitive neuroectodermal tumor, schwannoma, meningioma, atypical meningioma, anaplastic meningioma; and pituitary adenoma. Exemplary cancers are described in *Acta Neuropathol* (2008) 116:597–602 and *N Engl J Med*. 2009 Feb 19;360(8):765-73, the contents of which are each incorporated herein by reference.

In embodiments the disorder is glioblastoma.

In an embodiment the disorder is prostate cancer, *e.g.*, stage T1 (*e.g.*, T1a, T1b and T1c), T2 (*e.g.*, T2a, T2b and T2c), T3 (*e.g.*, T3a and T3b) and T4, on the TNM staging system. In embodiments the prostate cancer is grade G1, G2, G3 or G4 (where a higher number indicates greater difference from normal tissue). Types of prostate cancer include, *e.g.*, prostate adenocarcinoma, small cell carcinoma, squamous carcinoma, sarcomas, and transitional cell carcinoma.

Methods and compositions of the invention can be combined with art-known treatment. Art-known treatment for prostate cancer can include, *e.g.*, active surveillance, surgery (*e.g.*, radical prostatectomy, transurethral resection of the prostate, orchiectomy, and cryosurgery), radiation therapy including brachytherapy (prostate brachytherapy) and external beam radiation therapy, High-Intensity Focused Ultrasound (HIFU), chemotherapy, cryosurgery, hormonal therapy (*e.g.*, antiandrogens (*e.g.*, flutamide, bicalutamide, nilutamide and cyproterone acetate, ketoconazole, aminoglutethimide), GnRH antagonists (*e.g.*, Abarelix)), or a combination thereof.

All references described herein are expressly incorporated herein by reference.

## EXAMPLES

### Example 1 IDH1 cloning, mutagenesis, expression and purification

#### ***1. Wild type IDH1 was cloned into pET41a, creating His8 tag at C-terminus.***

The IDH1 gene coding region (cDNA) was purchased from Invitrogen in pENTR221 vector ([www.invitrogen.com](http://www.invitrogen.com), Cat#B-068487\_Ultimate\_ORF). Oligo

nucleotides were designed to PCR out the coding region of IDH1 with NdeI at the 5' end and XhoI at the 3'. (IDH1-f: TAATCATATGTCCAAAAAATCAGT (SEQ ID NO:1), IDH1-r: TAATCTCGAGTGAAAGTTTGGCCTGAGCTAGTT (SEQ ID NO:2)). The PCR product is cloned into the NdeI/XhoI cleaved pET41a vector. NdeI/XhoI cleavage of the vector pET41a releases the GST portion of the plasmid, and creating a C-terminal His8 tag (SEQ ID NO:3) without the N-terminal GST fusion. The original stop codon of IDH1 is change to serine, so the junction sequence in final IDH1 protein is: Ser-Leu-Glu-His-His-His-His-His-His-His-Stop (SEQ ID NO:4).

The C-terminal His tag strategy instead of N-terminal His tag strategy was chosen, because C-terminal tag might not negatively impact IDH1 protein folding or activity. See, *e.g.*, Xu X *et al*, J Biol Chem. 2004 Aug 6; 279(32):33946-57.

The sequence for pET41a-IDH1 plasmid is confirmed by DNA sequencing. **FIG. 1** shows detailed sequence verification of pET41a-IDH1 and alignment against published IDH1 CDS below.

## ***2. IDH1 site directed mutagenesis to create the IDHr132s and IDHr132h mutants.***

Site directed mutagenesis was performed to convert R132 to S or H, DNA sequencing confirmed that G395 is mutated to A (creating Arg→His mutation in the IDH1 protein), and C394 is mutated to A (creating Arg→Ser in the IDH1 protein). Detailed method for site directed mutagenesis is described in the user manual for QuikChange® MultiSite-Directed Mutagenesis Kit (Stratagene, cat# 200531). **FIG. 2** shows DNA sequence verification of such mutations. Highlighted nucleotides were successfully changed in the mutagenesis: G395→A mutation allows amino acid Arg132→His; C394→A mutation allows amino acid Arg132→Ser.

## ***3. IDH1 protein expression and purification.***

IDHwt, IDHR132S, and IDHR132H proteins were expressed in the *E. coli* strain Rosetta and purified according to the detailed procedure below. Active IDH1 proteins are in dimer form, and SEC column fraction/peak that correspond to the dimer form were collected for enzymology analysis and cross comparison of catalytic activities of these proteins.

### A. Cell culturing:

Cells were grown in LB (20 µg/ml Kanamycin) at 37°C with shaking until OD600 reaches 0.6. The temperature was changed to 18°C and protein was induced by adding IPTG to final concentration of 1 mM. Cells were collected 12-16 hours after IPTG induction.

B. Buffer system:

Lysis buffer: 20mM Tris, pH7.4, 0.1% Triton X-100, 500 mM NaCl, 1 mM PMSF, 5 mM β-mercaptoethanol, 10 % glycerol.

Ni-Column Buffer A: 20 mM Tris, pH7.4, 500mM NaCl, 5 mM β-mercaptoethanol, 10% glycerol.

Ni-column Buffer B: 20 mM Tris, pH7.4, 500 mM NaCl, 5 mM β-mercaptoethanol , 500 mM Imidazole, 10% glycerol

Gel filtration Buffer C: 200 mM NaCl, 50 mM Tris 7.5, 5 mM β-mercaptoethanol, 2 mM MnSO<sub>4</sub>, 10% glycerol.

C. Protein purification procedure

1. Cell pellet were resuspended in the lysis buffer (1gram cell/5-10 ml buffer).
2. Cells were broken by passing the cell through Microfluidizer with at a pressure of 15,000 psi for 3 times.
3. Soluble protein was collected from supernatant after centrifugation at 20,000g (Beckman Avanti J-26XP) for 30 min at 4°C.
4. 5-10 ml of Ni-column was equilibrated by Buffer A until the A280 value reached baseline. The supernatant was loaded onto a 5-ml Ni-Sepharose column (2 ml/min). The column was washed by 10-20 CV of washing buffer (90 % buffer A+10 % buffer B) until A280 reach the baseline (2 ml/min).
5. The protein was eluted by liner gradient of 10-100% buffer B (20 CV) with the flow rate of 2 ml/min and the sample fractions were collected as 2 ml/tube.
6. The samples were analyzed on SDS-PAGE gel.
7. The samples were collected and dialyzed against 200x Gel filtration buffer for 2 times (1 hour and > 4 hours).
8. The samples were concentrated to 10 ml.
9. 200 ml of S-200 Gel-filtration column was equilibrated by buffer C until the A280 value reached baseline. The samples were loaded onto Gel filtration column (0.5 ml/min).

10. The column was washed by 10 CV of buffer C, collect fractions as 2-4 ml/tube.
11. The samples were analyzed on SDS-PAGE gel and protein concentration was determined.

#### D. Protein purification results

The results for purification of wild type IDH1 are shown in **FIGs. 3, 4, 5A** and **5B**.

The results for purification of mutant IDH1R132S are shown in **FIGs. 6, 7, 8A** and **8B**.

The results for purification of wild type IDH1R132H are shown in **FIGs. 9, 10, 11A** and **11B**.

#### EXAMPLE 2 ENZYMOLOGY ANALYSIS OF IDH1 WILD TYPE AND MUTANTS

##### *1. Analysis of IDH1 wild-type and mutants R132H and R132S in the oxidative decarboxylation of isocitrate to $\alpha$ -Ketoglutarate ( $\alpha$ -KG).*

###### A. Methods

To determine the catalytic efficiency of enzymes in the oxidative decarboxylation of isocitrate to  $\alpha$ -Ketoglutarate ( $\alpha$ -KG) direction, reactions were performed to determine  $V_{max}$  and  $K_m$  for isocitrate. In these reactions, the substrate was varied while the cofactor was held constant at 500  $\mu$ M. All reactions were performed in 150 mM NaCl, 20 mM Tris-Cl, pH 7.5, 10% glycerol, and 0.03% (w/v) BSA). Reaction progress was followed by spectroscopy at 340 nM monitoring the change in oxidation state of the cofactor. Sufficient enzyme was added to give a linear change in absorbance for 10 minutes.

###### B. ICDH1 R132H and ICDH1 R132S are impaired for conversion of isocitrate to $\alpha$ -KG.

Michaelis-Menten plots for the relationship of isocitrate concentration to reaction velocity are presented in **FIGs. 12A-12C**. Kinetic parameters are summarized in the **Table 1**. All data was fit to the Hill equation by least-squares regression analysis.

###### **Table 1**



Enzyme	Vmax ( $\mu\text{mol}/\text{min}/\text{mg}$ )	Km ( $\mu\text{M}$ )	Hill Constant	Vmax/Km	Relative Catalytic Efficiency
Wt	30.5	56.8	1.8	0.537	100%
R132H	0.605	171.7	0.6	0.0035	0.35%
R132S	95	>1e6	0.479	<9.5e7	<.001%

Both mutant enzymes display a reduced Hill coefficient and an increase in Km for isocitrate, suggesting a loss of co-operativity in substrate binding and/or reduced affinity for substrate. R132H enzyme also displays a reduced Vmax, suggestive of a lower kcat. R132S displays an increase in Vmax, suggesting an increase in kcat, although this comes at the expense of a 20,000 fold increase in Km so that the overall effect on catalytic efficiency is a great decrease as compared to the wild-type enzyme. The relative catalytic efficiency, described as Vmax/Km, is dramatically lower for the mutants as compared to wild-type. The *in vivo* effect of these mutations would be to decrease the flux conversion of isocitrate to  $\alpha$ -KG.

C. The ICDH1 R132H and R132S mutants display reduced product inhibition in the oxidative decarboxylation of isocitrate to  $\alpha$ -Ketoglutarate ( $\alpha$ -KG).

A well-known regulatory mechanism for control of metabolic enzymes is feedback inhibition, in which the product of the reaction acts as a negative regulator for the generating enzyme. To examine whether the R132S or R132H mutants maintain this regulatory mechanism, the Ki for  $\alpha$ -KG in the oxidative decarboxylation of isocitrate to  $\alpha$ -ketoglutarate was determined. Data is presented in **FIGs. 13A-13C** and summarized in **Table 2**. In all cases,  $\alpha$ -KG acts as a competitive inhibitor of the isocitrate substrate. However, R132H and R132S display a 20-fold and 13-fold increase in sensitivity to feedback inhibition as compared to the wild-type enzyme.

**Table 2**

Enzyme	Ki ( $\mu\text{M}$ )
Wt	612.2
R132H	28.6
R132S	45.3

D. The effect of  $\text{MnCl}_2$  in oxidative decarboxylation of isocitrate to  $\alpha$ -Ketoglutarate ( $\alpha$ -KG).

MnCl<sub>2</sub> can be substituted with MgCl<sub>2</sub> to examine if there is any difference in oxidative decarboxylation of isocitrate to  $\alpha$ -Ketoglutarate ( $\alpha$ -KG).

E. The effect of R132 mutations on the inhibitory effect of oxalomalate on IDH1

The purpose of this example is to examine the susceptibility of IDH1R132S and IDH1R132H in oxidative decarboxylation of isocitrate to  $\alpha$ -Ketoglutarate ( $\alpha$ -KG) to the known IDH1 inhibitor oxalomalate. Experiments were performed to examine if R132 mutations circumvent the inhibition by oxalomalate.

Final concentrations: Tris 7.5 20 mM, NaCl 150 mM, MnCl<sub>2</sub> 2 mM, Glycerol 10%, BSA 0.03%, NADP 0.5 mM, IDH1 wt 1.5 ug/ml, IDH1R132S 30 ug/ml, IDH1R132H 60 ug/ml, DL-isocitrate (5 – 650 uM). The results are summarized in **FIG. 17** and Table 3. The R132S mutation displays approximately a two-fold increase in susceptibility to inhibition by oxalomalate, while the R132H mutation is essentially unaffected. In all three cases, the same fully competitive mode of inhibition with regards to isocitrate was observed.

**Table 3**

Enzyme	Oxalomalate Ki (uM)
wt	955.4
R132S	510
R132H	950.8

F. Forward reactions (isocitrate to  $\alpha$ -KG) of mutant enzyme do not go to completion.

Forward reactions containing ICDH1 R132S or ICDH1 R132H were assembled and reaction progress monitored by an increase in the OD340 of the reduced NADPH cofactor. It was observed (**FIG. 23**), that these reactions proceed in the forward direction for a period of time and then reverse direction and oxidize the cofactor reduced in the early stages of the reaction, essentially to the starting concentration present at the initiation of the experiment. Addition of further isocitrate re-initiated the forward reaction for a period of time, but again did not induce the reaction to proceed to completion. Rather, the system returned to initial concentrations of NADPH. This experiment suggested that the mutant enzymes were performing a reverse reaction other than the conversion of  $\alpha$ -KG to isocitrate.

**2. Analysis of IDH1 wild-type and mutants R132H and R132S in the reduction of  $\alpha$ -Ketoglutarate ( $\alpha$ -KG).**

**A. Methods**

To determine the catalytic efficiency of enzymes in the reduction of  $\alpha$ -Ketoglutarate ( $\alpha$ -KG), reactions were performed to determine  $V_{max}$  and  $K_m$  for  $\alpha$ -KG. In these reactions, substrate was varied while the cofactor was held constant at 500  $\mu$ M. All reactions were performed in 50 mM potassium phosphate buffer, pH 6.5, 10% glycerol, 0.03% (w/v) BSA, 5 mM  $MgCl_2$ , and 40 mM sodium hydrocarbonate. Reaction progress was followed by spectroscopy at 340 nM monitoring the change in oxidation state of the cofactor. Sufficient enzyme was added to give a linear change in absorbance for 10 minutes.

**B. The R132H and R132S mutant enzymes, but not the wild-type enzyme, support the reduction of  $\alpha$ -KG.**

To test the ability of the mutant and wild-type enzymes to perform the reduction of  $\alpha$ -KG, 40  $\mu$ g/ml of enzyme was incubated under the conditions for the reduction of  $\alpha$ -Ketoglutarate ( $\alpha$ -KG) as described above. Results are presented in **FIG. 14**. The wild-type enzyme was unable to consume NADPH, while R132S and R132H reduced  $\alpha$ -KG and consumed NADPH.

**C. The reduction of  $\alpha$ -KG by the R132H and R132S mutants occurs *in vitro* at physiologically relevant concentrations of  $\alpha$ -KG.**

To determine the kinetic parameters of the reduction of  $\alpha$ -KG performed by the mutant enzymes, a substrate titration experiment was performed, as presented in **FIGs. 15A-15B**. R132H maintained the Hill-type substrate interaction as seen in the oxidative decarboxylation of isocitrate, but displayed positive substrate co-operative binding. R132S showed a conversion to Michaelis-Menten kinetics with the addition of uncompetitive substrate inhibition, as compared to wild-type enzyme in the oxidative decarboxylation of isocitrate. The enzymatic parameters of the mutant enzyme are presented in **Table 4**. Since the wild-type enzyme did not consume measurable NADPH in the experiment described above, a full kinetic workup was not performed.

**Table 4**

Enzyme	Vmax (umol/min/mg)	Km (mM)	Hill Constant	Ki (mM)	Vmax/Km
R132H	1.3	0.965	1.8		1.35
R132S	2.7	0.181	0.479	24.6	14.92

The relative catalytic efficiency of reduction of  $\alpha$ -KG is approximately ten-fold higher in the R132S mutant than in the R132H mutant. The biological consequence is that the rate of metabolic flux should be greater in cells expressing R132S as compared to R132H.

D. Analysis of IDH1 wild-type and mutants R132H and R132S in the reduction of alpha-ketoglutarate with NADH.

In order to evaluate the ability of the mutant enzymes to utilize NADH in the reduction of alpha-ketoglutarate, the following experiment was conducted. Final concentrations: NaHCO<sub>3</sub> 40mM, MgCl<sub>2</sub> 5mM, Glycerol 10%, K<sub>2</sub>HPO<sub>4</sub> 50mM, BSA 0.03%, NADH 0.5mM, IDH1wt 5ug/ml, R132S 30ug/ml, R132H 60ug/ml, alpha-Ketoglutarate 5mM.

The results are shown in **FIG. 16** and **Table 5**. The R132S mutant demonstrated the ability to utilize NADH while the wild type and R132H show no measurable consumption of NADH in the presence of alpha-ketoglutarate.

**Table 5: Consumption of NADH by R132S in the presence of alpha-ketoglutarate**

	R132S		Mean	SD
<b>Rate (<math>\Delta</math>A/sec)</b>	0.001117	0.001088	0.001103	2.05E-05
<b>Umol/min/mg</b>	0.718328	0.699678	<b>0.709003</b>	0.013187

**Summary**

To understand how R132 mutations alter the enzymatic properties of IDH1, wild-type and R132H mutant IDH1 proteins were produced and purified from *E. coli*. When NADP<sup>+</sup>-dependent oxidative decarboxylation of isocitrate was measured using purified wild-type or R132H mutant IDH1 protein, it was confirmed that R132H mutation impairs the ability of IDH1 to catalyze this reaction (Yan, H. et al. N Engl J Med 360, 765-73 (2009); Zhao, S. et al. Science 324, 261-5 (2009)), as evident by the loss in binding affinity for both isocitrate and MgCl<sub>2</sub> along with a 1000-fold decrease

in catalytic turnover (**FIGs. 30A** and **30C**). In contrast, when NADPH-dependent reduction of  $\alpha$ KG was assessed using either wild-type or R132H mutant IDH1 protein, only R132H mutant could catalyze this reaction at a measurable rate (**FIGs. 30** and **30C**). Part of this increased rate of  $\alpha$ KG reduction results from an increase in binding affinity for both the cofactor NADPH and substrate  $\alpha$ KG in the R132H mutant IDH1 (**FIG. 30C**). Taken together, these data demonstrate that while the R132H mutation leads to a loss of enzymatic function for oxidative decarboxylation of isocitrate, this mutation also results in a gain of enzyme function for the NADPH-dependent reduction of  $\alpha$ KG.

## 2: Analysis of mutant IDH1

### **The R132H mutant does not result in the conversion of $\alpha$ -KG to isocitrate.**

Using standard experimental methods, an API2000 mass spectrometer was configured for optimal detection of  $\alpha$ -KG and isocitrate (Table 6). MRM transitions were selected and tuned such that each analyte was monitored by a unique transition. Then, an enzymatic reaction containing 1 mM  $\alpha$ -KG, 1 mM NADPH, and ICDH1 R132H were assembled and run to completion as judged by the decrease to baseline of the optical absorbance at 340 nM. A control reaction was performed in parallel from which the enzyme was omitted. Reactions were quenched 1:1 with methanol, extracted, and subjected to analysis by LC-MS/MS.

**FIG. 18A** presents the control reaction indicating that  $\alpha$ KG was not consumed in the absence of enzyme, and no detectable isocitrate was present. **FIG. 18B** presents the reaction containing R132H enzyme, in which the  $\alpha$ -KG has been consumed, but no isocitrate was detected. **FIG. 18C** presents a second analysis of the reaction containing enzyme in which isocitrate has been spiked to a final concentration of 1 mM, demonstrating that had  $\alpha$ -KG been converted to isocitrate at any appreciable concentration greater than 0.01%, the configured analytical system would have been capable of detecting its presence in the reaction containing enzyme. The conclusion from this experiment is that while  $\alpha$ -KG was consumed by R132H, isocitrate was not produced. This experiment indicates that one neoactivity of the R132H mutant is the reduction of  $\alpha$ -KG to a compound other than isocitrate.

Compound	Q1	Q3	DP	FP	EP	CEP	CE	CXP
$\alpha$ -KG	144.975	100.6	-6	-220	-10	-16	-10	-22
isocitrate	191.235	110.9	-11	-230	-4.5	-14	-16	-24
a-hydroxyglutarate	147.085	128.7	-11	-280	-10	-22	-12	-24

**The R132H mutant reduces  $\alpha$ -KG to 2-hydroxyglutaric acid.**

Using standard experimental methods, an API2000 mass spectrometer was configured for optimal detection 2-hydroxyglutarate (**Table 6** and **FIG. 19**). The reaction products of the control and enzyme-containing reactions from above were investigated for the presence of 2-hydroxyglutaric acid, **FIG. 20**. In the control reaction, no 2-hydroxyglutaric acid was detected, while in reaction containing R132H, 2-hydroxyglutaric acid was detected. This data confirms that one neoactivity of the R132H mutant is the reduction of  $\alpha$ -KG to 2-hydroxyglutaric acid.

To determine whether R132H mutant protein directly produced 2HG from  $\alpha$ KG, the product of the mutant IDH1 reaction was examined using negative ion mode triple quadrupole electrospray LC-MS. These experiments confirmed that 2HG was the direct product of NADPH-dependent  $\alpha$ KG reduction by the purified R132H mutant protein through comparison with a known metabolite standards (**FIG. 31A**). Conversion of  $\alpha$ KG to isocitrate was not observed.


One can determine the enantiomeric specificity of the reaction product through derivitization with DATAN (diacetyl-L-tartaric acid) and comparing the retention time to that of known R and S standards. This method is described in Struys *et al.* Clin Chem 50:1391-1395(2004). The stereo-specific production of either the R or S enantiomer of alpha-hydroxyglutaric acid by ICDH1 R132H may modify the biological activity of other enzymes present in the cell. The racemic production may also occur.

For example, one can measure the inhibitory effect of alpha-hydroxyglutaric acid on the enzymatic activity of enzymes which utilize  $\alpha$ -KG as a substrate. In one embodiment, alpha-hydroxyglutaric acid may be a substrate- or product- analogue inhibitor of wild-type ICDH1. In another embodiment alpha-hydroxyglutaric acid may be a substrate- or product- analogue inhibitor of HIF1 prolyl hydroxylase. In the former case, inhibition of wild type ICDH1 by the enzymatic product of R132H will reduce the circulating levels of  $\alpha$ KG in the cell. In the latter case, inhibition of HIF1

prolyl hydroxylase will result in the stabilization of HIF1 and an induction of the hypoxic response cohort of cellular responses.

**ICDH R132H reduces  $\alpha$ KG to the R-enantiomer of 2-hydroxyglutarate.**


There are two possible enantiomers of the ICDHR132H reductive reaction product, converting alpha-ketoglutarate to 2-hydroxyglutarate, with the chiral center being located at the alpha-carbon position. Exemplary products are depicted below.



These are referred to by those with knowledge in the art as the R (or pro-R) and S (or pro-S) enantiomers, respectively. In order to determine which form or both is produced as a result of the ICDH1 neoactivity described above, the relative amount of each chiral form in the reaction product was determined in the procedure described below.

Reduction of  $\alpha$ -KG to 2-HG was performed by ICDHR132H in the presence of NADPH as described above, and the reaction progress was monitored by a change in extinction coefficient of the nucleotide cofactor at 340 nM; once the reaction was judged to be complete, the reaction was extracted with methanol and dried down completely in a stream of nitrogen gas. In parallel, samples of chirally pure R-2-HG and a racemic mixture of R- and S-2-HG (produced by a purely chemical reduction of  $\alpha$ -KG to 2-HG) were resuspended in ddH<sub>2</sub>O, similarly extracted with methanol, and dried.

The reaction products or chiral standards were then resuspended in a solution of dichloromethane:acetic acid (4:1) containing 50 g/L DATAN and heated to 75°C for 30 minutes to promote the derivitization of 2-HG in the scheme described below:



After cooling to room temperature, the derivitization reactions were dried to completion and resuspended in ddH<sub>2</sub>O for analysis on an LC-MS/MS system. Analysis of reaction products and chiral standards was performed on an API2000 LC-MS/MS system using a 2 x 150 mM C18 column with an isocratic flow of 200  $\mu$ l/min of 90:10 (ammonium formate, pH 3.6:methanol) and monitoring the retention times of the 2-HG-DATAN complex using XIC and the diagnostic MRM transition of 363/147 in the negative ion mode.

It should be noted that retention times in the experiments described below are approximate and accurate to within +/- 1 minute; the highly reproducible peak seen at 4 minutes is an artefact of a column switching valve whose presence has no result on the conclusions drawn from the experiment.

Injection of the racemic mixture gave two peaks of equal area at retention times of 8 and 10 minutes (**FIG. 24A**), while injection of the R-2-HG standard resulted in a major peak of >95% area at 10 minutes and a minor peak <5% area at 8 minutes (**FIG. 24B**); indicating that the R-2-HG standard is approximately 95% R and 5% S. Thus, this method allows us to separate the R and S-2-HG chiral forms and to determine the relative amounts of each in a given sample. Coinjection of the racemic mixture and the R-2-HG standard resulted in two peaks at 8 and 10 minutes, with a larger peak at 10 minutes resulting from the addition of surplus pro-R-form (the standard) to a previously equal mixture of R- and S-2-HG (**FIG. 24C**). These experiments allow us to assign the 8 minute peak to the S-2-HG form and the 10 minute peak to the R-2-HG form.

Injection of the derivatized neoactivity enzyme reaction product alone yields a single peak at 10 minutes, suggesting that the neoactivity reaction product is chirally pure R-2-HG (**FIG. 24D**). Coinjection of the neoactivity reaction product with the R-



2-HG standard results in a major peak of >95% area at 10 minutes (**FIG. 24E**) and a single minor peak of <5% area at 8 minutes (previously observed in injection of the R-2-HG standard alone) confirming the chirality of the neoactivity product as R. Coinjection of a racemic mixture and the neoactivity reaction product (**FIG. 24F**) results in a 60% area peak at 10 minutes and a 40% area peak at 8 minutes; this deviation from the previously symmetrical peak areas observed in the racemate sample being due to the excess presence of R-2-HG form contributed by the addition of the neoactivity reaction product.

These experiments allow us to conclude that the ICDH1 neoactivity is a highly specific chiral reduction of  $\alpha$ -KG to R-2-HG.

### **Enzyme properties of other IDH1 mutations**

To determine whether the altered enzyme properties resulting from R132H mutation were shared by other R132 mutations found in human gliomas, recombinant R132C, R132L and R132S mutant IDH1 proteins were generated and the enzymatic properties assessed. Similar to R132H mutant protein, R132C, R132L, and R132S mutations all result in a gain-of-function for NADPH-dependent reduction of  $\alpha$ KG (data not shown). Thus, in addition to impaired oxidative decarboxylation of isocitrate, one common feature shared among the IDH1 mutations found in human gliomas is the ability to catalyze direct NADPH-dependent reduction of  $\alpha$ KG.

### **Identification of 2-HG production in glioblastoma cell lines containing the IDH-1 R132H mutant protein.**

*Generation of genetic engineered glioblastoma cell lines expressing wildtype or mutant IDH-1 protein.* A carboxy-terminal Myc-DDK-tagged open reading frame (ORF) clone of human isocitrate dehydrogenase 1 (IDH1; Ref. ID: NM\_005896) cloned in vector pCMV6 was obtained from commercial vendor Origen Inc. Vector pCMV6 contains both kanamycin and neomycin resistance cassettes for selection in both bacterial and mammalian cell systems. Standard molecular biology mutagenesis techniques were utilized to alter the DNA sequence at base pair 364 of the ORF to introduce base pair change from guanine to adenine resulting in a change in the amino acid code at position 132 from arginine (wt) to histidine (mutant; or R132H). Specific DNA sequence alteration was confirmed by standard methods for

DNA sequence analysis. Parental vector pCMV6 (no insert), pCMV6-wt IDH1 or pCMV6-R132H were transfected into immortalized human glioblastoma cell lines ATCC<sup>®</sup> CRL-2610 (LN-18) or HTB-14 (U-87) in standard growth medium (DMEM; Dulbecco's modified Eagles Medium containing 10 % fetal bovine serum). Approximately 24 hrs after transfection, the cell cultures were transitioned to DMEM containing G418 sodium salt at concentrations of either 750 ug/ml (CRL-2610) or 500 ug/ml (HTB-14) to select those cells in culture that expressed the integrated DNA cassette expressing both the neomycin selectable marker and the ORF for human wild type or R132H. Pooled populations of G418 resistant cells were generated and expression of either wild type IDH1 or R132 IDH1 was confirmed by standard Western blot analysis of cell lysates using commercial antibodies recognizing either human IDH1 antigen or the engineered carboxy-terminal MYC-DDK expression tag. These stable clonal pools were then utilized for metabolite preparation and analysis.

***Procedure for metabolite preparation and analysis.*** Glioblastoma cell lines (CRL-2610 and HTB-14) expressing wildtype or mutant IDH-1 protein were grown using standard mammalian tissue culture techniques on DMEM media containing 10% FCS, 25 mM glucose, 4 mM glutamine, and G418 antibiotic (CRL-2610 at 750 ug/mL; HTB-14 at 500 ug/mL) to insure ongoing selection to preserve the transfected mutant expression sequences. In preparation for metabolite extraction experiments, cells were passaged into 10 cm round culture dishes at a density of  $1 \times 10^6$  cells. Approximately 12 hours prior to metabolite extraction, the culture media was changed (8 mL per plate) to DMEM containing 10% dialyzed FCS (10,000 mwco), 5 mM glucose, 4 mM glutamine, and G-418 antibiotic as before; the dialyzed FCS removes multiple small molecules from the culture media and enables cell culture-specific assessment of metabolite levels. The media was again changed 2 hours prior to metabolite extraction. Metabolite extraction was accomplished by quickly aspirating the media from the culture dishes in a sterile hood, immediately placing the dishes in a tray containing dry ice to cool them to  $-80^{\circ}\text{C}$ , and as quickly as possible, adding 2.6 mL of 80% MeOH/20% water, pre-chilled to  $-80^{\circ}\text{C}$  in a dry-ice/acetone bath. These chilled, methanol extracted cells were then physically separated from the culture dish by scraping with a sterile polyethylene cell lifter (Corning #3008), brought into suspension and transferred to a 15 mL conical vial, then chilled to  $-20^{\circ}\text{C}$ . An additional 1.0 mL of 80% MeOH/20% water was applied to the chilled culture dish

and the cell lifting procedure repeated, to give a final extraction volume of 3.6 mL. The extracts were centrifuged at 20,000 x g for 30 minutes to sediment the cell debris, and 3.0 mL of the supernatants was transferred to a screw-cap freezer vial and stored at -80°C until ready for analysis.

In preparation for analysis, the extracts were removed from the freezer and dried on a nitrogen blower to remove methanol. The 100% aqueous samples were analyzed by LCMS as follows. The extract (10 µL) was injected onto a reverse-phase HPLC column (Synergi 150mm x 2 mm, Phenomenex Inc.) and eluted using a linear gradient of LCMS-grade methanol (Buffer B) in Aq. 10 mM tributylamine, 15 mM Acetic acid (Buffer A), running from 3% Buffer B to 95% Buffer B over 45 minutes at 200 µL/min. Eluted metabolite ions were detected using a triple-quadrupole mass spectrometer, tuned to detect in negative mode with multiple-reaction-monitoring mode transition set (MRM's) according to the molecular weights and fragmentation patterns for 38 known central metabolites, including 2-hydroxyglutarate (MRM parameters were optimized by prior infusion of known compound standards). Data was processed using Analyst Software (Applied Biosystems, Inc.) and metabolite signal intensities were converted into absolute concentrations using signal build-up curves from injected mixtures of metabolite standards at known concentrations. Final metabolite concentrations were reported as mean of at least three replicates, +/- standard deviation.

**Results.** Analyses reveal significantly higher levels of 2-HG in cells that express the IDH-1 R132H mutant protein. As shown in **FIG. 26A**, levels of 2-HG in CRL-2610 cell lines expressing the IDH-1 R132H mutant protein are approximately 28-fold higher than identical lines expressing the wild-type protein. Similarly, levels of 2-HG in HTB-14 cell lines expressing the IDH-1 R132H mutant protein are approximately 38-fold higher than identical lines expressing the wild-type protein, as shown in **FIG. 26B**.

### **Evaluation of 2-hydroxyglutarate (2-HG) production in human glioblastoma tumors containing mutations in isocitrate dehydrogenase 1 (IDH1) at amino acid 132.**

Heterozygous somatic mutations at nucleotide position 395 (amino acid codon 132) in the transcript encoding isocitrate dehydrogenase 1 (IDH1) can occur in brain tumors.

**Tissue source:** Human brain tumors were obtained during surgical resection, flash frozen in liquid nitrogen and stored at -80°C. Clinical classification of the tissue as gliomas was performed using standard clinical pathology categorization and grading.

**Genomic sequence analysis to identify brain tumor samples containing either wild type isocitrate dehydrogenase (IDH1) or mutations altering amino acid 132.** Genomic DNA was isolated from 50-100 mgs of brain tumor tissue using standard methods. A polymerase chain reaction (PCR) procedure was then performed on the isolated genomic DNA to amplify a 295 base pair fragment of the genomic DNA that contains both intron and 2<sup>nd</sup> exon sequences of human IDH1 (**FIG. 27**). In **FIG. 27**, intron sequence is shown in lower case font; 2<sup>nd</sup> exon IDH1 DNA sequence is shown in upper case font; forward (5') and reverse (3') primer sequences are shown in underlined font; guanine nucleotide mutated in a subset of human glioma tumors is shown in bold underlined font.

The amplified DNA fragment was then sequenced using standard protocols and sequence alignments were performed to classify the sequences as either wild type or mutant at the guanine nucleotide at base pair 170 of the amplified PCR fragment. Tumors were identified that contained genomic DNA having either two copies of guanine (wild type) or a mixed or monoallelic combination of one IDH1 allele containing guanine and the other an adenine (mutant) sequence at base pair 170 of the amplified product (**Table 15**). The nucleotide change results in a change at amino acid position 132 of human IDH1 protein from arginine (wild type) to histidine (mutant) as has been previously reported.

**Table 15.** Sequence variance at base pair 170 of the amplified genomic DNA from human glioma samples.

Sample ID	Base 170	IDH1 Amino Acid 132	Genotype
1102	G	arginine	wild type
1822	A	histidine	mutant
496	G	arginine	wild type
1874	A	histidine	mutant
816	A	histidine	mutant
534	G	arginine	wild type
AP-1	A	histidine	mutant
AP-2	A	histidine	mutant

***Procedure for metabolite preparation and analysis.*** Metabolite extraction was accomplished by adding a 10 X volume (m/v ratio) of -80 C methanol:water mix (80%:20%) to the brain tissue (approximately 100mgs) followed by 30 s homogenization at 4 C. These chilled, methanol extracted homogenized tissues were then centrifuged at 14,000 rpm for 30 minutes to sediment the cellular and tissue debris and the cleared tissue supernatants were transferred to a screw-cap freezer vial and stored at -80°C. For analysis, a 2X volume of tributylamine (10 mM) acetic acid (10 mM) pH 5.5 was added to the samples and analyzed by LCMS as follows. Sample extracts were filtered using a Millex-FG 0.20 micron disk and 10 µL were injected onto a reverse-phase HPLC column (Synergi 150mm x 2 mm, Phenomenex Inc.) and eluted using a linear gradient LCMS-grade methanol (50%) with 10 mM tributylamine and 10 mM acetic acid) ramping to 80 % methanol:10 mM tributylamine: 10 mM acetic acid over 6 minutes at 200 µL/min. Eluted metabolite ions were detected using a triple-quadrupole mass spectrometer, tuned to detect in negative mode with multiple-reaction-monitoring mode transition set (MRM's) according to the molecular weights and fragmentation patterns for 8 known central metabolites, including 2-hydroxyglutarate (MRM parameters were optimized by prior infusion of known compound standards). Data was processed using Analyst Software (Applied Biosystems, Inc.) and metabolite signal intensities were obtained by standard peak integration methods.

***Results.*** Analyses revealed dramatically higher levels of 2-HG in cells tumor samples that express the IDH-1 R132H mutant protein. Data is summarized in **Table 16** and **FIG. 28**.

Table 16

Sample ID	Primary Specimen Diagnosis	Grade	Tumor Cells in Tumor Foci (%)	Geno-type	Nucleo-tide change	Codon	2HG (□mole/g)	□KG (□mole/g)	Malate (□mole/g)	Fumarate (□mole/g)	Succinate (□mole/g)	Isocitrate (□mole/g)
1	Glioblastoma, residual/recurrent	WHO grade IV	n/a	wild type	wild type	R132	0.18	0.161	1.182	0.923	1.075	0.041
2	Glioblastoma	WHO grade IV	n/a	wild type	wild type	R132	0.16	0.079	1.708	1.186	3.156	0.100
3	Glioblastoma	WHO grade IV	n/a	wild type	wild type	R132	0.13	0.028	0.140	0.170	0.891	0.017
4	Oligoastrocytoma	WHO grade II	n/a	wild type	wild type	R132	0.21	0.016	0.553	1.061	1.731	0.089
5	Glioblastoma	WHO grade IV	n/a	mutant	G364A	R132H	16.97	0.085	1.091	0.807	1.357	0.058
6	Glioblastoma	WHO grade IV	n/a	mutant	G364A	R132H	19.42	0.023	0.462	0.590	1.966	0.073
7	Glioblastoma	WHO grade IV	n/a	mutant	G364A	R132H	31.56	0.068	0.758	0.503	2.019	0.093
8	Oligodendroglioma, anaplastic	WHO grade III	75	mutant	G364A	R132H	12.49	0.033	0.556	0.439	0.507	0.091
9	Oligodendroglioma, anaplastic	WHO grade III	90	mutant	G364A	R132H	4.59	0.029	1.377	1.060	1.077	0.574
10	Oligoastrocytoma	WHO grade II	n/a	mutant	G364A	R132H	6.80	0.038	0.403	0.503	1.561	0.065
11	Glioblastoma	WHO grade IV	n/a	wild type	wild type	R132	0.686	0.686	0.686	0.686	0.686	0.007
12	Glioblastoma	WHO grade IV	n/a	mutant	G364A	R132H	18.791	18.791	18.791	18.791	18.791	0.031
13	Glioblastoma	WHO grade IV	n/a	mutant	G364A	R132H	4.59	0.029	1.377	1.060	1.077	0.043
14	Glioblastoma	WHO grade IV	n/a	wild type	wild type	R132	0.199	0.046	0.180	0.170	0.221	0.014
15	Glioblastoma	WHO grade IV	n/a	mutant	C363G	R132G	13.827	0.030	0.905	0.599	1.335	0.046
16	Glioblastoma	WHO grade IV	n/a	mutant	G364A	R132H	28.364	0.068	0.535	0.488	2.105	0.054
17	Glioblastoma	WHO grade IV	n/a	mutant	C363A	R132S	9.364	0.029	1.038	0.693	2.151	0.121
18	Glioblastoma	WHO grade IV	n/a	wild type	wild type	R132	0.540	0.031	0.468	0.608	1.490	0.102
19	Glioma, malignant, astrocytoma	WHO grade IV	80	mutant	G364A	R132H	19.000	0.050	0.654	0.391	2.197	0.171
20	Oligodendroglioma	WHO grade III	80	wild type	wild type	R132	0.045	0.037	1.576	0.998	1.420	0.018
21	Glioma, malignant, astrocytoma	WHO grade	95	wild type	wild type	R132	0.064	0.034	0.711	0.710	2.105	0.165

		IV										
22	Glioblastoma	WHO grade IV	70	wild type	wild type	R132	0.171	0.041	2.066	1.323	0.027	0.072

To determine if 2HG production is characteristic of tumors harboring mutations in IDH1, metabolites were extracted from human malignant gliomas that were either wild-type or mutant for IDH1. It has been suggested that  $\alpha$ KG levels are decreased in cells transfected with mutant IDH1 (Zhao, S. et al. Science 324, 261-5 (2009)). The average  $\alpha$ KG level from 12 tumor samples harboring various R132 mutations was slightly less than the average  $\alpha$ KG level observed in 10 tumors which are wild-type for IDH1. This difference in  $\alpha$ KG was not statistically significant, and a range of  $\alpha$ KG levels was observed in both wild-type and mutant tumors. In contrast, increased 2HG levels were found in all tumors that contained an R132 IDH1 mutation. All R132 mutant IDH1 tumors examined had between 5 and 35  $\mu$ mol of 2HG per gram of tumor, while tumors with wild-type IDH1 had over 100 fold less 2HG. This increase in 2HG in R132 mutant tumors was statistically significant ( $p < 0.0001$ ). It was confirmed that (R)-2HG was the isomer present in tumor samples (data not shown). Together these data establish that the novel enzymatic activity associated with R132 mutations in IDH1 results in the production of 2HG in human brain tumors that harbor these mutations.

2HG is known to accumulate in the inherited metabolic disorder 2-hydroxyglutaric aciduria. This disease is caused by deficiency in the enzyme 2-hydroxyglutarate dehydrogenase, which converts 2HG to  $\alpha$ KG (Struys, E. A. et al. Am J Hum Genet 76, 358-60 (2005)). Patients with 2-hydroxyglutarate dehydrogenase deficiencies accumulate 2HG in the brain as assessed by MRI and CSF analysis, develop leukoencephalopathy, and have an increased risk of developing brain tumors (Aghili, M., Zahedi, F. & Rafiee, J Neurooncol 91, 233-6 (2009); Kolker, S., Mayatepek, E. & Hoffmann, G. F. Neuropediatrics 33, 225-31 (2002); Wajner, M., Latini, A., Wyse, A. T. & Dutra-Filho, C. S. J Inherit Metab Dis 27, 427-48 (2004)). Furthermore, elevated brain levels of 2HG result in increased ROS levels (Kolker, S. et al. Eur J Neurosci 16, 21-8 (2002); Latini, A. et al. Eur J Neurosci 17, 2017-22 (2003)), potentially contributing to an increased risk of cancer. The ability of 2HG to act as an NMDA receptor agonist may contribute to this effect (Kolker, S. et al. Eur J

Neurosci 16, 21-8 (2002)). 2HG may also be toxic to cells by competitively inhibiting glutamate and/or  $\alpha$ KG utilizing enzymes. These include transaminases which allow utilization of glutamate nitrogen for amino and nucleic acid biosynthesis, and  $\alpha$ KG-dependent prolyl hydroxylases such as those which regulate Hif1 $\alpha$  levels. Alterations in Hif1 $\alpha$  have been reported to result from mutant IDH1 protein expression (Zhao, S. et al. Science 324, 261-5 (2009)). Regardless of mechanism, it appears likely that the gain-of-function ability of cells to produce 2HG as a result of R132 mutations in IDH1 contributes to tumorigenesis. Patients with 2-hydroxyglutarate dehydrogenase deficiency have a high risk of CNS malignancy (Aghili, M., Zahedi, F. & Rafiee, E. J Neurooncol 91, 233-6 (2009)). The ability of mutant IDH1 to directly act on  $\alpha$ KG may explain the prevalence of IDH1 mutations in tumors from CNS tissue, which are unique in their high level of glutamate uptake and its ready conversion to  $\alpha$ KG in the cytosol (Tsacopoulos, M. J Physiol Paris 96, 283-8 (2002)), thereby providing high levels of substrate for 2HG production. The apparent co-dominance of the activity of mutant IDH1 with that of the wild-type enzyme is consistent with the genetics of the disease, in which only a single copy of the gene is mutated. As discussed above, the wild-type IDH1 could directly provide NADPH and  $\alpha$ KG to the mutant enzyme. These data also demonstrate that mutation of R132 to histidine, serine, cysteine, glycine or leucine share a common ability to catalyze the NADPH-dependent conversion of  $\alpha$ KG to 2HG. These findings help clarify why mutations at other amino acid residues of IDH1, including other residues essential for catalytic activity, are not found. Finally, these findings have clinical implications in that they suggest that 2HG production will identify patients with IDH1 mutant brain tumors. This will be important for prognosis as patients with IDH1 mutations live longer than patients with gliomas characterized by other mutations (Parsons, D. W. et al. Science 321, 1807-12 (2008)). In addition, patients with lower grade gliomas may benefit by the therapeutic inhibition of 2HG production. Inhibition of 2HG production by mutant IDH1 might slow or halt conversion of lower grade glioma into lethal secondary glioblastoma, changing the course of the disease.

**The reaction product of ICDH1 R132H reduction of  $\alpha$ -KG inhibits the oxidative decarboxylation of isocitrate by wild-type ICDH1.**



A reaction containing the wild-type ICDH1, NADP, and  $\alpha$ -KG was assembled (under conditions as described above) to which was added in a titration series either (R)-2-hydroxyglutarate or the reaction product of the ICDH1 R132H mutant reduction of  $\alpha$ -KG to 2-hydroxyglutarate. The reaction product 2-HG was shown to inhibit the oxidative decarboxylation of isocitrate by the wild-type ICDH1, while the (R)-2-hydroxyglutarate did not show any effect on the rate of the reaction. Since there are only two possible chiral products of the ICDH1 R132H mutant reduction of  $\alpha$ -KG to 2-HG, and the (R)-2-HG did not show inhibition in this assay, it follows that the product of the mutant reaction is the (S)-2-HG form. This experiment is presented in **FIG. 25**.

To determine the chirality of the 2HG produced, the products of the R132H reaction was derivatized with diacetyl-L-tartaric anhydride, which allowed separating the (S) and (R) enantiomers of 2HG by simple reverse-phase LC and detecting the products by tandem mass spectrometry (Struys, E. A., Jansen, E. E., Verhoeven, N. M. & Jakobs, C. Clin Chem 50, 1391-5 (2004)) (**FIG. 31B**). The peaks corresponding to the (S) and (R) isomers of 2HG were confirmed using racemic and R(-)-2HG standards. The reaction product from R132H co-eluted with R(-)-2HG peak, demonstrating that the R(-) stereoisomer is the product produced from  $\alpha$ KG by R132H mutant IDH1.

The observation that the reaction product of the mutant enzyme is capable of inhibiting a metabolic reaction known to occur in cells suggests that this reaction product might also inhibit other reactions which utilize  $\alpha$ -KG, isocitrate, or citrate as substrates or produce them as products in vivo or in vitro.

### EXAMPLE 3 METABOLOMICS ANALYSIS OF IDH1 WILD TYPE AND MUTANTS

Metabolomics research can provide mechanistic basis for why R132 mutations confer survival advantage for GBM patients carrying such mutations.

#### ***1. Metabolomics of GBM tumor cell lines: wild type vs R132 mutants***

Cell lines with R132 mutations can be identified and profiled. Experiments can be performed in proximal metabolite pool with a broad scope of metabolites.

#### ***2. Oxalomalate treatment of GBM cell lines***

Oxalomalate is a competitive inhibitor of IDH1. Change of NADPH (metabolomics) when IDH1 is inhibited by a small molecule can be examined.

***3. Metabolomics of primary GBM tumors: wild type vs R132 mutations***

Primary tumors with R132 mutations can be identified. Experiments can be performed in proximal metabolite pool with a broad scope of metabolites.

***4. Detection of 2-hydroxyglutarate in cells that overexpress IDH1 132 mutants***

Overexpression of an IDH1 132 mutant in cells may cause an elevated level of 2-hydroxyglutarate and/or a reduced level of alpha-ketoglutarate. One can perform a metabolomic experiment to demonstrate the consequence of this mutation on the cellular metabolite pool.

EXAMPLE 4 EVALUATION OF IDH1 AS A CANCER TARGET

shRNAmir inducible knockdown can be performed to examine the cellular phenotype and metabolomics profiles. HTS grade IDH1 enzymes are available. The IDH mutations described herein can be used for patient selection.

EXAMPLE 5 siRNAs

IDH1

Exemplary siRNAs are presented in the following tables. Art-known methods can be used to select other siRNAs. siRNAs can be evaluated, *e.g.*, by determining the ability of an siRNA to silence an IDH, *e.g.*, IDH1, *e.g.*, in an *in vitro* system, *e.g.*, in cultured cells, *e.g.*, HeLa cells or cultured glioma cells. siRNAs known in the art for silencing the target can also be used, see, *e.g.*, *Silencing of cytosolic NADP+ dependent isocitrate dehydrogenase by small interfering RNA enhances the sensitivity of HeLa cells toward stauropine*, Lee *et al.*, 2009, Free Radical Research, 43: 165-173.

The siRNAs in **Table 7** (with the exception of entry 1356) were generated using the siRNA selection tool available on the worldwide web at [jura.wi.mit.edu/bioc/siRNAext/](http://jura.wi.mit.edu/bioc/siRNAext/). (Yuan *et al.* Nucl. Acids. Res. 2004 32:W130-W134.) Other selection tools can be used as well. Entry 1356 was adapted from *Silencing of cytosolic NADP+ dependent isocitrate dehydrogenase by small interfering RNA enhances the sensitivity of HeLa cells toward stauropine*, Lee *et al.*, 2009, Free Radical Research, 43: 165-173.

The siRNAs in Tables 7, 8, 9, 10, 11, 12, 13 and 14 represent candidates spanning the IDH1 mRNA at nucleotide positions 628 and 629 according to the sequence at GenBank Accession No. NM\_005896.2 (SEQ ID NO:9, FIG. 22).

The RNAs in the tables can be modified, *e.g.*, as described herein. Modifications include chemical modifications to enhance properties, *e.g.*, resistance to degradation, or the use of overhangs. For example, either one or both of the sense and antisense strands in the tables can include an additional dinucleotide at the 3' end, *e.g.*, TT, UU, dTdT.

**Table 7. siRNAs targeting wildtype IDH1**

<b>Position on mRNA (FIG. 21B)</b>	<b>sense (5' to 3')</b>	<b>SEQ ID NO:</b>	<b>antisense (5' to 3')</b>	<b>SEQ ID NO:</b>
13	GGUUUCUGCAGAGUCUAC U	14	AGUAGACUCUGCAGAAAC C	15
118	CUCUUCGCCAGCAUAUCA U	16	AUGAUAUGCUGGCGAAGA G	17
140	GGCAGGCGAUAAACUACA U	18	AUGUAGUUUAUCGCCUGC C	19
145	GCGAUAAACUACAUUCAG U	20	ACUGAAUGUAGUUUAUCG C	21
199	GAAUCUAUUCACUGUCA A	22	UUGACAGUGAAUAGAUUU C	23
257	GUUCUGUGGUAGAGAUGC A	24	UGCAUCUCUACCACAGAA C	25
272	GCAAGGAGAUGAAAUGAC A	26	UGUCAUUUCAUCUCCUUG C	27
277	GGAGAUGAAAUGACACGA A	28	UUCGUGUCAUUUCAUCUC C	29
278	GAGAUGAAAUGACACGAA U	30	AUUCGUGUCAUUUCAUCU C	31
280	GAUGAAAUGACACGAAUC A	32	UGAUUCGUGUCAUUUCAU C	33
292	CGAAUCAUUUGGAAUUG A	34	UCAAUCCCAAUGAUUC G	35
302	GGGAAUUGAUUAAAGAGA A	36	UUCUCUUUAAUCAAUUCC C	37
332	CCUACGUGGAAUUGGAUC U	38	AGAUCCAAUCCACGUAG G	39
333	CUACGUGGAAUUGGAUCU A	40	UAGAUCCAAUCCACGUA G	41
345	GGAUUCACAUAGCUAUGA U	42	AUCAUAGCUAUGUAGAUC C	43
356	GCUAUGAUUUAGGCAUAG A	44	UCUAUGCCUAAAUCAUAG C	45
408	GGAUGCUGCAGAAGCUAU A	46	UAUAGCUUCUGCAGCAUC C	47
416	CAGAAGCUAUAAGAAGC A	48	UGCUCUUUAUAGCUUCU G	49
418	GAAGCUAUAAGAAGCAU A	50	UAUGCUUCUUUAUAGCUU C	51
432	GCAUAAUGUUGGCGUCAA A	52	UUUGACGCCAACAUUAUG C	53
467	CUGAUGAGAAGAGGGUUG A	54	UCAACCCUCUUCUCAUCA G	55
481	GUUGAGGAGUUCAAGUUG A	56	UCAACUUGAACUCCUCA C	57
487	GAGUUCAAGUUGAAACAA A	58	UUUGUUUCAACUUGAACU C	59
495	GUUGAAACAAUGUGGAA A	60	UUUCCACAUUUGUUUCA C	61
502	CAAAUGUGGAAAUACCA A	62	UUGGUGAUUCCACAUUU G	63
517	CCAAAUGGCACCAUACGA A	64	UUCGUAUGGUGCCAUUUG G	65
528	CAUACGAAAUUUCUGGG A	66	ACCCAGAAAUUUCGUAU U	67

	U		G	
560	GAGAAGCCAUUAUCUGCA A	68	UUGCAGAUAAUGGCUUCU C	69
614	CUAUCAUCAUAGGUCGUC A	70	UGACGACCUAUGAUGAUA G	71
618	CAUCAUAGGUCGUCAUGC U	72	AGCAUGACGACCUAUGAU G	73
621	CAUAGGUCGUCAUGCUUA U	74	AUAAGCAUGACGACCUAU G	75
691	GAGUAACCUACACACCA A	76	UUGGUGUGUAGGUUAUCU C	77
735	CCUGGUACAUAAACUUUGA A	78	UUCAAAGUUAUGUACCAG G	79
747	CUUUGAAGAAGGUGGUGG U	80	ACCACCACCUUCUUCAAA G	81
775	GGGAUGUAUUAACAAGAU A	82	UAUCUUGAUUAUACAUC C	83
811	GCACACAGUCCUUCCAA A	84	TUUGGAAGGAACUGUGUG C	85
818	GUUCCUUCCAAUAGGCUC U	86	AGAGCCAUUUGGAAGGAA C	87
844	GGUUGGCCUUUGUAUCUG A	88	UCAGAUACAAAGGCCAAC C	89
851	CUUUGUAUCUGAGACCA A	90	UUGGUGCUCAGAUACAAA G	91
882	GAAGAAAUAUGAUGGGCG U	92	ACGCCCAUCAUAUUUCU C	93
942	GUCCCAGUUUGAAGCUCA A	94	UUGAGCUUCAACUGGGA C	95
968	GGUAUGAGCAUAGGCUCA U	96	AUGAGCCUAUGCUCAUAC C	97
998	GGCCCAAGCUAUGAAAUC A	98	UGAUUUCAUAGCUUGGGC C	99
1001	CCCAAGCUAUGAAAUCAG A	100	UCUGAUUUCAUAGCUUGG G	101
1127	CAGAUGGCAAGACAGUAG A	102	UCUACUGUCUUGCCAUCU G	103
1133	GCAAGACAGUAGAAGCAG A	104	UCUGCUUCUACUGUCUUG C	105
1184	GCAUGUACCAGAAAGGAC A	106	UGUCCUUUCUGGUACAUG C	107
1214	CCAAUCCCAUUGCUUCCA U	108	AUGGAAGCAAUGGGAUUG G	109
1257	CCACAGAGCAAAGCUUGA U	110	AUCAAGCUUUGCUCUGUG G	111
1258	CACAGAGCAAAGCUUGAU A	112	UAUCAAGCUUUGCUCUGU G	113
1262	GAGCAAAGCUUGUAACA A	114	UUGUUAUCAAGCUUUGCU C	115
1285	GAGCUUGCCUUCUUUGCA A	116	UUGCAAAGAAGGCAAGCU C	117
1296	CUUUGCAAUUGCUUUGGA A	118	TUCCAAAGCAUUGCAA G	119
1301	CAAUUGCUUUGGAAGAAG U	120	ACUUCUCCAAAGCAUUU G	121
1307	CUUUGGAAGAAGUCUCUA U	122	AUAGAGACUUCUCCAAA G	123
1312	GAAGAAGUCUCUAUUGAG A	124	UCUCAAUAGAGACUUCU C	125

1315	GAAGUCUCUAUUGAGACA A	126	UUGUCUCAAUAGAGACUU C	127
1356	GGACUUGGCUGCUUGCAU U	128	AAUGCAAGCAGCCAAGUC C	129
1359	CUUGGCUGCUUGCAUAAA A	130	UUUAAUGCAAGCAGCCAA G	131
1371	CAUAAAAGGUUUACCCAA U	132	AUUGGGUAAAACCUUUAU G	133
1385	CCAAUGUGCAACGUUCUG A	134	UCAGAACGUUGCACAUG G	135
1390	GUGCAACGUUCUGACUAC U	136	AGUAGUCAGAACGUUGCA C	137
1396	CGUUCUGACUACUUGAAU A	138	UAUUCAAGUAGUCAGAAC G	139
1415	CAUUUGAGUUCUUGGAUA A	140	UUAUCCAUGAACUCAAAU G	141
1422	GUUCAUGGAUAAAACUUGG A	142	UCCAAGUUUAUCCAUGAA C	143
1425	CAUGGAUAAAACUUGGAGA A	144	UUCUCCAAGUUUAUCCA G	145
1455	CAAACUAGCUCAGGCCAA A	146	UUUGGCCUGAGCUAGUUU G	147
1487	CCUGAGCUAAGAAGGAUA A	148	UUAUCCUUCUAGCUCAG G	149
1493	CUAAGAAGGAUAAUUGUC U	150	AGACAAUUAUCCUUCUUA G	151
1544	CUGUGUACACUCAAGGA U	152	AUCCUUGAGUGUAACACA G	153
1546	GUGUUACACUCAAGGAUA A	154	UUAUCCUUGAGUGUAACA C	155
1552	CACUCAAGGAUAAAGGCA A	156	UUGCCUUUAUCCUUGAGU G	157
1581	GUAUUUGUUUAGAAGCC A	158	UGGCUUCUAAACAAAUUA C	159
1646	GUUAUUGCCACCUUUGUG A	160	UCACAAAGGUGGCAAUAA C	161
1711	CAGCCUAGGAAUUCGGUU A	162	UAACCGAAUCCUAGGCU G	163
1713	GCCUAGGAAUUCGGUUAG U	164	ACUAACCGAAUCCUAGG C	165
1714	CCUAGGAAUUCGGUUAGU A	166	UACUAACCGAAUCCUAG G	167
1718	GGAAUUCGGUUAGUACUC A	168	UGAGUACUAACCGAAUUC C	169
1719	GAAUUCGGUUAGUACUCA U	170	AUGAGUACUAACCGAAU C	171
1725	GGUUAGUACUCAUUUGUA U	172	AUACAAAUGAGUACUAAC C	173
1730	GUACUCAUUUGUAUUCAC U	174	AGUGAAUACAAAUGAGUA C	175
1804	GGUAAAUGAUAGCCACAG U	176	ACUGUGGCUAUCAUUUAC C	177
1805	GUAAAUGAUAGCCACAGU A	178	UACUGUGGCUAUCAUUUA C	179
1816	CCACAGUAUUGCUCCCUA A	180	UUAGGGAGCAAUCUGUG G	181
1892	GGGAAGUUCUGGUGUCAU A	182	UAUGACACCAGAACUCC C	183
1897	GUUCUGGUGUCAUAGAU A	184	AUAUCUAUGACACCAGAA A	185

	U		C	
1934	GCUGUGCAUUAACUUGC A	186	UGCAAGUUUAAUGCACAG C	187
1937	GUGCAUUAACUUGCACA U	188	AUGUGCAAGUUUAAUGCA C	189
1939	GCAUUAACUUGCACAUG A	190	UCAUGUGCAAGUUUAAUG C	191
1953	CAUGACUGGAACGAAGUA U	192	AUACUUCGUUCCAGUCAU G	193
1960	GGAACGAAGUAUGAGUGC A	194	UGCACUCAUACUUCGUUC C	195
1961	GAACGAAGUAUGAGUGCA A	196	UUGCACUCAUACUUCGUU C	197
1972	GAGUGCAACUCAAAUGUG U	198	ACACAUUUGAGUUGCACU C	199
1976	GCAACUCAAAUGUGUUGA A	200	UUCAACACAUUUGAGUUG C	201
1982	CAAUGUGUUGAAGAUAC U	202	AGUAUCUUCACACAUUU G	203
1987	GUGUUGAAGAUACUGCAG U	204	ACUGCAGUAUCUUCAACA C	205
1989	GUUGAAGAUACUGCAGUC A	206	UGACUGCAGUAUCUUCAA C	207
2020	CCUUGCUGAAUGUUUCCA A	208	UUGGAAACAUUCAGCAAG G	209
2021	CUUGCUGAAUGUUUCCA U	210	AUUGGAAACAUUCAGCAA G	211
2024	GCUGAAUGUUUCCAAUAG A	212	UCUAUUGGAAACAUUCAG C	213
2035	CCAAUAGACUAAAUCUG U	214	ACAGUAUUUAGUCUAUUG G	215
2067	GAGUUUGGAAUCCGGAU A	216	UAUCCGGAUUCCAAACU C	217
2073	GGAAUCCGGAUAAAUC U	218	AGUAUUUAUUCGGAUUC C	219
2074	GAAUCCGGAUAAAUCU A	220	UAGUAUUUAUUCGGAUU C	221
2080	GGAAUAAAUCUACCUGG A	222	UCCAGGUAGUAUUUAUUC C	223
2133	GGCCUGGCCUGAAUAUUA U	224	AUAAUAUUCAGGCCAGGC C	225
2134	GCCUGAAUAUUAUCUAC U	226	AGUAGUAUAAUAUUCAGG C	227
2136	CUGGCCUGAAUAUUAUC U	228	AGUAUAAUAUUCAGGCCA G	229
2166	CAUAUUUCAUCCAAGUGC A	230	UGCACUUGGAUGAAUAU G	231
2180	GUGCAAUAUGUAAGCUG A	232	UCAGCUUACAUUAUUGCA C	233
2182	GCAAUAUGUAAGCUGAA U	234	AUUCAGCUUACAUUAUUG C	235
2272	CACUAUCUUAUCUUCUCC U	236	AGGAGAAGUAAGAUAGU G	237
2283	CUUCUCCUGAACUGUUGA U	238	AUCAACAGUUCAGGAGAA G	239

**Table 8. siRNAs targeting wildtype IDH1**

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUC G	240	CGACCUAUGAUGAUAGGU U	241
612	ACCUAUCAUCAUAGGUCG U	242	ACGACCUAUGAUGAUAGG U	243
613	CCUAUCAUCAUAGGUCGU C	244	GACGACCUAUGAUGAUAG G	245
614	CUAUCAUCAUAGGUCGUC A	246	UGACGACCUAUGAUGAUA G	247
615	UAUCAUCAUAGGUCGUCA U	248	AUGACGACCUAUGAUGAU A	249
616	AUCAUCAUAGGUCGUCAU G	250	CAUGACGACCUAUGAUGA U	251
617	UCAUCAUAGGUCGUCAUG C	252	GCAUGACGACCUAUGAUG A	253
618	CAUCAUAGGUCGUCAUGC U	254	AGCAUGACGACCUAUGAU G	255
619	AUCAUAGGUCGUCAUGCU U	256	AAGCAUGACGACCUAUGA U	257
620	UCAUAGGUCGUCAUGCUU A	258	UAAGCAUGACGACCUAUG A	259
621	CAUAGGUCGUCAUGCUIA U	260	AUAAGCAUGACGACCUAU G	261
622	AUAGGUCGUCAUGCUIAU G	262	CAUAAGCAUGACGACCUA U	263
623	UAGGUCGUCAUGCUIAUG G	264	CCAUAAGCAUGACGACCU A	265
624	AGGUCGUCAUGCUIAUGG G	266	CCCAUAAGCAUGACGACC U	267
625	GGUCGUCAUGCUIAUGGG G	268	CCCCAUAAGCAUGACGAC C	269
626	GUCGUCAUGCUIAUGGGG A	270	UCCCAUAAGCAUGACGAC C	271
627	UCGUCAUGCUIAUGGGGA U	272	AUCCCAUAAGCAUGACGA C	273

**Table 9. siRNAs targeting G395A mutant IDH1 (SEQ ID NO:5) (equivalent to G629A of SEQ ID NO:9 (FIG. 21B))**

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUCA	274	UGACCUAUGAUGAUAGGUU	275
612	ACCUAUCAUCAUAGGUCAU	276	AUGACCUAUGAUGAUAGGU	277
613	CCUAUCAUCAUAGGUCAUC	278	GAUGACCUAUGAUGAUAGG	279
614	CUAUCAUCAUAGGUCAUCA	280	UGAUGACCUAUGAUGAUAG	281
615	UAUCAUCAUAGGUCAUCAU	282	AUGAUGACCUAUGAUGAUA	283



616	AUCAUCAUAGGUCAUCAUG	284	CAUGAUGACCUAUGAUGAU	285
617	UCAUCAUAGGUCAUCAUGC	286	GCAUGAUGACCUAUGAUGA	287
618	CAUCAUAGGUCAUCAUGCU	288	AGCAUGAUGACCUAUGAUG	289
619	AUCAUAGGUCAUCAUGCUU	290	AAGCAUGAUGACCUAUGAU	291
620	UCAUAGGUCAUCAUGCUUA	292	UAAGCAUGAUGACCUAUGA	293
621	CAUAGGUCAUCAUGCUUAU	294	AUAAGCAUGAUGACCUAUG	295
622	AUAGGUCAUCAUGCUUAUG	296	CAUAAGCAUGAUGACCUAU	297
623	UAGGUCAUCAUGCUUAUGG	298	CCAUAGCAUGAUGACCUA	299
624	AGGUCAUCAUGCUUAUGGG	300	CCCAUAGCAUGAUGACCU	301
625	GGUCAUCAUGCUUAUGGGG	302	CCCCAUAGCAUGAUGACC	303
626	GUUCAUCAUGCUUAUGGGGA	304	UCCCCAUAGCAUGAUGAC	305
627	UCAUCAUGCUUAUGGGGAU	306	AUCCCCAUAGCAUGAUGA	307

**Table 10.** siRNAs targeting C394A mutant IDH1 (SEQ ID NO:5) (equivalent to C628A of SEQ ID NO:9 (FIG. 21B)) (Arg132Ser (SEQ ID NO:8))

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUAG	308	CUACCUAUGAUGAUAGGUU	309
612	ACCUAUCAUCAUAGGUAGU	310	ACUACCUAUGAUGAUAGGU	311
613	CCUAUCAUCAUAGGUAGUC	312	GACUACCUAUGAUGAUAGG	313
614	CUAUCAUCAUAGGUAGUCA	314	UGACUACCUAUGAUGAUAG	315
615	UAUCAUCAUAGGUAGUCAU	316	AUGACUACCUAUGAUGAUA	317
616	AUCAUCAUAGGUAGUCAUG	318	CAUGACUACCUAUGAUGAU	319
617	UCAUCAUAGGUAGUCAUGC	320	GCAUGACUACCUAUGAUGA	321
618	CAUCAUAGGUAGUCAUGCU	322	AGCAUGACUACCUAUGAUG	323
619	AUCAUAGGUAGUCAUGCUU	324	AAGCAUGACUACCUAUGAU	325
620	UCAUAGGUAGUCAUGCUUA	326	UAAGCAUGACUACCUAUGA	327
621	CAUAGGUAGUCAUGCUUAU	328	AUAAGCAUGACUACCUAUG	329
622	AUAGGUAGUCAUGCUUAUG	330	CAUAAGCAUGACUACCUAU	331
623	UAGGUAGUCAUGCUUAUGG	332	CCAUAGCAUGACUACCUA	333
624	AGGUAGUCAUGCUUAUGGG	334	CCCAUAGCAUGACUACCU	335
625	GGUAGUCAUGCUUAUGGGG	336	CCCCAUAGCAUGACUACC	337
626	GUAGUCAUGCUUAUGGGGA	338	UCCCCAUAGCAUGACUAC	339
627	UAGUCAUGCUUAUGGGGAU	340	AUCCCCAUAGCAUGACUA	341

**Table 11.** siRNAs targeting C394U mutant IDH1 (SEQ ID NO:5) (equivalent to C628U of SEQ ID NO:9 (FIG. 21B)) (Arg132Cys (SEQ ID NO:8))

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUUG	342	CAACCUAUGAUGAUAGGUU	343
612	ACCUAUCAUCAUAGGUUGU	344	ACAACCUAUGAUGAUAGGU	345
613	CCUAUCAUCAUAGGUUGUC	346	GACAACCUAUGAUGAUAGG	347
614	CUAUCAUCAUAGGUUGUCA	348	UGACAACCUAUGAUGAUAG	349
615	UAUCAUCAUAGGUUGUCAU	350	AUGACAACCUAUGAUGAUA	351
616	AUCAUCAUAGGUUGUCAUG	352	CAUGACAACCUAUGAUGAU	353
617	UCAUCAUAGGUUGUCAUGC	354	GCAUGACAACCUAUGAUGA	355
618	CAUCAUAGGUUGUCAUGCU	356	AGCAUGACAACCUAUGAUG	357
619	AUCAUAGGUUGUCAUGCUU	358	AAGCAUGACAACCUAUGAU	359
620	UCAUAGGUUGUCAUGCUUA	360	UAAGCAUGACAACCUAUGA	361
621	CAUAGGUUGUCAUGCUUAU	362	AUAAGCAUGACAACCUAUG	363
622	AUAGGUUGUCAUGCUUAUG	364	CAUAAGCAUGACAACCUAU	365
623	UAGGUUGUCAUGCUUAUGG	366	CCAUAGCAUGACAACCUA	367
624	AGGUUGUCAUGCUUAUGGG	368	CCCAUAAGCAUGACAACCU	369
625	GGUUGUCAUGCUUAUGGGG	370	CCCCAUAAGCAUGACAACC	371
626	GUUGUCAUGCUUAUGGGGA	372	UCCCCAUAAGCAUGACAAC	373
627	UUGUCAUGCUUAUGGGGAU	374	AUCCCCAUAAGCAUGACAA	375

**Table 12.** siRNAs targeting C394G mutant IDH1 (SEQ ID NO:5) (equivalent to C628G of SEQ ID NO:9 (FIG. 21B)) (Arg132Gly (SEQ ID NO:8))

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUG G	376	CCACCUAUGAUGAUAGGU U	377
612	ACCUAUCAUCAUAGGUGG U	378	ACCACCUAUGAUGAUAGG U	379
613	CCUAUCAUCAUAGGUGGU C	380	GACCACCUAUGAUGAUAG G	381
614	CUAUCAUCAUAGGUGGUC A	382	UGACCACCUAUGAUGAUA G	383
615	UAUCAUCAUAGGUGGUCA U	384	AUGACCACCUAUGAUGAU A	385
616	AUCAUCAUAGGUGGUCAU G	386	CAUGACCACCUAUGAUGA U	387
617	UCAUCAUAGGUGGUCAUG C	388	GCAUGACCACCUAUGAUG A	389
618	CAUCAUAGGUGGUCAUGC U	390	AGCAUGACCACCUAUGAU G	391
619	AUCAUAGGUGGUCAUGCU	392	AAGCAUGACCACCUAUGA	393

	U		U	
620	UCAUAGGUGGUCAUGCUU A	394	UAAGCAUGACCACCUAUG A	395
621	CAUAGGUGGUCAUGCUUA U	396	AUAAGCAUGACCACCUAU G	397
622	AUAGGUGGUCAUGCUUAU G	398	CAUAAGCAUGACCACCUA U	399
623	UAGGUGGUCAUGCUUAUG G	400	CCAUAAGCAUGACCACCU A	401
624	AGGUUGUCAUGCUUAUGG G	402	CCCAUAAGCAUGACCACC U	403
625	GGUUGUCAUGCUUAUGGG G	404	CCCCAUAAGCAUGACCAC C	405
626	GUUGUCAUGCUUAUGGGG A	406	UCCCCAUAAGCAUGACCA C	407
627	UUGUCAUGCUUAUGGGGA U	408	AUCCCCAUAAGCAUGACC A	409

**Table 13.** siRNAs targeting G395C mutant IDH1 (SEQ ID NO:5) (equivalent to G629C of SEQ ID NO:9 (FIG. 21B)) (Arg132Pro (SEQ ID NO:8))

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUC G	410	CGACCUAUGAUGAUAGGU U	411
612	ACCUAUCAUCAUAGGUCG U	412	ACGACCUAUGAUGAUAGG U	413
613	CCUAUCAUCAUAGGUCGU C	414	GACGACCUAUGAUGAUAG G	415
614	CUAUCAUCAUAGGUCGUC A	416	UGACGACCUAUGAUGAUA G	417
615	UAUCAUCAUAGGUCGUCA U	418	AUGACGACCUAUGAUGAU A	419
616	AUCAUCAUAGGUCGUCAU G	420	CAUGACGACCUAUGAUGA U	421
617	UCAUCAUAGGUCGUCAUG C	422	GCAUGACGACCUAUGAUG A	423
618	CAUCAUAGGUCGUCAUGC U	424	AGCAUGACGACCUAUGAU G	425
619	AUCAUAGGUCGUCAUGCU U	426	AAGCAUGACGACCUAUGA U	427
620	UCAUAGGUCGUCAUGCUU A	428	UAAGCAUGACGACCUAUG A	429
621	CAUAGGUCGUCAUGCUUA U	430	AUAAGCAUGACGACCUAU G	431
622	AUAGGUCGUCAUGCUUAU G	432	CAUAAGCAUGACGACCUA U	433
623	UAGGUCGUCAUGCUUAUG G	434	CCAUAAGCAUGACGACCU A	435
624	AGGUCGUCAUGCUUAUGG G	436	CCCAUAAGCAUGACGACC U	437
625	GGUCGUCAUGCUUAUGGG G	438	CCCCAUAAGCAUGACGAC C	439

626	GUCGUCAUGCUUAUGGGG A	440	UCCCCAUAAGCAUGACGA C	441
627	UCGUCAUGCUUAUGGGGA U	442	AUCCCCAUAAGCAUGACG A	443

**Table 14.** siRNAs targeting G395U mutant IDH1 (SEQ ID NO:5) (equivalent to G629U of SEQ ID NO:9 (FIG. 21B)) (Arg132Leu (SEQ ID NO:8))

Position on mRNA (FIG. 21B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
611	AACCUAUCAUCAUAGGUC U	444	AGACCUAUGAUGAUAGGU U	445
612	ACCUAUCAUCAUAGGUCU U	446	AAGACCUAUGAUGAUAGG U	447
613	CCUAUCAUCAUAGGUCUU C	448	GAGACCUAUGAUGAUAG G	449
614	CUAUCAUCAUAGGUCUUC A	450	UGAAGACCUAUGAUGAUA G	451
615	UAUCAUCAUAGGUCUUCA U	452	AUGAAGACCUAUGAUGAU A	453
616	AUCAUCAUAGGUCUUCAU G	454	CAUGAAGACCUAUGAUGA U	455
617	UCAUCAUAGGUCUUCAUG C	456	GCAUGAAGACCUAUGAUG A	457
618	CAUCAUAGGUCUUCAUGC U	458	AGCAUGAAGACCUAUGAU G	459
619	AUCAUAGGUCUUCAUGCU U	460	AAGCAUGAAGACCUAUGA U	461
620	UCAUAGGUCUUCAUGCUU A	462	UAAGCAUGAAGACCUAUG A	463
621	CAUAGGUCUUCAUGCUUA U	464	AUAAGCAUGAAGACCUAU G	465
622	AUAGGUCUUCAUGCUUAU G	466	CAUAAGCAUGAAGACCUA U	467
623	UAGGUCUUCAUGCUUAUG G	468	CCAUAAGCAUGAAGACCU A	469
624	AGGUCUUCAUGCUUAUGG G	470	CCCAUAAGCAUGAAGACC U	471
625	GGUCUUCAUGCUUAUGGG G	472	CCCCAUAAGCAUGAAGAC C	473
626	GUCUUCAUGCUUAUGGGG A	474	UCCCCAUAAGCAUGAAGA C	475
627	UCUUCAUGCUUAUGGGGA U	476	AUCCCCAUAAGCAUGAAG A	477

## IDH2

Exemplary siRNAs are presented in the following tables. Art-known methods can be used to select other siRNAs. siRNAs can be evaluated, *e.g.*, by determining the ability of an siRNA to silence an *e.g.*, IDH2, *e.g.*, in an *in vitro* system, *e.g.*, in cultured cells, *e.g.*, HeLa cells or cultured glioma cells. *e.g.*,

The siRNAs in **Table 15** were generated using the siRNA selection tool available on the worldwide web at [jura.wi.mit.edu/bioc/siRNAext/](http://jura.wi.mit.edu/bioc/siRNAext/). (Yuan *et al.* Nucl. Acids. Res. 2004 32:W130-W134.) Other selection tools can be used as well. Entry 1356 was adapted from *Silencing of cytosolic NADP+ dependent isocitrate dehydrogenase by small interfering RNA enhances the sensitivity of HeLa cells toward staurosporine*, Lee *et al.*, 2009, Free Radical Research, 43: 165-173.

The siRNAs in Tables **16-23** represent candidates spanning the IDH2 mRNA at nucleotide positions 600, 601, and 602 according to the mRNA sequence presented at GenBank Accession No. NM\_002168.2 (Record dated August 16, 2009; GI28178831) (SEQ ID NO12, **FIG. 22B**; equivalent to nucleotide positions 514, 515, and 516 of the cDNA sequence represented by SEQ ID NO:11, **FIG. Fig. 22A**).

The RNAs in the tables can be modified, *e.g.*, as described herein. Modifications include chemical modifications to enhance properties, *e.g.*, resistance to degradation, or the use of overhangs. For example, either one or both of the sense and antisense strands in the tables can include an additional dinucleotide at the 3' end, *e.g.*, TT, UU, dTdT.

**Table 15. siRNAs targeting wildtype IDH2**

<b>Position on mRNA (FIG. 22B)</b>	<b>sense (5' to 3')</b>	<b>SEQ ID NO:</b>	<b>antisense (5' to 3')</b>	<b>SEQ ID NO:</b>
250	GUGAUGAGAUGACCCGUUU	478	AUACGGGUCAUCUCAUCAC	479
252	GAUGAGAUGACCCGUUUUA	480	UAAUACGGGUCAUCUCAUC	481
264	CGUAUUUAUCUGGCAGUUCA	482	UGAACUGCCAGAUAAUACG	483
274	GGCAGUUCAUCAAGGAGAA	484	UUCUCCUUGAUGAACUGCC	485
451	GUGUGGAAGAGUUCAAGCU	486	AGCUUGAACUCUCCACAC	487
453	GUGGAAGAGUUCAAGCUGA	488	UCAGCUUGAACUCUCCAC	489
456	GAAGAGUUCAAGCUGAAGA	490	UCUUCAGCUUGAACUCUUC	491
795	CAGUAUGCCAUCAGAGAAGA	492	UCUUCUGGAUGGCAUACUG	493
822	CUGUACAUGAGACCAAGA	494	UCUUGGUGCUCAUGUACAG	495
832	GCACCAAGAACACCAUACU	496	AGUAUGGUGUUCUUGGUGC	497
844	CCAUACUGAAAGCCUACGA	498	UCGUAGGCUUUCAGUAUGG	499
845	CAUACUGAAAGCCUACGAU	500	AUCGUAGGCUUUCAGUAUG	501
868	GUUUCAGGACAUCUCCA	502	UGGAAGAUGCCUUGAAAC	503
913	CCGACUUCGACAAGAAUAA	504	UUAUUCUUGUCGAAGUCGG	505
915	GACUUCGACAAGAAUAAGA	506	UCUUAUUCUUGUCGAAGUC	507
921	GACAAGAAUAAGAUCUGGU	508	ACCAGAUCUUAUUCUUGUC	509
949	GGCUCAUUGAUGACAUGGU	510	ACCAUGUCAUCAUGAGCC	511
1009	GCAAGAACUAUGACGGAGA	512	UCUCCGUCAUAGUUCUUGC	513
1010	CAAGAACUAUGACGGAGAU	514	AUCUCCGUCAUAGUUCUUG	515
1024	GAGAUGUGCAGUCAGACAU	516	AUGUCUGACUGCACAUUC	517
1096	CUGAUGGGAAGACGAUUGA	518	UCAAUCGUCUCCCAUCAG	519
1354	GCAAUGUGAAGCUGAACGA	520	UCGUUCAGCUUCACAUUGC	521
1668	CUGUAAUUUAUUUGCCCU	522	AGGGCAAUAUAAUUACAG	523
1694	CAUGGUGCCAUAUUUAGCU	524	AGCUAAUAUUGGCACCAUG	525
1697	GGUGCCAUAUUUAGCUACU	526	AGUAGCUAAUAUUGGCACC	527
1698	GUGCCAUAUUUAGCUACUA	528	UAGUAGCUAAUAUUGGCAC	529
1700	GCCAUAUUUAGCUACUAAA	530	UUUAGUAGCUAAUAUUGGC	531

**Table 16.** siRNAs targeting wildtype IDH2

<b>Position on mRNA (FIG. 22B)</b>	<b>sense (5' to 3')</b>	<b>SEQ ID NO:</b>	<b>antisense (5' to 3')</b>	<b>SEQ ID NO:</b>
584	GCCCAUCACCAUUGGCAGG	532	CCUGCCAAUGGUGAUGGGC	533
585	CCCAUCACCAUUGGCAGGC	534	GCCUGCCAAUGGUGAUGGG	535
586	CCAUCACCAUUGGCAGGCA	536	UGCCUGCCAAUGGUGAUGG	537
587	CAUCACCAUUGGCAGGCAC	538	GUGCCUGCCAAUGGUGAUG	539
588	AUCACCAUUGGCAGGCACG	540	CGUGCCUGCCAAUGGUGAU	541
589	UCACCAUUGGCAGGCACGC	542	GCGUGCCUGCCAAUGGUGA	543
590	CACCAUUGGCAGGCACGCC	544	GGCGUGCCUGCCAAUGGUG	545
591	ACCAUUGGCAGGCACGCC	546	GGGCGUGCCUGCCAAUGGU	547
592	CCAUUGGCAGGCACGCCCA	548	UGGGCGUGCCUGCCAAUGG	549
593	CAUUGGCAGGCACGCCCAU	550	AUGGGCGUGCCUGCCAAUG	551
594	AUUGGCAGGCACGCCCAUG	552	CAUGGGCGUGCCUGCCAAU	553
595	UUGGCAGGCACGCCCAUGG	554	CCAUGGGCGUGCCUGCCAA	555
596	UGGCAGGCACGCCCAUGGC	556	GCCAUGGGCGUGCCUGCCA	557
597	GGCAGGCACGCCCAUGGCG	558	CGCCAUGGGCGUGCCUGCC	559
598	GCAGGCACGCCCAUGGCGA	560	UCGCCAUGGGCGUGCCUGC	561
599	CAGGCACGCCCAUGGCGAC	562	GUCGCCAUGGGCGUGCCUG	563
600	AGGCACGCCCAUGGCGACC	564	GGUCGCCAUGGGCGUGCCU	565

**Table 17.** siRNAs targeting A514G mutant IDH2 (equivalent to A600G of SEQ ID

NO:12. (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCGGG	566	CCCGCCAAUGGUGAUGGGC	567
585	CCCAUCACCAUUGGCGGGC	568	GCCCGCCAAUGGUGAUGGG	569
586	CCAUCACCAUUGGCGGGCA	570	UGCCCGCCAAUGGUGAUGG	571
587	CAUCACCAUUGGCGGGCAC	572	GUGCCCGCCAAUGGUGAUG	573
588	AUCACCAUUGGCGGGCAGC	574	CGUGCCCGCCAAUGGUGAU	575
589	UCACCAUUGGCGGGCAGC	576	GCGUGCCCGCCAAUGGUGA	577
590	CACCAUUGGCGGGCAGGCC	578	GGCGUGCCCGCCAAUGGUG	579
591	ACCAUUGGCGGGCAGGCC	580	GGCGUGCCCGCCAAUGGU	581
592	CCAUUGGCGGGCAGGCCCA	582	UGGGCGUGCCCGCCAAUGG	583
593	CAUUGGCGGGCAGGCCCAU	584	AUGGGCGUGCCCGCCAAUG	585
594	AUUGGCGGGCAGGCCCAUG	586	CAUGGGCGUGCCCGCCAAU	587
595	UUGGCGGGCAGGCCCAUGG	588	CCAUGGGCGUGCCCGCCAA	589
596	UGGCGGGCAGGCCCAUGGC	590	GCCAUGGGCGUGCCCGCCA	591
597	GGCGGGCAGGCCCAUGGCG	592	CGCCAUGGGCGUGCCCGCC	593
598	GCGGGCAGGCCCAUGGCGA	594	UCGCCAUGGGCGUGCCCGC	595
599	CGGGCAGGCCCAUGGCGAC	596	GUCGCCAUGGGCGUGCCCG	597
600	GGGCAGGCCCAUGGCGACC	598	GGUCGCCAUGGGCGUGCCC	599



**Table 18.** siRNAs targeting A514U mutant IDH2 (equivalent to A600U of SEQ ID

NO:12, (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCUGG	600	CCAGCCAAUGGUGAUGGGC	601
585	CCCAUCACCAUUGGCUGGC	602	GCCAGCCAAUGGUGAUGGG	603
586	CCAUCACCAUUGGCUGGCA	604	UGCCAGCCAAUGGUGAUGG	605
587	CAUCACCAUUGGCUGGCAC	606	GUGCCAGCCAAUGGUGAUG	607
588	AUCACCAUUGGCUGGCACG	608	CGUGCCAGCCAAUGGUGAU	609
589	UCACCAUUGGCUGGCACGC	610	GCGUGCCAGCCAAUGGUGA	611
590	CACCAUUGGCUGGCACGCC	612	GGCGUGCCAGCCAAUGGUG	613
591	ACCAUUGGCUGGCACGCC	614	GGGCGUGCCAGCCAAUGGU	615
592	CCAUUGGCUGGCACGCCCA	616	UGGGCGUGCCAGCCAAUGG	617
593	CAUUGGCUGGCACGCCCAU	618	AUGGGCGUGCCAGCCAAUG	619
594	AUUGGCUGGCACGCCCAUG	620	CAUGGGCGUGCCAGCCAAU	621
595	UUGGCUGGCACGCCCAUGG	622	CCAUGGGCGUGCCAGCCAA	623
596	UGGCUGGCACGCCCAUGGC	624	GCCAUGGGCGUGCCAGCCA	625
597	GGCUGGCACGCCCAUGGCG	626	CGCCAUGGGCGUGCCAGCC	627
598	GCUGGCACGCCCAUGGCGA	628	UCGCCAUGGGCGUGCCAGC	629
599	CUGGCACGCCCAUGGCGAC	630	GUCGCCAUGGGCGUGCCAG	631
600	UGGCACGCCCAUGGCGACC	632	GGUCGCCAUGGGCGUGCCA	633

**Table 19.** siRNAs targeting G515A mutant IDH2 (equivalent to G601A of SEQ ID

NO:12, (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCAAG	634	CUUGCCAAUGGUGAUGGGC	635
585	CCCAUCACCAUUGGCAAGC	636	GCUUGCCAAUGGUGAUGGG	637
586	CCAUCACCAUUGGCAAGCA	638	UGCUUGCCAAUGGUGAUGG	639
587	CAUCACCAUUGGCAAGCAC	640	GUGCUUGCCAAUGGUGAUG	641
588	AUCACCAUUGGCAAGCAGC	642	CGUGCUUGCCAAUGGUGAU	643
589	UCACCAUUGGCAAGCACGC	644	GCGUGCUUGCCAAUGGUGA	645
590	CACCAUUGGCAAGCACGCC	646	GGCGUGCUUGCCAAUGGUG	647
591	ACCAUUGGCAAGCACGCC	648	GGGCGUGCUUGCCAAUGGU	649
592	CCAUUGGCAAGCACGCCCA	650	UGGGCGUGCUUGCCAAUGG	651
593	CAUUGGCAAGCACGCCCAU	652	AUGGGCGUGCUUGCCAAUG	653
594	AUUGGCAAGCACGCCCAUG	654	CAUGGGCGUGCUUGCCAAU	655
595	UUGGCAAGCACGCCCAUGG	656	CCAUGGGCGUGCUUGCCAA	657
596	UGGCAAGCACGCCCAUGGC	658	GCCAUGGGCGUGCUUGCCA	659
597	GGCAAGCACGCCCAUGGCG	660	CGCCAUGGGCGUGCUUGCC	661
598	GCAAGCACGCCCAUGGCGA	662	UCGCCAUGGGCGUGCUUGC	663
599	CAAGCACGCCCAUGGCGAC	664	GUCGCCAUGGGCGUGCUUG	665
600	AAGCACGCCCAUGGCGACC	666	GGUCGCCAUGGGCGUGCUU	667

**Table 20.** siRNAs targeting G515C mutant IDH2 (equivalent to G601C of SEQ ID

NO:12, (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCACG	668	CGUGCCAAUGGUGAUGGGC	669
585	CCCAUCACCAUUGGCACGC	670	GCGUGCCAAUGGUGAUGGG	671
586	CCAUCACCAUUGGCACGCA	672	UGCGUGCCAAUGGUGAUGG	673
587	CAUCACCAUUGGCACGCAC	674	GUGCGUGCCAAUGGUGAUG	675
588	AUCACCAUUGGCACGCACG	676	CGUGCGUGCCAAUGGUGAU	677
589	UCACCAUUGGCACGCACGC	678	GCGUGCGUGCCAAUGGUGA	679
590	CACCAUUGGCACGCACGCC	680	GGCGUGCGUGCCAAUGGUG	681
591	ACCAUUGGCACGCACGCCC	682	GGGCGUGCGUGCCAAUGGU	683
592	CCAUUGGCACGCACGCCCA	684	UGGGCGUGCGUGCCAAUGG	685
593	CAUUGGCACGCACGCCCAU	686	AUGGGCGUGCGUGCCAAUG	687
594	AUUGGCACGCACGCCCAUG	688	CAUGGGCGUGCGUGCCAAU	689
595	UUGGCACGCACGCCCAUGG	690	CCAUGGGCGUGCGUGCCAA	691
596	UGGCACGCACGCCCAUGGC	692	GCCAUGGGCGUGCGUGCCA	693
597	GGCACGCACGCCCAUGGCG	694	CGCCAUGGGCGUGCGUGCC	695
598	GCACGCACGCCCAUGGCGA	696	UCGCCAUGGGCGUGCGUGC	697
599	CACGCACGCCCAUGGCGAC	698	GUCGCCAUGGGCGUGCGUG	699
600	ACGCACGCCCAUGGCGACC	700	GGUCGCCAUGGGCGUGCGU	701

**Table 21.** siRNAs targeting G515U mutant IDH2 (equivalent to G601U of SEQ ID

NO:12, (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCAUG	702	CAUGCCAAUGGUGAUGGGC	703
585	CCCAUCACCAUUGGCAUGC	704	GCAUGCCAAUGGUGAUGGG	705
586	CCAUCACCAUUGGCAUGCA	706	UGCAUGCCAAUGGUGAUGG	707
587	CAUCACCAUUGGCAUGCAC	708	GUGCAUGCCAAUGGUGAUG	709
588	AUCACCAUUGGCAUGCACG	710	CGUGCAUGCCAAUGGUGAU	711
589	UCACCAUUGGCAUGCACGC	712	GCGUGCAUGCCAAUGGUGA	713
590	CACCAUUGGCAUGCACGCC	714	GGCGUGCAUGCCAAUGGUG	715
591	ACCAUUGGCAUGCACGCC	716	GGGCGUGCAUGCCAAUGGU	717
592	CCAUUGGCAUGCACGCCCA	718	UGGGCGUGCAUGCCAAUGG	719
593	CAUUGGCAUGCACGCCCAU	720	AUGGGCGUGCAUGCCAAUG	721
594	AUUGGCAUGCACGCCCAUG	722	CAUGGGCGUGCAUGCCAAU	723
595	UUGGCAUGCACGCCCAUGG	724	CCAUGGGCGUGCAUGCCAA	725
596	UGGCAUGCACGCCCAUGGC	726	GCCAUGGGCGUGCAUGCCA	727
597	GGCAUGCACGCCCAUGGCG	728	CGCCAUGGGCGUGCAUGCC	729
598	GCAUGCACGCCCAUGGCGA	730	UCGCCAUGGGCGUGCAUGC	731
599	CAUGCACGCCCAUGGCGAC	732	GUCGCCAUGGGCGUGCAUG	733
600	AUGCACGCCCAUGGCGACC	734	GGUCGCCAUGGGCGUGCAU	735

**Table 22.** siRNAs targeting G516C mutant IDH2 (equivalent to G602C of SEQ ID

NO:12, (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCAGC	736	GCUGCCAAUGGUGAUGGGC	737
585	CCCAUCACCAUUGGCAGCC	738	GGCUGCCAAUGGUGAUGGG	739
586	CCAUCACCAUUGGCAGCCA	740	UGGCUGCCAAUGGUGAUGG	741
587	CAUCACCAUUGGCAGCCAC	742	GUGGCUGCCAAUGGUGAUG	743
588	AUCACCAUUGGCAGCCACG	744	CGUGGCUGCCAAUGGUGAU	745
589	UCACCAUUGGCAGCCACGC	746	GCGUGGCUGCCAAUGGUGA	747
590	CACCAUUGGCAGCCACGCC	748	GGCGUGGCUGCCAAUGGUG	749
591	ACCAUUGGCAGCCACGCCCC	750	GGGCGUGGCUGCCAAUGGU	751
592	CCAUUGGCAGCCACGCCCA	752	UGGGCGUGGCUGCCAAUGG	753
593	CAUUGGCAGCCACGCCCAU	754	AUGGGCGUGGCUGCCAAUG	755
594	AUUGGCAGCCACGCCCAUG	756	CAUGGGCGUGGCUGCCAAU	757
595	UUGGCAGCCACGCCCAUGG	758	CCAUGGGCGUGGCUGCCAA	759
596	UGGCAGCCACGCCCAUGGC	760	GCCAUGGGCGUGGCUGCCA	761
597	GGCAGCCACGCCCAUGGCG	762	CGCCAUGGGCGUGGCUGCC	763
598	GCAGCCACGCCCAUGGCGA	764	UCGCCAUGGGCGUGGCUGC	765
599	CAGCCACGCCCAUGGCGAC	766	GUCGCCAUGGGCGUGGCUG	767
600	AGCCACGCCCAUGGCGACC	768	GGUCGCCAUGGGCGUGGCU	769

**Table 23.** siRNAs targeting G516U mutant IDH2 (equivalent to G602U of SEQ ID

NO:12, (FIG. 22B)

Position on mRNA (FIG. 22B)	sense (5' to 3')	SEQ ID NO:	antisense (5' to 3')	SEQ ID NO:
584	GCCCAUCACCAUUGGCAGU	770	ACUGCCAAUGGUGAUGGGC	771
585	CCCAUCACCAUUGGCAGUC	772	GACUGCCAAUGGUGAUGGG	773
586	CCAUCACCAUUGGCAGUCA	774	UGACUGCCAAUGGUGAUGG	775
587	CAUCACCAUUGGCAGUCAC	776	GUGACUGCCAAUGGUGAUG	777
588	AUCACCAUUGGCAGUCACG	778	CGUGACUGCCAAUGGUGAU	779
589	UCACCAUUGGCAGUCACGC	780	GCGUGACUGCCAAUGGUGA	781
590	CACCAUUGGCAGUCACGCC	782	GGCGUGACUGCCAAUGGUG	783
591	ACCAUUGGCAGUCACGCC	784	GGGCGUGACUGCCAAUGGU	785
592	CCAUUGGCAGUCACGCCCA	786	UGGGCGUGACUGCCAAUGG	787
593	CAUUGGCAGUCACGCCCAU	788	AUGGGCGUGACUGCCAAUG	789
594	AUUGGCAGUCACGCCCAUG	790	CAUGGGCGUGACUGCCAAU	791
595	UUGGCAGUCACGCCCAUGG	792	CCAUGGGCGUGACUGCCAA	793
596	UGGCAGUCACGCCCAUGGC	794	GCCAUGGGCGUGACUGCCA	795
597	GGCAGUCACGCCCAUGGCG	796	CGCCAUGGGCGUGACUGCC	797
598	GCAGUCACGCCCAUGGCGA	798	UCGCCAUGGGCGUGACUGC	799
599	CAGUCACGCCCAUGGCGAC	800	GUCGCCAUGGGCGUGACUG	801
600	AGUCACGCCCAUGGCGACC	802	GGUCGCCAUGGGCGUGACU	803

**EXAMPLE 6 STRUCTURAL ANALYSIS OF R132H MUTANT IDH1**

To define how R132 mutations alter the enzymatic properties of IDH1, the crystal structure of R132H mutant IDH1 bound to  $\alpha$ KG, NADPH, and  $\text{Ca}^{2+}$  was solved at 2.1 Å resolution.

The overall quaternary structure of the homodimeric R132H mutant enzyme adopts the same closed catalytically competent conformation (shown as a monomer in **FIG. 29A**) that has been previously described for the wild-type enzyme (Xu, X. et al. *J Biol Chem* 279, 33946-57 (2004)). NADPH is positioned as expected for hydride transfer to  $\alpha$ KG in an orientation that would produce R(-)-2HG, consistent with our chiral determination of the 2HG product.

Two important features were noted by the change of R132 to histidine: the effect on catalytic conformation equilibrium and the reorganization of the active-site. Locating atop a  $\beta$ -sheet in the relatively rigid small domain, R132 acts as a gate-keeper residue and appears to orchestrate the hinge movement between the open and closed conformations. The guanidinium moiety of R132 swings from the open to the closed conformation with a distance of nearly 8 Å. Substitution of histidine for arginine is likely to change the equilibrium in favor of the closed conformation that forms the catalytic cleft for cofactor and substrate to bind efficiently, which partly explains the high-affinity for NADPH exhibited by the R132H mutant enzyme. This feature may be advantageous for the NADPH-dependent reduction of  $\alpha$ KG to R(-)-2HG in an environment where NADPH concentrations are low. Secondly, closer examination of the catalytic pocket of the mutant IDH1 structure in comparison to the wild-type enzyme showed not only the expected loss of key salt-bridge interactions between the guanidinium of R132 and the  $\alpha/\beta$  carboxylates of isocitrate, as well as changes in the network that coordinates the metal ion, but also an unexpected reorganization of the active-site. Mutation to histidine resulted in a significant shift in position of the highly conserved residues Y139 from the A subunit and K212' from the B subunit (**FIG. 29B**), both of which are thought to be critical for catalysis of this enzyme family (Aktas, D. F. & Cook, P. F. *Biochemistry* 48, 3565-77 (2009)). In particular, the hydroxyl moiety of Y139 now occupies the space of the  $\beta$ -carboxylate of isocitrate. In addition, a significant repositioning of  $\alpha$ KG compared to isocitrate where the distal carboxylate of  $\alpha$ KG now points upward to make new contacts with N96 and S94 was observed. Overall, this single R132 mutation results in formation of a distinct active site compared to wild-type IDH1.

#### EXAMPLE 7 MATERIALS AND METHODS

##### *Summary*

R132H, R132C, R132L and R132S mutations were introduced into human IDH1 by standard molecular biology techniques. 293T and the human glioblastoma cell lines U87MG and LN-18 were cultured in DMEM, 10% fetal bovine serum. Cells were transfected and selected using standard techniques. Protein expression levels were determined by Western blot analysis using IDHc antibody (Santa Cruz Biotechnology), IDH1 antibody (proteintech), MYC tag antibody (Cell Signaling

Technology), and IDH2 antibody (Abcam). Metabolites were extracted from cultured cells and from tissue samples according to close variants of a previously reported method (Lu, W., Kimball, E. & Rabinowitz, J. D. *J Am Soc Mass Spectrom* 17, 37-50 (2006)), using 80% aqueous methanol (-80 °C) and either tissue scraping or homogenization to disrupt cells. Enzymatic activity in cell lysates was assessed by following a change in NADPH fluorescence over time in the presence of isocitrate and NADP, or  $\alpha$ KG and NADPH. For enzyme assays using recombinant IDH1 enzyme, proteins were produced in *E. coli* and purified using Ni affinity chromatography followed by Sephacryl S-200 size-exclusion chromatography. Enzymatic activity for recombinant IDH1 protein was assessed by following a change in NADPH UV absorbance at 340 nm using a stop-flow spectrophotometer in the presence of isocitrate and NADP or  $\alpha$ KG and NADPH. Chirality of 2HG was determined as described previously (Struys, E. A., Jansen, E. E., Verhoeven, N. M. & Jakobs, C. *Clin Chem* 50, 1391-5 (2004)). For crystallography studies, purified recombinant IDH1 (R132H) at 10 mg/mL in 20 mM Tris pH 7.4, 100 mM NaCl was pre-incubated for 60 min with 10 mM NADPH, 10 mM calcium chloride, and 75 mM  $\alpha$ KG. Crystals were obtained at 20°C by vapor diffusion equilibration using 3  $\mu$ L drops mixed 2:1 (protein:precipitant) against a well-solution of 100 mM MES pH 6.5, 20% PEG 6000. Patient tumor samples were obtained after informed consent as part of a UCLA IRB-approved research protocol. Brain tumor samples were obtained after surgical resection, snap frozen in isopentane cooled by liquid nitrogen and stored at -80 C. The IDH1 mutation status of each sample was determined using standard molecular biology techniques as described previously (Yan, H. et al. *N Engl J Med* 360, 765-73 (2009)). Metabolites were extracted and analyzed by LC-MS/MS as described above. Full methods are available in the supplementary material.

### ***Supplementary methods***

**Cloning, Expression, and Purification of ICDH1 wt and mutants in *E. coli*.** The open reading frame (ORF) clone of human isocitrate dehydrogenase 1 (cDNA) (IDH1; ref. ID NM\_005896) was purchased from Invitrogen in pENTR221 (Carlsbad, CA) and Origene Inc. in pCMV6 (Rockville, MD). To transfect cells with wild-type or mutant IDH1, standard molecular biology mutagenesis techniques were utilized to alter the DNA sequence at base pair 395 of the ORF in pCMV6 to introduce base pair change from guanine to adenine, which resulted in a change in the amino acid code at



position 132 from arginine (wt) to histidine (mutant; or R132H), and confirmed by standard DNA sequencing methods. For 293T cell transfection, wild-type and R132H mutant IDH1 were subcloned into pCMV-Sport6 with or without a carboxy-terminal Myc-DDK-tag. For stable cell line generation, constructs in pCMV6 were used. For expression in *E. coli*, the coding region was amplified from pENTR221 by PCR using primers designed to add NDEI and XHO1 restrictions sites at the 5' and 3' ends respectively. The resultant fragment was cloned into vector pET41a (EMD Biosciences, Madison, WI) to enable the *E. coli* expression of C-terminus His8-tagged protein. Site directed mutagenesis was performed on the pET41a-ICH1D1 plasmid using the QuikChange® MultiSite-Directed Mutagenesis Kit (Stratagene, La Jolla, CA) to change G395 to A, resulting in the Arg to His mutation. R132C, R132L and R132S mutants were introduced into pET41a-ICH1D1 in an analogous way.

Wild-type and mutant proteins were expressed in and purified from the *E. coli* Rosetta™ strain (Invitrogen, Carlsbad, CA) as follows. Cells were grown in LB (20 µg/ml Kanamycin) at 37°C with shaking until OD600 reaches 0.6. The temperature was changed to 18°C and protein expression was induced by adding IPTG to final concentration of 1 mM. After 12-16 hours of IPTG induction, cells were resuspended in Lysis Buffer (20mM Tris, pH7.4, 0.1% Triton X-100, 500 mM NaCl, 1 mM PMSF, 5 mM β-mercaptoethanol, 10 % glycerol) and disrupted by microfluidation. The 20,000g supernatant was loaded on metal chelate affinity resin (MCAC) equilibrated with Nickel Column Buffer A (20 mM Tris, pH7.4, 500mM NaCl, 5 mM β-mercaptoethanol, 10% glycerol) and washed for 20 column volumes. Elution from the column was effected by a 20 column-volume linear gradient of 10% to 100% Nickel Column Buffer B (20 mM Tris, pH7.4, 500 mM NaCl, 5 mM β-mercaptoethanol, 500 mM Imidazole, 10% glycerol) in Nickel Column Buffer A). Fractions containing the protein of interest were identified by SDS-PAGE, pooled, and dialyzed twice against a 200-volume excess of Gel Filtration Buffer (200 mM NaCl, 50 mM Tris 7.5, 5 mM β-mercaptoethanol, 2 mM MnSO<sub>4</sub>, 10% glycerol), then concentrated to 10 ml using Centricon (Millipore, Billerica, MA) centrifugal concentrators. Purification of active dimers was achieved by applying the concentrated eluent from the MCAC column to a Sephacryl S-200 (GE Life Sciences, Piscataway, NJ) column equilibrated with Gel Filtration Buffer and eluting the column with 20 column volumes of the same buffer. Fractions corresponding to the retention time of the dimeric protein were identified by SDS-PAGE and pooled for storage at -80°C.

**Cell lines and Cell Culture.** 293T cells were cultured in DMEM (Dulbecco's modified Eagles Medium) with 10% fetal bovine serum and were transfected using pCMV-6-based IDH-1 constructs in six-well plates with Fugene 6 (Roche) or Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions. Parental vector pCMV6 (no insert), pCMV6-wt IDH1 or pCMV6-R132H were transfected into human glioblastoma cell lines (U87MG; LN-18 (ATCC, HTB-14 and CRL-2610; respectively) cultured in DMEM with 10 % fetal bovine serum. Approximately 24 hrs after transfection, the cell cultures were transitioned to medium containing G418 sodium salt at concentrations of either 500 ug/ml (U87MG) or 750 ug/ml (LN-18) to select stable transfectants. Pooled populations of G418 resistant cells were generated and expression of either wild-type IDH1 or R132 IDH1 was confirmed by standard Western blot analysis.

**Western blot.** For transient transfection experiments in 293 cells, cells were lysed 72 hours after transfection with standard RIPA buffer. Lysates were separated by SDS-PAGE, transferred to nitrocellulose and probed with goat-anti-IDHc antibody (Santa Cruz Biotechnology sc49996) or rabbit-anti-MYC tag antibody (Cell Signaling Technology #2278) and then detected with HRP-conjugated donkey anti-goat or HRP-conjugated goat-anti-rabbit antibody (Santa Cruz Biotechnology sc2004). IDH1 antibody to confirm expression of both wild-type and R132H IDH1 was obtained from Proteintech. The IDH2 mouse monoclonal antibody used was obtained from Abcam.

**Detection of isocitrate,  $\alpha$ KG, and 2HG in purified enzyme reactions by LC-MS/MS.** Enzyme reactions performed as described in the text were run to completion as judged by measurement of the oxidation state of NADPH at 340 nm. Reactions were extracted with eight volumes of methanol, and centrifuged to remove precipitated protein. The supernatant was dried under a stream of nitrogen and resuspended in H<sub>2</sub>O. Analysis was conducted on an API2000 LC-MS/MS (Applied Biosystems, Foster City, CA). Sample separation and analysis was performed on a 150 x 2 mm, 4 uM Synergi Hydro-RP 80 A column, using a gradient of Buffer A (10

mM tributylamine, 15 mM acetic acid, 3% (v/v) methanol, in water) and Buffer B (methanol) using MRM transitions.

**Cell lysates based enzyme assays.** 293T cell lysates for measuring enzymatic activity were obtained 48 hours after transfection with M-PER lysis buffer supplemented with protease and phosphatase inhibitors. After lysates were sonicated and centrifuged at 12,000g, supernatants were collected and normalized for total protein concentration. To measure IDH oxidative activity, 3  $\mu$ g of lysate protein was added to 200  $\mu$ l of an assay solution containing 33 mM Tris-acetate buffer (pH 7.4), 1.3 mM  $MgCl_2$ , 0.33 mM EDTA, 100  $\mu$ M  $\beta$ -NADP, and varying concentrations of D-(+)-*threo*-isocitrate. Absorbance at 340 nm, reflecting NADPH production, was measured every 20 seconds for 30 min on a SpectraMax 190 spectrophotometer (Molecular Devices). Data points represent the mean activity of 3 replicates per lysate, averaged among 5 time points centered at every 5 min. To measure IDH reductive activity, 3  $\mu$ g of lysate protein was added to 200  $\mu$ l of an assay solution which contained 33 mM Tris-acetate (pH 7.4), 1.3 mM  $MgCl_2$ , 25  $\mu$ M  $\beta$ -NADPH, 40 mM  $NaHCO_3$ , and 0.6 mM  $\alpha$ KG. The decrease in 340 nm absorbance over time was measured to assess NADPH consumption, with 3 replicates per lysate.

**Recombinant IDH1 Enzyme Assays.** All reactions were performed in standard enzyme reaction buffer (150 mM NaCl, 20 mM Tris-Cl, pH 7.5, 10% glycerol, 5 mM  $MgCl_2$  and 0.03% (w/v) bovine serum albumin). For determination of kinetic parameters, sufficient enzyme was added to give a linear reaction for 1 to 5 seconds. Reaction progress was monitored by observation of the reduction state of the cofactor at 340 nm in an SFM-400 stopped-flow spectrophotometer (BioLogic, Knoxville, TN). Enzymatic constants were determined using curve fitting algorithms to standard kinetic models with the Sigmaplot software package (Systat Software, San Jose, CA).

**Determination of chirality of reaction products from enzyme reactions and tumors.** Enzyme reactions were run to completion and extracted with methanol as described above, then derivatized with enantiomerically pure tartaric acid before resolution and analysis by LC-MS/MS. After being thoroughly dried, samples were resuspended in freshly prepared 50 mg/ml (2*R*,3*R*)-(+)-Tartaric acid in dichloromethane:acetic acid (4:1) and incubated for 30 minutes at 75°C. After cooling

to room temperature, samples were briefly centrifuged at 14,000g, dried under a stream of nitrogen, and resuspended in H<sub>2</sub>O. Analysis was conducted on an API200 LC-MS/MS (Applied Biosystems, Foster City, CA), using an isocratic flow of 90:10 (2 mM ammonium formate, pH 3.6:MeOH) on a Luna C18(2) 150 x 2 mm, 5  $\mu$ M column. Tartaric-acid derivatized 2HG was detected using the 362.9/146.6 MRM transition and the following instrument settings: DP -1, FP -310, EP -4, CE-12, CXP-26. Analysis of the (R)-2HG standard, 2HG racemic mixture, and methanol-extracted tumor biomass (q.v.) was similarly performed.

**Crystallography conditions.** Crystals were obtained at 20°C by vapor diffusion equilibration using 3  $\mu$ L drops mixed 2:1 (protein:precipitant) against a well-solution of 100 mM MES pH 6.5, 20% PEG 6000.

**Protein characterization.** Approximately 90 mg of human cytosolic isocitrate dehydrogenase (HcIDH) was supplied to Xtal BioStructures by Agios. This protein was an engineered mutant form, R132S, with an 11-residue C-terminal affinity-purification tag (sequence SLEHHHHHHHH). The calculated monomeric molecular weight was 48.0 kDa and the theoretical pI was 6.50. The protein, at about 6 mg/mL concentration, was stored in 1-mL aliquots in 50 mM Tris-HCl (pH 7.4), 500 mM NaCl, 5 mM  $\beta$ -mercaptoethanol and 10% glycerol at -80°C. As shown in **FIG. 32A**, SDS-PAGE was performed to test protein purity and an anti-histidine Western blot was done to demonstrate the protein was indeed his-tagged. A sample of the protein was injected into an FPLC size-exclusion column to evaluate the sample purity and to determine the polymeric state in solution. **FIG. 32B** is a chromatogram of this run showing a single peak running at an estimated 87.6 kDa, suggesting IDH exists as a dimer at pH 7.4. Prior to crystallization, the protein was exchanged into 20 mM Tris-HCl (pH 7.4) and 100 mM NaCl using Amicon centrifugal concentrators. At this time, the protein was also concentrated to approximately 15 mg/mL. At this protein concentration and ionic strength, the protein tended to form a detectable level of precipitate. After spinning out the precipitate, the solution was stable at ~10 mg/mL at 4 °C.

**Initial attempts at crystallization.** The strategy for obtaining diffraction-quality crystals was derived from literature conditions, specifically “Structures of Human

Cytosolic NADP-dependent Isocitrate Dehydrogenase Reveal a Novel Self-regulatory Mechanism of Activity,” Xu, *et al.* (2005) *J.Biol.Chem.* 279: 33946-56. In this study, two crystal forms of HcIDH wildtype protein were produced. One contained their “binary complex”, IDH-NADP, which crystallized from hanging drops in the tetragonal space group  $P4_32_12$ . The drops were formed from equal parts of protein solution (15 mg/mL IDH, 10 mM NADP) and precipitant consisting of 100 mM MES (pH 6.5) and 12% PEG 20000. The other crystal form contained their “quaternary complex”, IDH-NADP/isocitrate/ $Ca^{2+}$ , which crystallized in the monoclinic space group  $P2_1$  using 100 mM MES (pH 5.9) and 20% PEG 6000 as the precipitant. Here they had added 10 mM DL-isocitrate and 10 mM calcium chloride to the protein solution. First attempts at crystallizing the R132S mutant in this study centered around these two reported conditions with little variation. The following lists the components of the crystallization that could be varied; several different combinations of these components were tried in the screening process.

In the protein solution:

HcIDH(R132S)	always ~10 mg/mL or ~0.2 mM
Tris-HCl (pH 7.4)	always 20 mM
NaCl	always 100 mM
NADP <sup>+</sup> /NADPH	absent or 5 mM NADP <sup>+</sup> (did not try NADPH)
DL-isocitric acid, trisodium salt	absent or 5 mM
calcium chloride	absent or 10 mM

In the precipitant: 100 mM MES (pH 6.5) and 12% PEG 20000

**OR**

100 mM MES (pH 6.0) and 20% PEG 6000

Drop size: always 3  $\mu$ L

Drop ratios: 2:1, 1:1 or 1:2 (protein:precipitant)

Upon forming the hanging drops, a milky precipitate was always observed. On inspection after 2-4 days at 20 °C most drops showed dense precipitation or phase separation. In some cases, the precipitate subsided and it was from these types of drops small crystals had grown, for example, as shown in **FIG. 33**.

**Crystal optimization.** Once bonafide crystals were achieved, the next step was to optimize the conditions to obtain larger and more regularly-shaped crystals of IDH-NADP/isocitrate/Ca<sup>2+</sup> in a timely and consistent manner. The optimal screen focused on varying the pH from 5.7 to 6.2, the MES concentration from 50 to 200 mM and the PEG 6000 concentration from 20 to 25%. Also, bigger drops were set up (5-6  $\mu$ l) and the drop ratios were again varied. These attempts failed to produce larger, diffraction-quality crystals but did reproduce the results reported above. Either a dense precipitate, oily phase separation or small crystals were observed.

**Using  $\alpha$ -Ketoglutarate.** Concurrent to the optimization of the isocitrate crystals, other screens were performed to obtain crystals of IDH(R132S) complexed with  $\alpha$ -ketoglutarate instead. The protein solution was consistently 10 mg/mL IDH in 20 mM Tris-HCl (pH 7.4) and 100 mM NaCl. The following were added in this order: 5 mM NADP, 5 mM  $\alpha$ -ketoglutaric acid (free acid, pH balanced with NaOH) and 10 mM calcium chloride. The protein was allowed to incubate with these compounds for at least an hour before the drops were set up. The precipitant was either 100 mM

MES (pH 6.5) and 12% PEG 20000 or 100 mM MES (pH 6.5) and 20% PEG 6000. Again, precipitation or phase separation was primarily seen, but in some drops small crystals did form. At the edge of one of the drops, a single large crystal formed, pictured below. This was the single crystal used in the following structure determination. **FIG. 34** shows crystal obtained from a protein solution contained 5 mM NADP, 5 mM  $\alpha$ -ketoglutarate, 10 mM Ca<sup>2+</sup>. Precipitant contained 100 mM MES (pH 6.5) and 12% PEG 20000.

**Cryo conditions.** In order to ship the crystal to the X-ray source and protect it during cryo-crystallography, a suitable cryo-protectant was needed. Glycerol is quite widely used and was the first choice. A cryo solution was made, basically as a mixture of the protein buffer and precipitant solution plus glycerol: 20 mM Tris-HCl (pH 7.5), 100 mM NaCl, 5 mM NADP, 5 mM  $\alpha$ -ketoglutaric acid, 10 mM calcium chloride, 100 mM MES (pH 6.5), 12% PEG 20000 and either 12.5% glycerol or 25% glycerol. The crystal was transferred to the cryo solution in two steps. First, 5  $\mu$ L of the 12.5% glycerol solution was added directly to the drop and incubated for 10 minutes, watching for possible cracking of the crystal. The liquid was removed from the drop and 10  $\mu$ L of the 25% glycerol solution was added on top of the crystal. Again, this incubated for 10 minutes, harvested into a nylon loop and plunged into liquid nitrogen. The crystal was stored submerged in a liquid nitrogen dewar for transport.

**Data collection and processing.** The frozen crystal was mounted on a Rigaku RAXIS IV X-ray instrument under a stream of nitrogen gas at temperatures near  $-170$  °C. A 200° dataset was collected with the image plate detector using 1.54 Å wavelength radiation from a rotating copper anode home source, 1° oscillations and 10 minute exposures. The presence of 25% glycerol as a cryoprotectant was sufficient for proper freezing, as no signs of crystal cracking (split spots or superimposed lattices) were observed. A diffuse ring was observed at 3.6 Å resolution, most likely caused by icing. The X-ray diffraction pattern showed clear lattice planes and reasonable spot separation, although the spacing along one reciprocal axis was rather small ( $b = 275.3$ ). The data was indexed to 2.7 Å resolution into space group  $P2_12_12$  with HKL2000 (Otwinowski and Minor, 1997). Three structures for HcIDH are known, designated the closed form (1T0L), the open form (1T09 subunit A) and semi-open form (1T09 subunit B). Molecular replacement was performed with the

CCP4 program PHASER (Bailey, 1994) using only the protein atoms from these three forms. Only the closed form yielded a successful molecular replacement result with 6 protein subunits in the asymmetric unit. The unit cell contains approximately 53.8% solvent.

**Model refinement.** Using the CCP4 program REFMAC5, rigid-body refinement was performed to fit each of the 6 IDH subunits in the asymmetric unit. This was followed by rigid-body refinement of the three domains in each protein subunit. Restrained refinement utilizing non-crystallographic symmetry averaging of related pairs of subunits yielded an initial structure with  $R_{cryst}$  of 33% and  $R_{free}$  of 42%. Model building and real-space refinement were performed using the graphics program COOT (Emsley and Cowtan, 2004). A difference map was calculated and this showed strong electron density into which six individual copies of the NADP ligand and calcium ion were manually fit with COOT. Density for the  $\alpha$ -ketoglutarate structure was less defined and was fit after the binding-site protein residues were fit using a  $2F_o - F_c$  composite omit map. Automated Ramachandran-plot optimization coupled with manual real-space density fitting was applied to improve the overall geometry and fit. A final round of restrained refinement with NCS yielded an  $R_{cryst}$  of 30.1% and  $R_{free}$  of 35.2%.

a, Å	b, Å	c, Å	$\alpha$	$\beta$	$\gamma$	Unit cell volume, Å <sup>3</sup>	Z
116.14	275.30	96.28	90°	90°	90°	$3.08 \times 10^6$	24

Reflections in working set / test set	68,755 / 3,608 (5.0%)
$R_{cryst}$	30.1%
$R_{free}$	35.2%



X-ray data and refinement statistics for IDH(R132S)-NADP/ $\alpha$ -ketoglurate/ $\text{Ca}^{2+}$ 

Crystal parameters	
Space group	<i>P2<sub>1</sub>2<sub>1</sub>2</i>
Unit cell dimensions	
a, b, c, Å	116.139, 275.297, 96.283
$\alpha$ , $\beta$ , $\gamma$ , °	90.0, 90.0, 90.0
Volume, Å <sup>3</sup>	3,078,440
No. protein molecules in asymmetric unit	6
No. protein molecules in unit cell, Z	24
Data collection	
Beam line	
Date of collection	Apr 25, 2009
$\lambda$ , Å	1.5418
Detector	Rigaku Raxis IV
Data set ( $\phi$ ), °	200
Resolution, Å	25-2.7 (2.8-2.7)
Unique reflections ( <i>N</i> , <i>F</i> > 0)	73,587
Completeness, %	85.4 (48.4)
$\langle I \rangle / \sigma I$	9.88 (1.83)

R-merge	0.109 (0.33)
Redundancy	4.3 (1.8)
Mosaicity	0.666
Wilson B factor	57.9
Anisotropy B factor, $\text{\AA}^2$	-1.96
Refinement Statistics	
Resolution limit, $\text{\AA}$	20.02-2.70
No. of reflections used for R-work <sup>a</sup> / R-free <sup>b</sup>	68,755 / 3608
Protein atoms	19788
Ligand atoms	348
No. of waters	357
Ions etc.	6
Matthews coeff. $\text{\AA}^3$ / Dalton	2.68
Solvent, %	53.8
R-work <sup>a</sup> / R-free <sup>b</sup> , (%)	30.1 / 35.2
Figure-of-merit <sup>c</sup>	0.80 (0.74)
Average B factors	31.0
Coordinates error (Luzzati plot), $\text{\AA}$	0.484
R.M.S. deviations	

Bond lengths, Å	0.026
Bond angles, °	2.86

Completeness and *R*-merge are given for all data and for data in the highest resolution shell. Highest shell values are in parentheses.

<sup>a</sup>*R* factor =  $\sum_{hkl} |F_o - F_c| / \sum_{hkl} F_o$ , where  $F_o$  and  $F_c$  are the observed and calculated structure factor amplitudes, respectively for all reflections *hkl* used in refinement.

<sup>b</sup>*R*-free is calculated for 5% of the data that were not used in refinement.

<sup>c</sup>Figure of merit =  $\sqrt{x^2 + y^2}$ , where  $x = (\sum_0^{2\pi} P(\alpha) \cos \alpha) / (\sum_0^{2\pi} P(\alpha))$ ,  $y = (\sum_0^{2\pi} P(\alpha) \sin \alpha) / (\sum_0^{2\pi} P(\alpha))$ , and the phase probability  $P(\alpha) = \exp(A \cos \alpha + B \sin \alpha + C \cos(2\alpha) + D \sin(2\alpha))$ , where *A*, *B*, *C*, and *D* are the Hendrickson-Lattman coefficients and  $\alpha$  is the phase.

#### Stereochemistry of IDH(R132S)-NADP/ $\alpha$ -ketoglurate/Ca<sup>2+</sup>

Ramachandran plot statistics	No. of amino acids	% of Residues
Residues in most favored regions [A, B, L]	1824	82.2
Residues in additional allowed regions [a, b, l, p]	341	15.4
Residues in generously allowed regions [-a, -b, -l, -p]	38	1.7
Residues in disallowed regions	17	0.8
Number of non-glycine and non-proline residues	2220	100
Number of end-residues (excl. Gly and Pro)	387	
Number of glycine residues	198	
Number of proline residues	72	
Total number of residues	2877	
Overall <G> -factor <sup>d</sup> score (> -1.0)	-0.65	

Generated by PROCHECK (Laskowski RA, MacArthur MW, Moss DS, Thornton JM (1993) J Appl Crystallogr 26:283-291.)

<sup>d</sup> G-factors for main-chain and side-chain dihedral angles, and main-chain covalent forces (bond lengths and bond angles). Values should be ideally -0.5 or above -1.0.

Radiation wavelength, Å	1.54
Resolution, Å (outer shell)	20-2.70 (2.80-2.70)
Unique reflections	73,587
Completeness (outer shell)	85.4% (48.4%)
Redundancy (outer shell)	4.3 (1.8)
$R_{\text{merge}}$ (outer shell)	10.9% (33%)
$\langle I \rangle / \langle \sigma(I) \rangle$ (outer shell)	9.88 (1.83)

**Clinical Specimens, metabolite extraction and analysis.** Human brain tumors were obtained during surgical resection, snap frozen in isopentane cooled by liquid nitrogen and stored at -80 C. Clinical classification of the tissue was performed using standard clinical pathology categorization and grading as established by the WHO. Genomic sequence analysis was deployed to identify brain tumor samples containing either wild-type isocitrate dehydrogenase (IDH1) or mutations altering amino acid 132. Genomic DNA was isolated from 50-100 mgs of brain tumor tissue using standard methods. A polymerase chain reaction on the isolated genomic DNA was used to amplify a 295 base pair fragment of the genomic DNA that contains both the intron and 2<sup>nd</sup> exon sequences of human IDH1 and mutation status assessed by standard molecular biology techniques.

Metabolite extraction was accomplished by adding a 10x volume (m/v ratio) of -80 °C methanol:water mix (80%:20%) to the brain tissue (approximately 100mgs) followed by 30 s homogenization at 4 C. These chilled, methanol extracted homogenized tissues were then centrifuged at 14,000 rpm for 30 minutes to sediment the cellular and tissue debris and the cleared tissue supernatants were transferred to a screw-cap freezer vial and stored at -80 °C. For analysis, a 2X volume of tributylamine (10 mM) acetic acid (10 mM) pH 5.5 was added to the samples and analyzed by LCMS as follows. Sample extracts were filtered using a Millex-FG 0.20 micron disk and 10 µL were injected onto a reverse-phase HPLC column (Synergi 150mm x 2 mm,

Phenomenex Inc.) and eluted using a linear gradient LCMS-grade methanol (50% with 10 mM tributylamine and 10 mM acetic acid) ramping to 80 % methanol:10 mM tributylamine: 10 mM acetic acid over 6 minutes at 200  $\mu$ L/min. Eluted metabolite ions were detected using a triple-quadrupole mass spectrometer, tuned to detect in negative mode with multiple-reaction-monitoring mode transition set (MRM's) according to the molecular weights and fragmentation patterns for 8 known central metabolites, including 2-hydroxyglutarate as described above. Data was processed using Analyst Software (Applied Biosystems, Inc.) and metabolite signal intensities were obtained by standard peak integration methods.




EXAMPLE 9 COMPOUNDS THAT INHIBIT IDH1 R132H



Assays were conducted in a volume of 76  $\mu$ l assay buffer (150 mM NaCl, 10 mM MgCl<sub>2</sub>, 20 mM Tris pH 7.5, 0.03% bovine serum albumin) as follows in a standard 384-well plate: To 25  $\mu$ l of substrate mix (8  $\mu$ M NADPH, 2 mM aKG), 1  $\mu$ l of test compound was added in DMSO. The plate was centrifuged briefly, and then 25  $\mu$ l of enzyme mix was added (0.2  $\mu$ g/ml ICDH1 R132H) followed by a brief centrifugation and shake at 100 RPM. The reaction was incubated for 50 minutes at room temperature, then 25  $\mu$ l of detection mix (30  $\mu$ M resazurin, 36  $\mu$ g/ml ) was added and the mixture further incubated for 5 minutes at room temperature. The conversion of resazurin to resorufin was detected by fluorescent spectroscopy at Ex544 Em590 c/o 590.

**Table 24a** shows the wild type vs mutant selectivity profile of 5 examples of IDH1R132H inhibitors. The IDH1wt assay was performed at 1x Km of NADPH as opposed to IDHR132H at 10x or 100x Km of NADPH. The second example showed no inhibition, even at 100  $\mu$ M. Also, the first example has IC<sub>50</sub>=5.74  $\mu$ M but is shifted significantly when assayed at 100x Km, indicating direct NADPH-competitive inhibitor. The selectivity between wild type vs mutant could be >20-fold.



**Table 24a**






STRUCTURE	LDHa IC50	LDHb IC50	ICDH IC50 ( $\mu$ M) @ 4 $\mu$ M (10x Km) NADPH	ICDH IC50 ( $\mu$ M) @ 40 $\mu$ M NADPH	IC50 Ratio (40/4)	IDH1wt IC50 @ 1x Km ( $\mu$ M)

	25.43	64.07	5.74	>100	17.42	<b>16.22</b>
	5.92	17.40	12.26	41.40	3.38	<b>NO inhibition</b>
	8.61	>100	12.79	14.70	1.15	<b>19.23</b>






	33.75	>100	14.98	19.17	1.28	<b>46.83</b>
	12.76	>100	23.80	33.16	1.39	<b>69.33</b>






Additional exemplary compounds that inhibit IDH1R132H are provided below in Table 24b.






Compound	No.
	<b>1</b>
	<b>2</b>






Compound	No.
	3
	4
	5
	6
	7













Compound	No.
	8
	9
	10
	11
	12






Compound	No.
	13
	14
	15
	16
	17






Compound	No.
	18
	19
	20
	21
	22

Compound	No.
	23
	24
	25
	26
	27






Compound	No.
	28
	29
	30
	31
	32






Compound	No.
	33
	34
	35
	36
	37






Compound	No.
	38
	39
	40
	41
	42






Compound	No.
	43
	44
	45
	46
	47













Compound	No.
	48
	49
	50
	51
	52






Compound	No.
	53
	54
	55
	56
	57






Compound	No.
	58
	59
	60
	61
	62

Compound	No.
	63
	64
	65
	66
	67






Compound	No.
	68
	69
	70
	71
	72

Compound	No.
	73
	74
	75
	76
	77

Compound	No.
	78
	79
	80
	81
	82

Compound	No.
	83
	84
	85
	86
	87



Compound	No.
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C4=CC=C(F)C=C4)CC3</chem>	88
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C4=CC=C(C)C=C4)CC3</chem>	89
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C4=CC=C(OC)C=C4)CC3</chem>	90
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C4=CC=C(OC)C=C4)CC3</chem>	91
 <chem>CCCC1=CC=C(N1S(=O)(=O)C2=CC=C(C=C2)C(=O)N3CCN(C4=CC=C(C=C4)CCN5CCCC5)CC3</chem>	92

EXAMPLE 10. The mutant enzyme IDH2-R172K has elevated NADPH reductive catalysis activity as compared to wildtype IDH2 enzyme.

NADPH reduction activity was measured for the enzymes IDH2-R172K, IDH2-wildtype, IDH1-R132H and IDH1-wildtype. The final reactant concentrations for each reaction were as follows: 20 mM Tris 7.5, 150 mM NaCl, 2 mM MnCl<sub>2</sub>, 10% glycerol, 0.03% BSA, enzyme (1-120 µg/mL), 1 mM NADPH, and 5 mM αKG (alpha ketoglutarate). The resulting specific activities (µmol/min/mg) are presented in the graph in FIG. 35. The results indicate that the mutant IDH2 has elevated reductive activity as compared to wildtype IDH2, even though both the mutant and wildtype IDH2 enzymes were able to make 2HG (2-hydroxyglutarate) at saturating levels of reactants αKG and NADPH.

EXAMPLE 11: 2-HG accumulates in AML with IDH1/2 mutations

Patients and clinical data

Peripheral blood and bone marrow were collected from AML patients at the time of diagnosis and at relapse, following REB approved informed consent. The cells were separated by ficol hypaque centrifugation, and stored at -150° C in 10% DMSO, 40% FCS and 50% alpha-MEM medium. Patient sera were stored at -80° C. Cytogenetics and molecular testing were performed in the diagnostic laboratory of the University Health Network (Toronto, Canada). A subgroup of patients (n=132) was given consistent initial treatment using a standard induction and consolidation chemotherapy regimen consisting of daunorubicin and cytarabine.

IDH1 and IDH2 Genotyping

DNA was extracted from leukemic cells and cell lines using the Qiagen Puregene kit (Valencia CA). For a subset of samples (n=96), RNA was extracted from leukemic cells using a Qiagen RNeasy kit, and reverse transcribed into cDNA for IDH1 and IDH2 genotyping. IDH1 and IDH2 genotype was determined at the Analytical Genetics Technology Centre at the University Health Network (Toronto, Canada) using a Sequenom MassARRAY™ platform (Sequenom, San Diego, CA). Positive results were confirmed by direct sequencing on an ABI PRISM 3130XL genetic analyzer (Applied Biosystems, Foster City, CA).

Cell lines

AML cell lines (OCI/AML-1, OCI/AML-2, OCI/AML-3, OCI/AML-4, OCI/AML-5, HL-60, MV-4-11, THP-1, K562, and KG1A) and 5637 cells were obtained from the laboratory of Mark Minden (Ontario Cancer Institute, Toronto, Canada). Primary AML cells were cultured in alpha-MEM media supplemented with 20% fetal bovine serum, and 10% 5637 cell conditioned media as previously described<sup>13</sup>. Growth curves were generated by counting viable cells as assessed by trypan blue exclusion on a Vi-CELL automated cell counter (Beckman Coulter, Fullerton, CA).

#### Expression / purification of IDH1 and IDH2 proteins

The human IDH1 cDNA (ref. ID NM\_005896) and IDH2 cDNA (ref. ID NM\_002168) were purchased from OriGene Technologies (Rockville, MD). For expression in *E. coli*, the coding region was amplified by PCR using primers designed to add NDEI and XHO1 restrictions sites at the 5' and 3' ends respectively. The resultant fragments for IDH1 (full length) and IDH2 (residues 40-452) were cloned into vector pET41a (EMD Biosciences, Madison, WI) to enable the *E. coli* expression of C-terminal His8-tagged protein. Site directed mutagenesis was performed on the pET41a-IDH1 and pET41a-IDH2 plasmid using the QuikChange® Lightning Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA) to change C394 to T in the IDH1 cDNA, resulting in the R132C mutation, and to change G515 to A in the IDH2 cDNA, resulting in the R172K mutation. Wild-type and mutant IDH1 proteins were expressed in and purified from the *E. coli* Rosetta™ (DE3) strain according to manufacturer's instructions (Invitrogen, Carlsbad, CA). Overexpression of IDH2 protein was accomplished by co-transfection of expression plasmids encoding respective IDH2 clones and pG-KJE8 expressing chaperone proteins.

#### IDH1/2 activity assays

Enzymatic activity was assessed by following the change in NADPH absorbance at 340 nm over time in an SFM-400 stopped-flow spectrophotometer (BioLogic, Knoxville, TN) in the presence of isocitrate and NADP<sup>+</sup> (forward reaction), or  $\alpha$ -KG and NADPH (reverse reaction). All reactions were performed in standard enzyme reaction buffer (150 mM NaCl, 20 mM Tris-Cl, pH 7.5, 10mM MgCl<sub>2</sub> and 0.03% (w/v) bovine serum albumin). For determination of kinetic parameters, sufficient enzyme was added to give a linear reaction for 1 to 5 seconds. Enzymatic binding constants were determined using curve fitting algorithms to standard kinetic models with the Sigmaplot software package (Systat Software, San Jose, CA). For determination of *k*<sub>cat</sub>, enzyme was incubated with 5X *K*<sub>m</sub> of substrate and cofactor;

consumption of NADPH or NADP was determined by a change in the OD<sub>340</sub> over time. In both cases an extinction coefficient of 6200 M<sup>-1</sup> cm<sup>-1</sup> was used for NADPH.

#### 2-HG and metabolite analysis

Metabolites were extracted from cultured cells, primary leukemic cells, and sera using 80% aqueous methanol (-80°C) as previously described. For cell extraction, frozen biopsies were thawed quickly at 37°C, and an aliquot of 2 million cells was spun down at 4 °C. The pellet was resuspended in -80°C 80% methanol. For serum extraction, 1 ml of serum was thawed quickly and mixed with 4 ml -80° C methanol. All extracts were spun at 13000 rpm at 4 °C to remove precipitate, dried at room temperature, and stored at -80° C until analysis by LC-MS. Metabolite levels (2-HG, α-KG, succinate, fumarate, and malate) were determined by ion paired reverse phase LC coupled to negative mode electrospray triple-quadropole MS using multiple reaction monitoring, and integrated elution peaks were compared with metabolite standard curves for absolute quantification as described.

#### Statistical analysis

Fisher's exact test was used to test for differences in categorical variables between IDH1/2 wt and IDH1/2 mutant patients. One way ANOVA followed by a student's t-test with correction for multiple comparisons was used to test for differences in IDH1 activity and metabolite concentrations. Differences with p<0.05 were considered significant.

### **Results**

In order to investigate the role of IDH1 R132 mutations in AML, leukemic cells obtained at initial presentation, from a series of 145 AML patients treated at the Princess Margaret Hospital with the aim of identifying mutant samples in our viable cell tissue bank were genotyped. Heterozygous IDH1 R132 mutations were found in 11 (8%) of these patients (Table 25). The spectrum of IDH1 mutations observed in AML appears to differ from that seen in CNS tumors. In the CNS, the majority of mutations (80-90%) are IDH1 R132H substitutions, whereas 5, 4, and 2 patients with IDH1 R132H, R132C, and R132G mutations, respectively (Table 25), were observed. In four cases, leukemic cells were also available from samples taken at the time of relapse. The IDH1 mutation was retained in 4/4 of these samples (Table 25). One of the patients harboring an IDH1 mutation had progressed to AML from an earlier

myelodysplastic syndrome (MDS). When cells from the prior MDS in this patient were analyzed, IDH1 was found to be wild-type. An additional 14 patients with MDS were genotyped, and all patients were found to be wild-type for IDH1, suggesting that IDH1 mutations are not a common feature of this disease. In samples from a subset of IDH1 mutant patients (n=8), reverse transcribed RNA was used for genotyping in order to assess the relative expression of mutant and wild-type alleles. Sequenom genotyping showed balanced allele peaks for these samples, indicating that both the wild-type and mutant genes are expressed. Ten established AML cell lines were also genotyped (OCI/AML-1, OCI/AML-2, OCI/AML-3, OCI/AML-4, OCI/AML-5, HL-60, MV-4-11, THP-1, K562, and KG1A) and none carried an IDH1 R132 mutation. Table 25: Identification of 13 AML patients bearing an IDH1 R132 or IDH2 R172 mutation\*

**Table 25**

Patient ID	Mutation	Amino acid change	FAB subtype	NPM1 and FLT3 status	Cytogenetic profile	Genotype at relapse	2-HG level (ng/2x10 <sup>6</sup> cells)
<b>IDH1 mutations</b>							
090108	G/A	R132H	M4	na	Normal	na	2090
090356	G/A	R132H	na	na	na	na	1529
0034	C/T	R132C	M5a	Normal	Normal	na	10285
0086	C/G	R132G	M2	Normal	Normal	na	10470
0488	C/T	R132C	M0	Normal	Normal	R132C	13822
8587	G/A	R132H	na	Normal	Normal	na	5742
8665	C/T	R132C	M1	na	Normal	na	7217
8741	G/A	R132H	M4	NPM1	Normal	R132H	6419
9544	C/G	R132G	na	na	Normal	R132G	4962
0174268	G/A	R132H	M1	NPM1	Normal	R132H	8464
090148	C/T	R132C	M1	na	46,-xx,-i(7)-(p10)-[20]	na	na
<b>IDH2 mutations</b>							
9382	G/A	R172K	M0	Normal	Normal	na	19247
0831	G/A	R172K	M1	Normal	Normal	na	15977

\* NPM1 denotes nucleophosmin 1, and FLT3 FMS-related tyrosine kinase 3. na indicates that some data was not available for some patients.

<u>Patient ID</u>	<u>Mutation</u>	<u>Amino acid change</u>	<u>FAB subtype</u>	<u>NPM1 and FLT3 status</u>	<u>Cytogenetic profile</u>	<u>Genotype at relapse</u>	<u>2-HG level (ng/2x10<sup>6</sup> cells)</u>
-------------------	-----------------	--------------------------	--------------------	-----------------------------	----------------------------	----------------------------	---

<b>IDH1 mutations</b>							
<u>090108</u>	<u>G/A</u>	<u>R132H</u>	<u>M4</u>	<u>na</u>	<u>Normal</u>	<u>na</u>	<u>2090</u>
<u>090356</u>	<u>G/A</u>	<u>R132H</u>	<u>na</u>	<u>na</u>	<u>na</u>	<u>na</u>	<u>1529</u>
<u>0034</u>	<u>C/T</u>	<u>R132C</u>	<u>M5a</u>	<u>Normal</u>	<u>Normal</u>	<u>na</u>	<u>10285</u>
<u>0086</u>	<u>C/G</u>	<u>R132G</u>	<u>M2</u>	<u>Normal</u>	<u>Normal</u>	<u>na</u>	<u>10470</u>
<u>0488</u>	<u>C/T</u>	<u>R132C</u>	<u>M0</u>	<u>Normal</u>	<u>Normal</u>	<u>R132C</u>	<u>13822</u>
<u>8587</u>	<u>G/A</u>	<u>R132H</u>	<u>na</u>	<u>Normal</u>	<u>Normal</u>	<u>na</u>	<u>5742</u>
<u>8665</u>	<u>C/T</u>	<u>R132C</u>	<u>M1</u>	<u>na</u>	<u>Normal</u>	<u>na</u>	<u>7217</u>
<u>8741</u>	<u>G/A</u>	<u>R132H</u>	<u>M4</u>	<u>NPM1</u>	<u>Normal</u>	<u>R132H</u>	<u>6419</u>
<u>9544</u>	<u>C/G</u>	<u>R132G</u>	<u>na</u>	<u>na</u>	<u>Normal</u>	<u>R132G</u>	<u>4962</u>
<u>0174268</u>	<u>G/A</u>	<u>R132H</u>	<u>M1</u>	<u>NPM1</u>	<u>Normal</u>	<u>R132H</u>	<u>8464</u>
<u>090148</u>	<u>C/T</u>	<u>R132C</u>	<u>M1</u>	<u>na</u>	<u>46. xx. i(7) (p10) [20]</u>	<u>na</u>	<u>na</u>
<b>IDH2 mutations</b>							
<u>9382</u>	<u>G/A</u>	<u>R172K</u>	<u>M0</u>	<u>Normal</u>	<u>Normal</u>	<u>na</u>	<u>19247</u>
<u>0831</u>	<u>G/A</u>	<u>R172K</u>	<u>M1</u>	<u>Normal</u>	<u>Normal</u>	<u>na</u>	<u>15877</u>

\* NPM1 denotes nucleophosmin 1, and FLT FMS-related tyrosine kinase 3. na indicates that some data was not available for some patients.

A metabolite screening assay to measure 2-HG in this set of AML samples was set up. Levels of 2-HG were approximately 50-fold higher in samples harboring an IDH1 R132 mutation (Table 25, Figure 36A, Table 26). 2-HG was also elevated in the sera of patients with IDH1 R132 mutant AML (Figure 36B). There was no relationship between the specific amino acid substitution at residue 132 of IDH1 and the level of 2-HG in this group of patients.

Table 26: Metabolite concentrations in individual IDH1/2 mutant and wild-type AML cells\*

Sample	IDH1/2 Genotype	2-HG (ng/2x10 <sup>6</sup> cells)	α-KG (ng/2x10 <sup>6</sup> cells)	Malate (ng/2x10 <sup>6</sup> cells)	Fumarate (ng/2x10 <sup>6</sup> cells)	Succinate (ng/2x10 <sup>6</sup> cells)
<u>0034</u>	<u>R132C</u>	<u>10295</u>	<u>125</u>	<u>102</u>	<u>239</u>	<u>2651</u>
<u>0086</u>	<u>R132G</u>	<u>10470</u>	<u>124</u>	<u>258</u>	<u>229</u>	<u>3043</u>
<u>0488</u>	<u>R132C</u>	<u>13822</u>	<u>95</u>	<u>184</u>	<u>199</u>	<u>2671</u>
<u>8587</u>	<u>R132H</u>	<u>5742</u>	<u>108</u>	<u>97</u>	<u>95</u>	<u>1409</u>
<u>8665</u>	<u>R132C</u>	<u>7217</u>	<u>137</u>	<u>118</u>	<u>120</u>	<u>1648</u>
<u>8741</u>	<u>R132H</u>	<u>6419</u>	<u>87</u>	<u>66</u>	<u>61</u>	<u>938</u>
<u>9544</u>	<u>R132G</u>	<u>4962</u>	<u>95</u>	<u>76</u>	<u>72</u>	<u>1199</u>
<u>0174268</u>	<u>R132H</u>	<u>8464</u>	<u>213</u>	<u>323</u>	<u>318</u>	<u>2287</u>
<u>090356</u>	<u>R132H</u>	<u>1529</u>	<u>138</u>	<u>657</u>	<u>366</u>	<u>1462</u>
<u>090108</u>	<u>R132H</u>	<u>2090</u>	<u>Na</u>	<u>246</u>	<u>941</u>	<u>3560</u>

<del>090148</del> †	R132C	na	Na	na	Na	Na
<del>8741</del> ‡	R132H	2890	131	113	106	1509
<del>9554</del> ‡	R132G	7448	115	208	227	2658
<del>0174268</del> ‡	R132H	964	72	134	138	2242
<del>0488</del> ‡	R132C	7511	85	289	310	3448
9382	R172K	19247	790	821	766	5481
0831	R172K	15877	350	721	708	5144
157	Wild type	212	121	484	437	3057
202	Wild type	121	57	161	136	1443
205	Wild type	147	39	162	153	1011
209	Wild type	124	111	167	168	1610
239	Wild type	112	106	305	361	1436
277	Wild type	157	61	257	257	2029
291	Wild type	113	118	124	128	1240
313	Wild type	116	75	161	181	1541
090158	Wild type	411	217	658	647	3202
090156	Wild type	407	500	1278	1275	6091

† IDH1/2 denotes isocitrate dehydrogenase 1 and 2, 2-HG 2-hydroxy glutarate, and  $\alpha$ -KG alpha-ketoglutarate.

Metabolite measurements were not available for all patients.

‡ metabolic measurements were not made due to limited patient sample

‡ indicates samples obtained at relapse.

Sample	IDH1/2 Genotype	2-HG (ng / 2x10 <sup>6</sup> cells)	$\alpha$ -KG (ng / 2x10 <sup>6</sup> cells)	Malate (ng / 2x10 <sup>6</sup> cells)	Fumarate (ng / 2x10 <sup>6</sup> cells)	Succinate (ng / 2x10 <sup>6</sup> cells)
0034	R132C	10285	125	192	239	2651
0086	R132G	10470	124	258	229	3043
0488	R132C	13822	95	184	193	2671
8587	R132H	5742	108	97	95	1409
8665	R132C	7217	137	118	120	1648
8741	R132H	6419	87	66	61	938
9544	R132G	4962	95	76	72	1199
0174268	R132H	8464	213	323	318	2287
090356	R132H	1529	138	657	366	1462
090108	R132H	2090	Na	246	941	3560
090148†	R132C	na	Na	na	Na	Na
8741‡	R132H	2890	131	113	106	1509
9554‡	R132G	7448	115	208	227	2658
0174268‡	R132H	964	72	134	138	2242
0488‡	R132C	7511	85	289	310	3448
9382	R172K	19247	790	821	766	5481
0831	R172K	15877	350	721	708	5144
157	Wild type	212	121	484	437	3057
202	Wild type	121	57	161	136	1443
205	Wild type	147	39	162	153	1011

<u>209</u>	Wild type	<u>124</u>	<u>111</u>	<u>167</u>	<u>168</u>	<u>1610</u>
<u>239</u>	Wild type	<u>112</u>	<u>106</u>	<u>305</u>	<u>361</u>	<u>1436</u>
<u>277</u>	Wild type	<u>157</u>	<u>61</u>	<u>257</u>	<u>257</u>	<u>2029</u>
<u>291</u>	Wild type	<u>113</u>	<u>118</u>	<u>124</u>	<u>128</u>	<u>1240</u>
<u>313</u>	Wild type	<u>116</u>	<u>75</u>	<u>151</u>	<u>181</u>	<u>1541</u>
<u>090158</u>	Wild type	<u>411</u>	<u>217</u>	<u>658</u>	<u>647</u>	<u>3202</u>
<u>090156</u>	Wild type	<u>407</u>	<u>500</u>	<u>1276</u>	<u>1275</u>	<u>6091</u>

\* IDH1/2 denotes isocitrate dehydrogenase 1 and 2, 2-HG 2-hydroxy glutarate, and  $\alpha$ -KG alpha-ketoglutarate. Metabolite measurements were not available for all patients.

† metabolic measurements were not made due to limited patient sample

‡ indicates samples obtained at relapse.

Two samples harboring wild-type IDH1 also showed high levels of 2-HG (Table 25). The high 2-HG concentration prompted sequencing of the IDH2 gene in these two AML samples, which established the presence of IDH2 R172K mutations in both samples (Table 25).

Evaluation of the clinical characteristics of patients with or without IDH1/2 mutations revealed a significant correlation between IDH1/2 mutations and normal karyotype ( $p=0.05$ ), but no other differences between these two groups (Table 27). Notably, there was no difference in treatment response for a subgroup of patients who received consistent treatment ( $n=136$ ). These findings are consistent with the initial report identifying IDH1 mutations in AML.

Table 27: Characteristics of IDH1/2 mutant and wild-type patients\*

Variable	IDH1/2 Wild-type (N=132)	IDH1/2 Mutant (N=13)	P-Value
Age (yr)	58.8 ±16.2	52.6 ±7.0	0.17†
Sex (% male)	63 (70/132)	62 (8/13)	0.77‡
WBC at diagnosis ( $10^9$ cells/L)	40.7 ±50.6	28.7 ±34.1	0.38†
Initial treatment response (% complete remission)	70 (85/122)	62 (8/13)	0.54‡
Cytogenetic profile (% normal)	62 (72/117)	92 (11/12)	0.05‡
<b>Additional mutations</b>			
— FLT3 (%)	17 (8/47)	0 (0/8)	0.58‡
— NPM1 (%)	30 (14/47)	25 (2/8)	1.0‡

\* For plus-minus values, the value indicates the mean, and  $\pm$  indicates the standard deviation. IDH1/2 denotes isocitrate dehydrogenase 1 and 2, WBC white blood cell count, FLT3 FMS-related tyrosine kinase 3, and NPM1 nucleophosmin 1.

† P-value was calculated using the student's t test.



‡ P-value was calculated using Fisher's exact test.

<u>Variable</u>	<u>IDH1/2 Wild-type (N=132)</u>	<u>IDH1/2 Mutant (N=13)</u>	<u>P Value</u>
<u>Age (yr)</u>	<u>58.8 ±16.2</u>	<u>52.6 ±7.0</u>	<u>0.17†</u>
<u>Sex (% male)</u>	<u>53 (70/132)</u>	<u>62 (8/13)</u>	<u>0.77‡</u>
<u>WBC at diagnosis (10<sup>9</sup> cells/L)</u>	<u>40.7 ±50.6</u>	<u>28.7 ±34.1</u>	<u>0.38†</u>
<u>Initial treatment response (% complete remission)</u>	<u>70 (85/122)</u>	<u>62 (8/13)</u>	<u>0.54‡</u>
<u>Cytogenetic profile (% normal)</u>	<u>62 (72/117)</u>	<u>92 (11/12)</u>	<u>0.05‡</u>
<u>Additional mutations</u>			
— <u>FLT3 (%)</u>	<u>17 (8/47)</u>	<u>0 (0/8)</u>	<u>0.58‡</u>
— <u>NPM1 (%)</u>	<u>30 (14/47)</u>	<u>25(2/8)</u>	<u>1.0‡</u>

\* For plus-minus values, the value indicates the mean, and ± indicates the standard deviation. IDH1/2 denotes isocitrate dehydrogenase 1 and 2. WBC white blood cell count. FLT3 FMS-related tyrosine kinase 3, and NPM1 nucleophosmin 1.

† P-value was calculated using the student's t-test.

‡ P-value was calculated using Fisher's exact test.

Panels of AML cells from wild-type and IDH1 mutant patients were cultured *in vitro*. There was no difference in the growth rates or viability of the IDH1 R132 mutant and wild-type cells, with both groups showing high variability in their ability to proliferate in culture, as is characteristic of primary AML cells (Figure 36C). There was no relationship between 2-HG levels in the IDH1 R132 mutant cells and their growth rate or viability in culture. After 14 days in culture, the mutant AML cells retained their IDH1 R132 mutations (11/11), and continued to accumulate high levels of 2-HG (Figure 36A), further confirming that IDH1 R132 mutations lead to the production and accumulation of 2-HG in AML cells.

To investigate the effect of IDH1/2 mutations on the concentration of cellular metabolites proximal to the IDH reaction,  $\alpha$ -KG, succinate, malate, and fumarate levels were measured in AML cells with IDH1/2 mutations and in a set of wild-type AML cells matched for AML subtype and cytogenetic profile. None of the metabolites were found to be greatly altered in the IDH1 mutants compared to the IDH1 wild-type cells (Figure 27, Supplementary Table 26). The mean level of  $\alpha$ -KG was not altered in the IDH1/2 mutant AML cells, suggesting that the mutation does not decrease the concentration of this metabolite as has been previously hypothesized.

To confirm that the R132C mutation of IDH1, and the R172K mutation of IDH2 confer a novel enzymatic activity that produces 2-HG, recombinant mutant enzymes were assayed for the NADPH-dependent reduction of  $\alpha$ -KG. When samples were analyzed by LC-MS upon completion of the enzyme assay, 2-HG was identified as the end product for both the IDH1 R132C and IDH2 R172K mutant enzymes (Figure 38). No isocitrate was detectable by LC-MS, indicating that 2-HG is the sole product of this reaction (Figure 38). This observation held true even when the reductive reaction was performed in buffer containing  $\text{NaHCO}_3$  saturated with  $\text{CO}_2$ .

A large proportion of IDH1 mutant patients in AML have an IDH1 R132C mutation (Table 25). In order to biochemically characterize mutant IDH1 R132C, the enzymatic properties of recombinant R132C protein were assessed *in vitro*. Kinetic analyses showed that the R132C substitution severely impairs the oxidative decarboxylation of isocitrate to  $\alpha$ -KG, with a significant decrease in  $k_{\text{cat}}$ , even though the affinity for the co-factor  $\text{NADP}^+$  remains essentially unchanged (Table 28). However, unlike the R132H mutant enzyme described previously the R132C mutation leads to a dramatic loss of affinity for isocitrate ( $K_M$ ), and a drop in net isocitrate metabolism efficiency ( $k_{\text{cat}}/K_M$ ) of more than six orders of magnitude (Table 28). This suggests a potential difference in the substrate-level regulation of enzyme activity in the context of AML. While substitution of cysteine at R132 inactivates the canonical conversion of isocitrate to  $\alpha$ -KG, the IDH1 R132C mutant enzyme acquires the ability to catalyze the reduction of  $\alpha$ -KG to 2-HG in an NADPH dependent manner (Figure 39). This reductive reaction of mutant IDH1 R132C is highly efficient ( $k_{\text{cat}}/K_M$ ) compared to the wild-type enzyme, due to the considerable increase in binding affinity of both the NADPH and  $\alpha$ -KG substrates ( $K_M$ ) (Table 28).

Table 28: Kinetic parameters of the IDH1 R132C mutant enzyme

Oxidative ( $\rightarrow$ NADPH)	WT	R132C
$K_{M,NADP^+}$ ( $\mu\text{M}$ )	49	21
$K_{M,\text{isocitrate}}$ ( $\mu\text{M}$ )	57	$8.7 \times 10^4$
$K_{M,MgCl_2}$ ( $\mu\text{M}$ )	29	$4.5 \times 10^2$
$K_{\alpha\text{KG}}$ ( $\mu\text{M}$ )	$6.1 \times 10^2$	61
$k_{\text{cat}}$ ( $\text{s}^{-1}$ )	$1.3 \times 10^5$	$7.1 \times 10^2$

$k_{cat}/K_{M, isoc}(M^{-1}s^{-1})$	$2.3 \times 10^9$	$8.2 \times 10^3$
<b>Reductive (<math>\rightarrow</math> NADP<sup>+</sup>)</b>	<b>WT</b>	<b>R132C</b>
$K_{M, NADPH}(\mu M)$	n/a*	0.3
$K_{M, aKG}(\mu M)$	n/a	295
$k_{cat}(s^{-1})$	~7 (est.)	$5.5 \times 10^2$

\* n/a indicates no measureable activity

<b>Oxidative (<math>\rightarrow</math> NADPH)</b>	<b>WT</b>	<b>R132C</b>
$K_{M, NADP+}(\mu M)$	49	21
$K_{M, isocitrate}(\mu M)$	57	$8.7 \times 10^4$
$K_{M, MgCl_2}(\mu M)$	29	$4.5 \times 10^2$
$K_{i, aKG}(\mu M)$	$6.1 \times 10^2$	61
$k_{cat}(s^{-1})$	$1.3 \times 10^5$	$7.1 \times 10^2$
$k_{cat}/K_{M, isoc}(M^{-1}s^{-1})$	$2.3 \times 10^9$	$8.2 \times 10^3$
<b>Reductive (<math>\rightarrow</math> NADP<sup>+</sup>)</b>	<b>WT</b>	<b>R132C</b>
$K_{M, NADPH}(\mu M)$	n/a*	0.3
$K_{M, aKG}(\mu M)$	n/a	295
$k_{cat}(s^{-1})$	~7 (est.)	$5.5 \times 10^2$

\* n/a indicates no measureable activity



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 4 columns: APPLICATION NUMBER (15/589,615), FILING OR 371(C) DATE (05/08/2017), FIRST NAMED APPLICANT (Lenny Dang), ATTY. DOCKET NO./TITLE (AGS-013C2)

CONFIRMATION NO. 7017

FORMALITIES LETTER

148106
GOODWIN PROCTER LLP
PATENT ADMINISTRATOR
100 Northern Avenue
BOSTON, MA 02210



Date Mailed: 07/28/2017

NOTICE OF INCOMPLETE REPLY (NONPROVISIONAL)

Filing Date Granted

The U.S. Patent and Trademark Office has received your reply on 07/20/2017 to the Notice to File Missing Parts (Notice) mailed 05/24/2017 and it has been entered into the nonprovisional application. The reply, however, does not include the following items required in the Notice. A complete reply must be timely filed to prevent ABANDONMENT of the above-identified application. Replies should be mailed to: Mail Stop Missing Parts, Commissioner for Patents, P.O. Box 1450, Alexandria VA 22313-1450.

Applicant is given TWO MONTHS from the date of the Notice to File Missing Parts (Notice) mailed 05/24/2017 within which to file all required items and pay any fees required below to avoid abandonment. Extensions of time may be obtained by filing a petition accompanied by the extension fee under the provisions of 37 CFR 1.136(a).

Items Required to Avoid Abandonment:

The required items noted below SHOULD be filed along with any items required above. The filing date of this nonprovisional application will be the date of receipt of the items required above.

The application is informal since it does not comply with the regulations for the reason(s) indicated below.

The required item(s) identified below must be timely submitted to avoid abandonment:

- A substitute specification in compliance with 37 CFR 1.52, 1.121(b)(3), and 1.125, is required. The substitute specification must be submitted with markings and be accompanied by a clean version (without markings) as set forth in 37 CFR 1.125(c) and a statement that the substitute specification contains no new matter (see 37 CFR 1.125(b)). The specification, claims, and/or abstract page(s) submitted is not acceptable and cannot be scanned or properly stored because:
- The application papers (including any electronically submitted papers) are not in compliance with 37 CFR 1.52 because pages 179-181,183 contain text that is unreadable or of insufficient clarity. Application papers (including any electronically submitted papers) must be presented in a form having sufficient clarity and contrast between the background of the paper and the writing thereon to permit the Office to electronically reproduce the papers by use of digital imaging and optical character recognition. See 37 CFR 1.52(a)(1)(v).

Applicant is cautioned that correction of the above items may cause the specification and drawings page count to exceed 100 pages. If the specification and drawings exceed 100 pages, applicant will need to submit the required application size fee.

Items Required To Avoid Processing Delays:

Applicant is notified that the above-identified application contains the deficiencies noted below. No period for reply is set forth in this notice for correction of these deficiencies. However, if a deficiency relates to the inventor's oath or declaration, the applicant must file an oath or declaration in compliance with 37 CFR 1.63, or a substitute statement in compliance with 37 CFR 1.64, executed by or with respect to each actual inventor no later than the expiration of the time period set in the "Notice of Allowability" to avoid abandonment. See 37 CFR 1.53(f).

- A properly executed inventor's oath or declaration has not been received for the following inventor(s):  
Lenny Dang

Replies must be received in the USPTO within the set time period or must include a proper Certificate of Mailing or Transmission under 37 CFR 1.8 with a mailing or transmission date within the set time period. For more information and a suggested format, see Form PTO/SB/92 and MPEP 512.

Replies should be mailed to:

Mail Stop Missing Parts  
Commissioner for Patents  
P.O. Box 1450  
Alexandria VA 22313-1450

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web, including a copy of this Notice and selecting the document description "Applicant response to Pre-Exam Formalities Notice".  
<https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at 1-866-217-9197 or visit our website at <http://www.uspto.gov/ebc>.

If you are not using EFS-Web to submit your reply, you must include a copy of this notice.

Questions about the contents of this notice and the requirements it sets forth should be directed to the Office of Data Management, Application Assistance Unit, at (571) 272-4000 or (571) 272-4200 or 1-888-786-0101.

/sareebuddin/  
.....

## Electronic Patent Application Fee Transmittal

<b>Application Number:</b>	15589615			
<b>Filing Date:</b>	08-May-2017			
<b>Title of Invention:</b>	METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS			
<b>First Named Inventor/Applicant Name:</b>	Lenny Dang			
<b>Filer:</b>	Ioana Davies/Jeannie Le			
<b>Attorney Docket Number:</b>	AGS-013C2			
Filed as Large Entity				
<b>Filing Fees for Utility under 35 USC 111(a)</b>				
<b>Description</b>	<b>Fee Code</b>	<b>Quantity</b>	<b>Amount</b>	<b>Sub-Total in USD(\$)</b>
<b>Basic Filing:</b>				
<b>Pages:</b>				
<b>Claims:</b>				
<b>Miscellaneous-Filing:</b>				
<b>Petition:</b>				
<b>Patent-Appeals-and-Interference:</b>				
<b>Post-Allowance-and-Post-Issuance:</b>				
<b>Extension-of-Time:</b>				

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Extension - 1 month with \$0 paid	1251	1	200	200
<b>Miscellaneous:</b>				
<b>Total in USD (\$)</b>				<b>200</b>

## Electronic Acknowledgement Receipt

<b>EFS ID:</b>	30000763
<b>Application Number:</b>	15589615
<b>International Application Number:</b>	
<b>Confirmation Number:</b>	7017
<b>Title of Invention:</b>	METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS
<b>First Named Inventor/Applicant Name:</b>	Lenny Dang
<b>Customer Number:</b>	148106
<b>Filer:</b>	Ioana Davies
<b>Filer Authorized By:</b>	
<b>Attorney Docket Number:</b>	AGS-013C2
<b>Receipt Date:</b>	07-AUG-2017
<b>Filing Date:</b>	08-MAY-2017
<b>Time Stamp:</b>	14:58:07
<b>Application Type:</b>	Utility under 35 USC 111(a)

### Payment information:

Submitted with Payment	yes
Payment Type	CARD
Payment was successfully received in RAM	\$200
RAM confirmation Number	080817INTEFSW14585000
Deposit Account	
Authorized User	

The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows:



<b>File Listing:</b>					
<b>Document Number</b>	<b>Document Description</b>	<b>File Name</b>	<b>File Size(Bytes)/ Message Digest</b>	<b>Multi Part /.zip</b>	<b>Pages (if appl.)</b>
1	Applicant Response to Pre-Exam Formalities Notice	Response_to_Notice_to_File_Missing_Parts_of_Application.pdf	17774 a3f3d64354ef47a9140bdeca463e135ae892347f	no	1
<b>Warnings:</b>					
<b>Information:</b>					
2	Extension of Time	One_Month_Request_for_Extension_of_Time_Under_37_CFR_1136a.pdf	23035 f0db0acaebb6068853ae8d85ce593bba45559663	no	1
<b>Warnings:</b>					
<b>Information:</b>					
3	Specification	Substitute_Specification_clean.pdf	1310112 0f18bdc2b0129837a8c5904b4da320145985a801	no	183
<b>Warnings:</b>					
<b>Information:</b>					
4	Specification	Substitute_Specification_marked.pdf	1333678 7836afb6bd651d39af7b001aacfd70f61d6b338d	no	185
<b>Warnings:</b>					
<b>Information:</b>					
5	Applicant Response to Pre-Exam Formalities Notice	Copy_of_Notice.pdf	246829 1f900a3e2942a0fe5a7edecaa702919bb788d11a	no	2
<b>Warnings:</b>					
<b>Information:</b>					
6	Fee Worksheet (SB06)	fee-info.pdf	30570 13ceb03396521d750666ef1f472ab0ad82d9b8e8	no	2
<b>Warnings:</b>					
<b>Information:</b>					

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

**New Applications Under 35 U.S.C. 111**

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

**National Stage of an International Application under 35 U.S.C. 371**

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

**New International Application Filed with the USPTO as a Receiving Office**

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 4 columns: APPLICATION NUMBER (15/589,615), FILING OR 371(C) DATE (05/08/2017), FIRST NAMED APPLICANT (Lenny Dang), ATTY. DOCKET NO./TITLE (AGS-013C2)

CONFIRMATION NO. 7017

FORMALITIES LETTER



148106
GOODWIN PROCTER LLP
PATENT ADMINISTRATOR
100 Northern Avenue
BOSTON, MA 02210

Date Mailed: 07/28/2017

NOTICE OF INCOMPLETE REPLY (NONPROVISIONAL)

Filing Date Granted

The U.S. Patent and Trademark Office has received your reply on 07/20/2017 to the Notice to File Missing Parts (Notice) mailed 05/24/2017 and it has been entered into the nonprovisional application. The reply, however, does not include the following items required in the Notice. A complete reply must be timely filed to prevent ABANDONMENT of the above-identified application. Replies should be mailed to: Mail Stop Missing Parts, Commissioner for Patents, P.O. Box 1450, Alexandria VA 22313-1450.

Applicant is given TWO MONTHS from the date of the Notice to File Missing Parts (Notice) mailed 05/24/2017 within which to file all required items and pay any fees required below to avoid abandonment. Extensions of time may be obtained by filing a petition accompanied by the extension fee under the provisions of 37 CFR 1.136(a).

Items Required to Avoid Abandonment:

The required items noted below SHOULD be filed along with any items required above. The filing date of this nonprovisional application will be the date of receipt of the items required above.

The application is informal since it does not comply with the regulations for the reason(s) indicated below.

The required item(s) identified below must be timely submitted to avoid abandonment:

- A substitute specification in compliance with 37 CFR 1.52, 1.121(b)(3), and 1.125, is required. The substitute specification must be submitted with markings and be accompanied by a clean version (without markings) as set forth in 37 CFR 1.125(c) and a statement that the substitute specification contains no new matter (see 37 CFR 1.125(b)). The specification, claims, and/or abstract page(s) submitted is not acceptable and cannot be scanned or properly stored because:
• The application papers (including any electronically submitted papers) are not in compliance with 37 CFR 1.52 because pages 179-181,183 contain text that is unreadable or of insufficient clarity. Application papers (including any electronically submitted papers) must be presented in a form having sufficient clarity and contrast between the background of the paper and the writing thereon to permit the Office to electronically reproduce the papers by use of digital imaging and optical character recognition. See 37 CFR 1.52(a)(1)(v).

Applicant is cautioned that correction of the above items may cause the specification and drawings page count to exceed 100 pages. If the specification and drawings exceed 100 pages, applicant will need to submit the required application size fee.

Items Required To Avoid Processing Delays:

Applicant is notified that the above-identified application contains the deficiencies noted below. No period for reply is set forth in this notice for correction of these deficiencies. However, if a deficiency relates to the inventor's oath or declaration, the applicant must file an oath or declaration in compliance with 37 CFR 1.63, or a substitute statement in compliance with 37 CFR 1.64, executed by or with respect to each actual inventor no later than the expiration of the time period set in the "Notice of Allowability" to avoid abandonment. See 37 CFR 1.53(f).

- A properly executed inventor's oath or declaration has not been received for the following inventor(s):  
Lenny Dang

Replies must be received in the USPTO within the set time period or must include a proper Certificate of Mailing or Transmission under 37 CFR 1.8 with a mailing or transmission date within the set time period. For more information and a suggested format, see Form PTO/SB/92 and MPEP 512.

Replies should be mailed to:

Mail Stop Missing Parts  
Commissioner for Patents  
P.O. Box 1450  
Alexandria VA 22313-1450

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web, including a copy of this Notice and selecting the document description "Applicant response to Pre-Exam Formalities Notice".  
<https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at 1-866-217-9197 or visit our website at <http://www.uspto.gov/ebc>.

If you are not using EFS-Web to submit your reply, you must include a copy of this notice.

Questions about the contents of this notice and the requirements it sets forth should be directed to the Office of Data Management, Application Assistance Unit, at **(571) 272-4000** or **(571) 272-4200** or **1-888-786-0101**.

/sareebuddin/

---

**PATENT APPLICATION FEE DETERMINATION RECORD**

Substitute for Form PTO-875

Application or Docket Number  
15/589,615

**APPLICATION AS FILED - PART I**

(Column 1)		(Column 2)	SMALL ENTITY		OR	OTHER THAN SMALL ENTITY	
FOR	NUMBER FILED	NUMBER EXTRA	RATE(\$)	FEE(\$)		RATE(\$)	FEE(\$)
BASIC FEE (37 CFR 1.16(a), (b), or (c))	N/A	N/A	N/A			N/A	280
SEARCH FEE (37 CFR 1.16(k), (l), or (m))	N/A	N/A	N/A			N/A	600
EXAMINATION FEE (37 CFR 1.16(o), (p), or (q))	N/A	N/A	N/A			N/A	720
TOTAL CLAIMS (37 CFR 1.16(i))	18	minus 20 = *			OR	x 80 =	0.00
INDEPENDENT CLAIMS (37 CFR 1.16(h))	1	minus 3 = *				x 420 =	0.00
APPLICATION SIZE FEE (37 CFR 1.16(s))	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$310 (\$155 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).						800
MULTIPLE DEPENDENT CLAIM PRESENT (37 CFR 1.16(j))							0.00
* If the difference in column 1 is less than zero, enter "0" in column 2.			TOTAL			TOTAL	2400

**APPLICATION AS AMENDED - PART II**

AMENDMENT A	(Column 1)	(Column 2)	(Column 3)	SMALL ENTITY		OR	OTHER THAN SMALL ENTITY	
	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE(\$)	ADDITIONAL FEE(\$)		RATE(\$)	ADDITIONAL FEE(\$)
Total (37 CFR 1.16(i))	*	Minus **	=	x	=	OR	x	=
Independent (37 CFR 1.16(h))	*	Minus ***	=	x	=	OR	x	=
Application Size Fee (37 CFR 1.16(s))						OR		
FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))						OR		
				TOTAL ADD'L FEE		OR	TOTAL ADD'L FEE	
AMENDMENT B	(Column 1)	(Column 2)	(Column 3)	SMALL ENTITY		OR	OTHER THAN SMALL ENTITY	
	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE(\$)	ADDITIONAL FEE(\$)		RATE(\$)	ADDITIONAL FEE(\$)
Total (37 CFR 1.16(i))	*	Minus **	=	x	=	OR	x	=
Independent (37 CFR 1.16(h))	*	Minus ***	=	x	=	OR	x	=
Application Size Fee (37 CFR 1.16(s))						OR		
FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))						OR		
				TOTAL ADD'L FEE		OR	TOTAL ADD'L FEE	

\* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.  
 \*\* If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".  
 \*\*\* If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".  
 The "Highest Number Previously Paid For" (Total or Independent) is the highest found in the appropriate box in column 1.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 6 columns: APPLICATION NUMBER, FILING or 371(c) DATE, GRP ART UNIT, FIL FEE REC'D, ATTY DOCKET NO, TOT CLAIMS, IND CLAIMS. Values: 15/589,615, 05/08/2017, 2540, AGS-013C2, 18, 1

CONFIRMATION NO. 7017

UPDATED FILING RECEIPT



148106
GOODWIN PROCTER LLP
PATENT ADMINISTRATOR
100 Northern Avenue
BOSTON, MA 02210

Date Mailed: 07/28/2017

Receipt is acknowledged of this non-provisional patent application. The application will be taken up for examination in due course. Applicant will be notified as to the results of the examination. Any correspondence concerning the application must include the following identification information: the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please submit a written request for a Filing Receipt Correction. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections

Inventor(s)

Lenny Dang, Boston, MA;
Valeria Fantin, Burlingame, CA;
Stefan Gross, Brookline, MA;
Hyun G. Jang, Waltham, MA;
Shengfang Jin, Newton, MA;
Francesco G. Salituro, Marlborough, MA;
Jeffrey O. Saunders, Lincoln, MA;
Shin-San Michael Su, Boston, MA;
Katharine Yen, Wellesley, MA;

Applicant(s)

Lenny Dang, Boston, MA;
Valeria Fantin, Burlingame, CA;
Stefan Gross, Brookline, MA;
Hyun G. Jang, Waltham, MA;
Shengfang Jin, Newton, MA;
Francesco G. Salituro, Marlborough, MA;
Jeffrey O. Saunders, Lincoln, MA;
Shin-San Michael Su, Boston, MA;
Katharine Yen, Wellesley, MA;

Assignment For Published Patent Application

Agios Pharmaceuticals, Inc., Cambridge, MA

Power of Attorney: None

Domestic Priority data as claimed by applicant

This application is a CON of 13/939,519 07/11/2013

which is a CON of 13/256,396 11/29/2011 ABN  
which is a 371 of PCT/US10/27253 03/12/2010  
which claims benefit of 61/266,929 12/04/2009  
and claims benefit of 61/253,820 10/21/2009  
and claims benefit of 61/229,689 07/29/2009  
and claims benefit of 61/227,649 07/22/2009  
and claims benefit of 61/220,543 06/25/2009  
and claims benefit of 61/180,609 05/22/2009  
and claims benefit of 61/173,518 04/28/2009  
and claims benefit of 61/160,664 03/16/2009  
and claims benefit of 61/160,253 03/13/2009

**Foreign Applications** for which priority is claimed (You may be eligible to benefit from the **Patent Prosecution Highway** program at the USPTO. Please see <http://www.uspto.gov> for more information.) - None.

*Foreign application information must be provided in an Application Data Sheet in order to constitute a claim to foreign priority. See 37 CFR 1.55 and 1.76.*

**Permission to Access Application via Priority Document Exchange:** Yes

**Permission to Access Search Results:** Yes

Applicant may provide or rescind an authorization for access using Form PTO/SB/39 or Form PTO/SB/69 as appropriate.

**Projected Publication Date:** Incomplete Reply

**Non-Publication Request:** No

**Early Publication Request:** No

**Title**

METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS

**Preliminary Class**

**Statement under 37 CFR 1.55 or 1.78 for AIA (First Inventor to File) Transition Applications:** No

## **PROTECTING YOUR INVENTION OUTSIDE THE UNITED STATES**

Since the rights granted by a U.S. patent extend only throughout the territory of the United States and have no effect in a foreign country, an inventor who wishes patent protection in another country must apply for a patent in a specific country or in regional patent offices. Applicants may wish to consider the filing of an international application under the Patent Cooperation Treaty (PCT). An international (PCT) application generally has the same effect as a regular national patent application in each PCT-member country. The PCT process **simplifies** the filing of patent applications on the same invention in member countries, but **does not result** in a grant of "an international patent" and does not eliminate the need of applicants to file additional documents and fees in countries where patent protection is desired.

Almost every country has its own patent law, and a person desiring a patent in a particular country must make an application for patent in that country in accordance with its particular laws. Since the laws of many countries differ in various respects from the patent law of the United States, applicants are advised to seek guidance from specific foreign countries to ensure that patent rights are not lost prematurely.

Applicants also are advised that in the case of inventions made in the United States, the Director of the USPTO must issue a license before applicants can apply for a patent in a foreign country. The filing of a U.S. patent application serves as a request for a foreign filing license. The application's filing receipt contains further information and guidance as to the status of applicant's license for foreign filing.

Applicants may wish to consult the USPTO booklet, "General Information Concerning Patents" (specifically, the section entitled "Treaties and Foreign Patents") for more information on timeframes and deadlines for filing foreign patent applications. The guide is available either by contacting the USPTO Contact Center at 800-786-9199, or it can be viewed on the USPTO website at <http://www.uspto.gov/web/offices/pac/doc/general/index.html>.

For information on preventing theft of your intellectual property (patents, trademarks and copyrights), you may wish to consult the U.S. Government website, <http://www.stopfakes.gov>. Part of a Department of Commerce initiative, this website includes self-help "toolkits" giving innovators guidance on how to protect intellectual property in specific countries such as China, Korea and Mexico. For questions regarding patent enforcement issues, applicants may call the U.S. Government hotline at 1-866-999-HALT (1-866-999-4258).

**LICENSE FOR FOREIGN FILING UNDER  
Title 35, United States Code, Section 184  
Title 37, Code of Federal Regulations, 5.11 & 5.15**

**GRANTED**

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 37 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Bureau of Industry and Security, Department of Commerce (15 CFR parts 730-774); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

**NOT GRANTED**

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed



from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15(b).

---

### **SelectUSA**

The United States represents the largest, most dynamic marketplace in the world and is an unparalleled location for business investment, innovation, and commercialization of new technologies. The U.S. offers tremendous resources and advantages for those who invest and manufacture goods here. Through SelectUSA, our nation works to promote and facilitate business investment. SelectUSA provides information assistance to the international investor community; serves as an ombudsman for existing and potential investors; advocates on behalf of U.S. cities, states, and regions competing for global investment; and counsels U.S. economic development organizations on investment attraction best practices. To learn more about why the United States is the best country in the world to develop technology, manufacture products, deliver services, and grow your business, visit <http://www.SelectUSA.gov> or call +1-202-482-6800.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office  
Address: COMMISSIONER FOR PATENTS  
P.O. Box 1450  
Alexandria, Virginia 22313-1450  
www.uspto.gov

APPLICATION NUMBER	FILING OR 371(C) DATE	FIRST NAMED APPLICANT	ATTY. DOCKET NO./TITLE
15/589,615	05/08/2017	Lenny Dang	AGS-013C2

148106  
GOODWIN PROCTER LLP  
PATENT ADMINISTRATOR  
100 Northern Avenue  
BOSTON, MA 02210

**CONFIRMATION NO. 7017**

**37 CFR 1.48 ACKNOWLEDGEMENT  
LETTER**



Date Mailed: 07/28/2017

**NOTICE OF ACCEPTANCE OF REQUEST UNDER 37 CFR 1.48(a)**

This is in response to the applicant's request under 37 CFR 1.48(a) submitted on 07/20/2017.

The request under 37 CFR 1.48(a) to correct the inventorship, to correct or update the name of an inventor, or to correct the order of names of joint inventors is accepted.

Questions about the contents of this notice and the requirements it sets forth should be directed to the Office of Data Management, Application Assistance Unit, at (571) 272-4000 or (571) 272-4200 or 1-888-786-0101.

/sareebuddin/

## RAW SEQUENCE LISTING

Loaded by SCORE, no errors detected.

Application Serial Number: 15589615

Source: OPAP

Date Processed by SCORE: 07/25/2017

***ENTERED***

SEQUENCE LISTING

<110> Dang, Lenny  
Fantin, Valeria  
Gross, Stefan  
Jang Gyung, Hyun  
Jin, Shengfang  
Salituro G., Francesco  
Saunders O., Jeffrey  
Su, Shinsan  
Yen, Katherine

<120> METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED  
DISORDERS

<130> c2081-7013US

<140> US 15/589,615

<141> 2017-05-08

<150> 13256396

<151> 2012-02-03

<150> 61/266,929

<151> 2009-12-04

<150> 61/253,820

<151> 2009-10-21

<150> 61/229,689

<151> 2009-07-29

<150> 61/227,649

<151> 2009-07-22

<150> 61/220,543

<151> 2009-06-25

<150> 61/180,609

<151> 2009-05-22

<150> 61/173,518

<151> 2009-04-28

<150> 61/160,664

<151> 2009-03-16

<150> 61/160,253

<151> 2009-03-13

<160> 804

<170> PatentIn version 3.5

<210> 1

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 1

taatcatatg tccaaaaaaaa tcagt

25

<210> 2

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 2

taatctcgag tgaaagtttg gcctgagcta gtt

33

<210> 3

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic 8xHis tag

<400> 3

His His His His His His His His

1

5

<210> 4

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 4  
Ser Leu Glu His His His His His His His His  
1 5 10 .

<210> 5  
<211> 1245  
<212> DNA  
<213> Homo sapiens

<400> 5  
atgtccaaaa aatcagtg cggttctgtg gtagagatgc aaggagatga aatgacacga  
60  
atcatttggg aattgattaa agagaaactc atttttccct acgtggaatt ggatctacat  
120  
agctatgatt taggcataga gaatcgtgat gccaccaacg accaagtcac caaggatgct  
180  
gcagaagcta taaagaagca taatgttggc gtcaaagtgc cactatcac tctgatgag  
240  
aagaggggtg aggagttcaa gttgaaaca atgtggaaat caccaaatgg caccatcga  
300  
aatattctgg gtggcacggt cttcagagaa gccattatct gcaaaaatat cccccgctt  
360  
gtgagtggat gggtaaaacc tatcatcata ggtcgtcatg cttatgggga tcaatacaga  
420  
gcaactgatt ttgttgttcc tgggcctgga aaagtagaga taacctacac accaagtgc  
480  
ggaacccaaa aggtgacata cctggtacat aactttgaag aagggtgtg tgttgccatg  
540  
gggatgtata atcaagataa gtcaattgaa gattttgcac acagttcctt ccaaatggct  
600  
ctgtctaagg gttggccttt gtatctgagc accaaaaaca ctattctgaa gaaatgat  
660  
gggcgtttta aagacatctt tcaggagata tatgacaagc agtacaagtc ccagttttaa  
720  
gctcaaaaga tctggtatga gcataggctc atcgacgaca tgggtggcca agctatgaaa  
780  
tcagagggag gcttcatctg ggctgtaaa aactatgatg gtgacgtgca gtcggactct

840

gtggcccaag ggtatggctc tctcggcatg atgaccagcg tgctggtttg tccagatggc  
900

aagacagtag aagcagaggc tgcccacggg actgtaacct gtcactaccg catgtaccag  
960

aaaggacagg agacgtccac caatcccatt gcttccattt ttgcctggac cagaggggta  
1020

gcccacagag caaagcttga taacaataaa gagcttgctt tctttgcaaa tgctttggaa  
1080

gaagtctcta ttgagacaat tgaggctggc ttcattgacca aggacttggc tgcttgcatt  
1140

aaaggtttac ccaatgtgca acgttctgac tacttgaata catttgagtt catggataaa  
1200

cttggagaaa acttgaagat caaactagct caggccaaac tttaa  
1245

<210> 6

<211> 1297

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
polynucleotide

<400> 6

atgtccaaaa aaatcagtgg cggttctgtg gtagagatgc aaggagatga aatgacacga  
60

atcatttggg aattgattaa agagaaactc atttttccct acgtggaatt ggatctacat  
120

agctatgatt taggcataga gaatcgtgat gccaccaacg accaagtcac caaggatgct  
180

gcagaagcta taaagaagca taatgttggc gtcaaattgtg ccaactatcac tcttgatgag  
240

aagaggggtg aggagttcaa gttgaaacaa atgtggaaat caccaaatgg caccatacga  
300

aatattctgg gtggcacggt cttcagagaa gccattatct gcaaaaatat cccccggctt  
360

gtgagtggat gggtaaaacc tatcatcata ggtcgcatg cttatgggga tcaatacaga  
420

gcaactgatt ttgttgttcc tgggctgga aaagtagaga taacctacac accaagtgac  
480

ggaacccaaa aggtgacata cctggtacat aactttgaag aagggtggtg tgttgccatg  
540

gggatgtata atcaagataa gtcaattgaa gattttgcac acagttcctt ccaaattggct  
600

ctgtctaagg gttggccttt gtatctgagc accaaaaaca ctattctgaa gaaatatgat  
660

gggcgtttta aagacatctt tcaggagata tatgacaagc agtacaagtc ccagtttgaa  
720

gctcaaaaga tctggtatga gcataggctc atcgacgaca tgggtggcca agctatgaaa  
780

tcagagggag gcttcatctg ggctgtaaa aactatgatg gtgacgtgca gtcggactct  
840

gtggcccaag ggtatggctc tctcggcatg atgaccagcg tgctggtttg tccagatggc  
900

aagacagtag aagcagaggc tgcccacggg actgtaacc gtcactaccg catgtaccag  
960

aaaggacagg agacgtccac caatcccatt gcttccattt ttgcctggac cagagggtta  
1020

gcccacagag caaagcttga taacaataaa gagcttgctt tctttgcaaa tgctttggaa  
1080

gaagtctcta ttgagacaat tgaggctggc ttcatgacca aggacttggc tgcttgcatt  
1140

aaaggtttac ccaatgtgca acgttctgac tacttgaata catttgagtt catggataaa  
1200

cttgagaaa acttgaagat caaactagct caggccaaac tttcactcga gcaccaccac  
1260

caccaccacc accactaatt gattaalacc taggctg  
1297

<210> 7

<211> 1245

<212> DNA



Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

<b>REQUEST FOR TRANSFER OF A COMPUTER READABLE FORM UNDER 37 CFR 1.821(e)</b>			
Application No.:	15/589,615-Conf. #7017	First Named Inventor:	Lenny Dang
Filing Date:	May 8, 2017	Practitioner Docket No.:	AGS-013C2
Title of the Invention:	METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS		
<p>The sequence information in the paper copy or PDF file of the Sequence Listing submitted in this application in accordance with 37 CFR 1.821(c),</p> <p><input type="checkbox"/> as part of the originally-filed specification of this application</p> <p><input checked="" type="checkbox"/> as a separate amendment of the specification filed</p> <p style="padding-left: 40px;"><input checked="" type="checkbox"/> herewith</p> <p style="padding-left: 40px;"><input type="checkbox"/> on _____</p> <p>is identical to the sequence information in the</p> <p><input type="checkbox"/> only computer readable form</p> <p><input checked="" type="checkbox"/> last filed computer readable form</p> <p><input type="checkbox"/> other computer readable form, which was filed on _____</p> <p>in application number <u>13/256,396</u>, which was filed on <u>November 13, 2011</u>.</p> <p>This computer readable form was compliant with 37 CFR 1.821-1.825, and applicant hereby requests that it be used as the computer readable form for the present application, in accordance with 37 CFR 1.821(e).</p>			
The above referenced paper copy or PDF file of the Sequence Listing contains no new matter.			
<p>A sequence listing text file submitted via EFS-Web that complies with the requirements of 37 CFR 1.824(a) (2)-(6) and (b) (i.e., is a compliant sequence listing ASCII text file), serves as both the paper copy required by 37 CFR 1.821(c) and the CRF required by 37 CFR 1.821(e). If a user submits a compliant sequence listing ASCII text file via EFS-Web, the U.S. Patent and Trademark Office will not carry out a request to use a compliant computer readable "Sequence Listing" that is already on file for another application pursuant to 37 CFR 1.821(e) but will use the sequence listing submitted with the application as originally filed via EFS-Web.</p>			
It is understood that upon the transfer of a copy of the computer readable form to this application, the U.S. Patent and Trademark Office will update the copy of the computer readable form to reflect the application number and filing date for this application.			
Signature	/Ioana Davies/	Date	2017-07-20
Name (Print/Typed)	Ioana Davies	Practitioner Registration Number	75,817
<p><b>Note:</b> This form must be signed in accordance with 37 CFR 1.33. See 37 CFR 1.4(d) for signature requirements and certifications. Submit multiple forms if more than one signature is required.*</p>			
<p><input type="checkbox"/> *Total of <u>1</u> forms are submitted.</p>			

1189804\_1.TXT  
SEQUENCE LISTING

<110> Dang, Lenny  
Fantin, Valeria  
Gross, Stefan  
Jang Gyung, Hyun  
Jin, Shengfang  
Salituro G., Francesco  
Saunders O., Jeffrey  
Su, Shinsan  
Yen, Katherine

<120> METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS

<130> c2081-7013US

<140> 13/256,396

<141> 2011-09-13

<150> 61/266,929

<151> 2009-12-04

<150> 61/253,820

<151> 2009-10-21

<150> 61/229,689

<151> 2009-07-29

<150> 61/227,649

<151> 2009-07-22

<150> 61/220,543

<151> 2009-06-25

<150> 61/180,609

<151> 2009-05-22

<150> 61/173,518

<151> 2009-04-28

<150> 61/160,664

<151> 2009-03-16

<150> 61/160,253

<151> 2009-03-13

<160> 804

<170> PatentIn version 3.5

<210> 1

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 1

taatcatatg tccaaaaaaa tcagt

25

<210> 2  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 2  
 taatctcgag tgaaagtttg gcctgagcta gtt

33

<210> 3  
 <211> 8  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 8xHis tag

<400> 3  
 His His His His His His His His  
 1 5

<210> 4  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 peptide

<400> 4  
 Ser Leu Glu His His His His His His His His  
 1 5 10

<210> 5  
 <211> 1245  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 atgtccaaaa aatcagtg cggttctgtg gtagagatgc aaggagatga aatgacacga 60  
 atcatttggg aattgattaa agagaaactc atttttccct acgtggaatt ggatctacat 120  
 agctatgatt taggcataga gaatcgtgat gccaccaacg accaagtcac caaggatgct 180  
 gcagaagcta taaagaagca taatgttggc gtcaaatgtg cactatcac tcctgatgag 240  
 aagagggttg aggagttaa gttgaaaca atgtggaat caccaaatgg caccatacga 300  
 aatattctgg gtggcacggt cttcagagaa gccattatct gcaaaaatat cccccggctt 360  
 gtgagtggat gggtaaaacc tatcatcata ggtcgtcatg cttatgggga tcaatacaga 420  
 gcaactgatt ttgttgttcc tgggcctgga aaagtagaga taacctacac accaagtgc 480

## 1189804\_1.TXT

ggaacccaaa aggtgacata cctggtacat aactttgaag aagtggtgg tgttgccatg 540  
 gggatgtata atcaagataa gtcaattgaa gatthttgcac acagttcctt ccaaatggct 600  
 ctgtctaagg gttggccttt gtatctgagc accaaaaaca ctattctgaa gaaatatgat 660  
 gggcgthtta aagacatctt tcaggagata tatgacaagc agtacaagtc ccagthttgaa 720  
 gctcaaaaaga tctggtatga gcataggctc atcgacgaca tggthggccca agctatgaaa 780  
 tcagagggag gcttcatctg ggctgtaaa aactatgatg gtgacgtgca gtcggactct 840  
 gtggcccaag ggtatggctc tctcggcatg atgaccagcg tgctggthttg tccagatggc 900  
 aagacagtag aagcagaggc tgcccacggg actgtaaccg gtcactaccg catgtaccag 960  
 aaaggacagg agacgtccac caatcccatt gcttccattt thgcctggac cagagggthta 1020  
 gccacagag caaagcttga taacaataaa gagctthgct tctthgcaaa thgctthggaa 1080  
 gaagtctcta thgagacaat thgaggtggc thcatgacca aggactthggc thgctthgatt 1140  
 aaagthttac ccaatgtgca acgthctgac thctthgaata cthttgagth catggataaa 1200  
 cthggagaaa actthgaagat caaactagct caggccaaaac thtaa 1245

<210> 6  
 <211> 1297  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 6  
 atgtccaaaa aatcagthg cgthtctgtg gtagagatgc aaggagatga aatgacacga 60  
 atcattthggg aattgattaa agagaaaactc atthttccct acgthggaatt ggatctacat 120  
 agctatgatt taggcataga gaatcgtgat gccaccaacg accaagthc acaggtatgct 180  
 gcagaagcta taaagaagca taatgthggc gtcaaatgtg ccactatcac tctgathgag 240  
 aagagggthg aggagthcaa gthgaaacaa atgtggaaat caccaaatgg caccatacga 300  
 aatattctgg thggcacggc cthcagagaa gccattatct gcaaaaatath cccccggct 360  
 gthgathggat gggthaaaacc thcatcata gthcgtcatg cthathgggga thcaatacaga 420  
 gcaactgatt thgthgthcc thggcctgga aaagthgaga thacctacac accaagthgac 480  
 ggaacccaaa aggtgacata cctggtacath aactthgaa aagthggtgg thgttgccatg 540  
 gggatgtata atcaagataa gtcaattgaa gatthttgcac acagthcctt ccaaatggct 600  
 ctgtctaagg gthggcctth gtatctgagc accaaaaaca ctattctgaa gaaatatgat 660  
 gggcgthtta aagacatctt tcaggagata thgacaagc agtacaagtc ccagthttgaa 720  
 gctcaaaaaga tctggtatga gcataggctc atcgacgaca thgtggccca agctatgaaa 780

## 1189804\_1.TXT

tcagagggag gcttcatctg ggccctgtaaa aactatgatg gtgacgtgca gtcggactct	840
gtggcccaag ggtatggctc tctcggcatg atgaccagcg tgctggtttg tccagatggc	900
aagacagtag aagcagaggc tgcccacggg actgtaaccc gtcactaccg catgtaccag	960
aaaggacagg agacgtccac caatcccatt gcttccattt ttgcctggac cagagggtta	1020
gcccacagag caaagcttga taacaataaa gagcttgccct tctttgcaaa tgctttggaa	1080
gaagtctcta ttgagacaat tgaggctggc ttcattgacca aggacttggc tgcttgatt	1140
aaaggtttac ccaatgtgca acgttctgac tacttgaata catttgagtt catggataaa	1200
cttgagaaa acttgaagat caaactagct caggccaac tttactcga gcaccaccac	1260
caccaccacc accactaatt gattaatacc taggctg	1297

&lt;210&gt; 7

&lt;211&gt; 1245

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Synthetic consensus sequence

&lt;400&gt; 7

atgtccaaaa aaatcagtgg cggttctgtg gtagagatgc aaggagatga aatgacacga	60
atcatttggg aattgattaa agagaaactc atttttccct acgtggaatt ggatctacat	120
agctatgatt taggcataga gaatcgtgat gccaccaacg accaagtcac caaggatgct	180
gcagaagcta taaagaagca taatgttggc gtcaaatgtg ccaactatcac tcctgatgag	240
aagagggttg aggagttcaa gttgaaacaa atgtggaaat caccaaatgg caccatacga	300
aatattctgg gtggcacggc cttcagagaa gccattatct gcaaaaatat cccccggctt	360
gtgagtggat gggtaaaacc tatcatcata ggtcgtcatg cttatgggga tcaatacaga	420
gcaactgatt ttgttgttcc tgggcctgga aaagtagaga taacctacac accaagtgc	480
ggaacccaaa aggtgacata cctggtacat aactttgaag aaggtgggtg tgttgccatg	540
gggatgtata atcaagataa gtcaattgaa gattttgcac acagttcctt ccaaatggct	600
ctgtctaagg gttggccttt gtatctgagc accaaaaaca ctattctgaa gaaatatgat	660
gggcgtttta aagacatctt tcaggagata tatgacaagc agtacaagtc ccagtttgaa	720
gctcaaaaga tctggtatga gcataggctc atcgacgaca tgggtggcca agctatgaaa	780
tcagagggag gcttcatctg ggccctgtaaa aactatgatg gtgacgtgca gtcggactct	840
gtggcccaag ggtatggctc tctcggcatg atgaccagcg tgctggtttg tccagatggc	900
aagacagtag aagcagaggc tgcccacggg actgtaaccc gtcactaccg catgtaccag	960
aaaggacagg agacgtccac caatcccatt gcttccattt ttgcctggac cagagggtta	1020

1189804\_1.TXT

gcccacagag caaagcttga taacaataaa gagcttgcct tctttgcaaa tgctttggaa 1080  
 gaagtctcta ttgagacaat tgaggctggc ttcattgacca aggacttggc tgcttgatt 1140  
 aaaggtttac ccaatgtgca acgttctgac tacttgaata catttgagtt catggataaa 1200  
 cttggagaaa acttgaagat caaactagct caggccaaac tttma 1245

<210> 8  
 <211> 1245  
 <212> DNA  
 <213> Homo sapiens

<400> 8  
 atgtccaaaa aaatcagtgg cggttctgtg gtagagatgc aaggagatga aatgacacga 60  
 atcatttggg aattgattaa agagaaactc atttttccct acgtggaatt ggatctacat 120  
 agctatgatt taggcataga gaatcgtgat gccaccaacg accaagtcac caaggatgct 180  
 gcagaagcta taaagaagca taatgttggc gtcaaatgtg ccactatcac tctgatgag 240  
 aagagggttg aggagttaa gttgaaacaa atgtggaaat caccaaatgg caccatacga 300  
 aatattctgg gtggcacggt cttcagagaa gccattatct gcaaaaatat cccccggctt 360  
 gtgagtggat gggtaaaacc tatcatcata ggtcgtcatg cttatgggga tcaatacaga 420  
 gcaactgatt ttgttgttcc tgggcctgga aaagtagaga taacctacac accaagtgac 480  
 ggaacccaaa aggtgacata cctggtacat aactttgaag aagggtgggg tggtgcatg 540  
 gggatgtata atcaagataa gtcaattgaa gatthttgac acagttcctt ccaaatggct 600  
 ctgtctaagg gttggccttt gtatctgagc accaaaaaca ctattctgaa gaaatgatg 660  
 gggcgthtta aagacatctt tcaggagata tatgacaagc agtacaagtc ccagthtgaa 720  
 gctcaaaaga tctggtatga gcataggctc atcgacgaca tgggtggcca agctatgaaa 780  
 tcagaggag gcttcatctg ggctgtgaaa aactatgatg gtgacgtgca gtcggactct 840  
 gtggcccaag ggtatggctc tctcggcatg atgaccagcg tgctggthtg tccagatggc 900  
 aagacagtag aagcagaggc tgcccacggg actgtaacc gtactaccg catgtaccag 960  
 aaaggacagg agacgtccac caatcccatt gcttccattt ttgcctggac cagagggtta 1020  
 gcccacagag caaagcttga taacaataaa gagcttgcct tctttgcaaa tgctttggaa 1080  
 gaagtctcta ttgagacaat tgaggctggc ttcattgacca aggacttggc tgcttgatt 1140  
 aaaggtttac ccaatgtgca acgttctgac tacttgaata catttgagtt catggataaa 1200  
 cttggagaaa acttgaagat caaactagct caggccaaac tttma 1245

<210> 9  
 <211> 2339  
 <212> DNA  
 <213> Homo sapiens

## 1189804\_1.TXT

<400> 9  
cctgtggtcc cgggtttctg cagagtctac ttcagaagcg gaggcactgg gagtccggtt 60  
tgggattgcc aggctgtggt tgtgagtctg agcttgtgag cggctgtggc gccccaactc 120  
ttcgccagca tatcatcccg gcaggcgata aactacattc agttgagtct gcaagactgg 180  
gaggaactgg ggtgataaga aatctattca ctgtcaaggt ttattgaagt caaaatgtcc 240  
aaaaaatca gtggcggttc tgtggtagag atgcaaggag atgaaatgac acgaatcatt 300  
tgggaattga ttaaagagaa actcattttt ccctacgtgg aattggatct acatagctat 360  
gatttaggca tagagaatcg tgatgccacc aacgaccaag tcaccaagga tgctgcagaa 420  
gctataaaga agcataatgt tggcgtcaaa tgtgccacta tcaactctga tgagaagagg 480  
gttgaggagt tcaagttgaa acaaatgtgg aaatcaccaa atggcaccat acgaaatatt 540  
ctgggtggca cggctctcag agaagccatt atctgcaaaa atatcccccg gcttgtgagt 600  
ggatgggtaa aacctatcat cataggctcg catgcttatg gggatcaata cagagcaact 660  
gattttgttg ttcctgggcc tggaaaagta gagataacct acacaccaag tgacggaacc 720  
caaaaggaga catacctggt acataacttt gaagaagggtg gtggtgttgc catggggatg 780  
tataatcaag ataagtcaat tgaagatfff gcacacagtt ccttccaaat ggctctgtct 840  
aagggttggc ctttgtatct gagcaccaa aacactattc tgaagaaata tgatgggcgt 900  
tttaagaca tctttcagga gatatatgac aagcagtaca agtcccagtt tgaagtcaa 960  
aagatctggt atgagcatag gctcatcgac gacatggtgg cccaagctat gaaatcagag 1020  
ggaggcttca tctgggcctg taaaaactat gatggtgacg tgcagtcgga ctctgtggcc 1080  
caagggtatg gctctctcgg catgatgacc agcgtgctgg tttgtccaga tggcaagaca 1140  
gtagaagcag aggctgcca cgggactgta acccgtcact accgcatgta ccagaaagga 1200  
caggagacgt ccaccaatcc cattgcttcc atttttgcct ggaccagagg gttagcccac 1260  
agagcaaagc ttgataacaa taaagagcct gccttctttg caaatgcttt ggaagaagtc 1320  
tctattgaga caattgaggc tggcttcatg accaaggact tggctgcttg cattaaaggt 1380  
ttaccaatg tgcaacgttc tgactacttg aatacatttg agttcatgga taaacttgga 1440  
gaaaacttga agatcaaaact agctcaggcc aaactttaag ttcataactg agctaagaag 1500  
gataattgtc ttttggtaac taggtctaca ggtttacatt tttctgtggt aactcaagg 1560  
ataaaggcaa aatcaatfff gtaatftgtt tagaagccag agtttatctt ttctataagt 1620  
ttacagcctt tttcttatat atacagttat tgccaccttt gtgaacatgg caaggactt 1680  
ttttacaatt tttatfttat tttctagtag cagcctagga attcggtttag tactcatttg 1740  
tattcactgt cactftttct catgttctaa ttataaatga ccaaaatcaa gattgctcaa 1800  
aagggtaaat gatagccaca gtattgctcc ctaaaatattg cataaagtag aaattcactg 1860

1189804\_1.TXT

ctttcccctc ctgtccatga ccttggggcac aggggaagttc tgggtgcata gatatcccgt 1920  
 tttgtgaggt agagctgtgc attaaacttg cacatgactg gaacgaagta tgagtgaac 1980  
 tcaaagtgtg tgaagatact gcagtcattt ttgtaaagac cttgctgaat gtttccaata 2040  
 gactaaatac tgtttaggcc gcaggagagt ttggaatccg gaataaatac tacctggagg 2100  
 tttgtcctct ccatttttct ctttctctc ctggcctggc ctgaatatta tactactcta 2160  
 aatagcatat ttcaccaag tgcaataatg taagctgaat cttttttgga cttctgctgg 2220  
 cctgttttat ttctttata taaatgtgat ttctcagaaa ttgatattaa acactatctt 2280  
 atcttctcct gaactgttga ttttaattaa aattaagtgc taattaccaa aaaaaaaaa 2339

<210> 10  
 <211> 452  
 <212> PRT  
 <213> Homo sapiens

<400> 10  
 Met Ala Gly Tyr Leu Arg Val Val Arg Ser Leu Cys Arg Ala Ser Gly  
 1 5 10 15

Ser Arg Pro Ala Trp Ala Pro Ala Ala Leu Thr Ala Pro Thr Ser Gln  
 20 25 30

Glu Gln Pro Arg Arg His Tyr Ala Asp Lys Arg Ile Lys Val Ala Lys  
 35 40 45

Pro Val Val Glu Met Asp Gly Asp Glu Met Thr Arg Ile Ile Trp Gln  
 50 55 60

Phe Ile Lys Glu Lys Leu Ile Leu Pro His Val Asp Ile Gln Leu Lys  
 65 70 75 80

Tyr Phe Asp Leu Gly Leu Pro Asn Arg Asp Gln Thr Asp Asp Gln Val  
 85 90 95

Thr Ile Asp Ser Ala Leu Ala Thr Gln Lys Tyr Ser Val Ala Val Lys  
 100 105 110

Cys Ala Thr Ile Thr Pro Asp Glu Ala Arg Val Glu Glu Phe Lys Leu  
 115 120 125

Lys Lys Met Trp Lys Ser Pro Asn Gly Thr Ile Arg Asn Ile Leu Gly  
 130 135 140

Gly Thr Val Phe Arg Glu Pro Ile Ile Cys Lys Asn Ile Pro Arg Leu  
 145 150 155 160



1189804\_1.TXT

Val Pro Gly Trp Thr Lys Pro Ile Thr Ile Gly Arg His Ala His Gly  
 165 170

Asp Gln Tyr Lys Ala Thr Asp Phe Val Ala Asp Arg Ala Gly Thr Phe  
 180 185 190

Lys Met Val Phe Thr Pro Lys Asp Gly Ser Gly Val Lys Glu Trp Glu  
 195 200 205

Val Tyr Asn Phe Pro Ala Gly Gly Val Gly Met Gly Met Tyr Asn Thr  
 210 215 220

Asp Glu Ser Ile Ser Gly Phe Ala His Ser Cys Phe Gln Tyr Ala Ile  
 225 230 235 240

Gln Lys Lys Trp Pro Leu Tyr Met Ser Thr Lys Asn Thr Ile Leu Lys  
 245 250 255

Ala Tyr Asp Gly Arg Phe Lys Asp Ile Phe Gln Glu Ile Phe Asp Lys  
 260 265 270

His Tyr Lys Thr Asp Phe Asp Lys Asn Lys Ile Trp Tyr Glu His Arg  
 275 280 285

Leu Ile Asp Asp Met Val Ala Gln Val Leu Lys Ser Ser Gly Gly Phe  
 290 295 300

Val Trp Ala Cys Lys Asn Tyr Asp Gly Asp Val Gln Ser Asp Ile Leu  
 305 310 315 320

Ala Gln Gly Phe Gly Ser Leu Gly Leu Met Thr Ser Val Leu Val Cys  
 325 330 335

Pro Asp Gly Lys Thr Ile Glu Ala Glu Ala Ala His Gly Thr Val Thr  
 340 345 350

Arg His Tyr Arg Glu His Gln Lys Gly Arg Pro Thr Ser Thr Asn Pro  
 355 360 365

Ile Ala Ser Ile Phe Ala Trp Thr Arg Gly Leu Glu His Arg Gly Lys  
 370 375 380

Leu Asp Gly Asn Gln Asp Leu Ile Arg Phe Ala Gln Met Leu Glu Lys  
 385 390 395 400

Val Cys Val Glu Thr Val Glu Ser Gly Ala Met Thr Lys Asp Leu Ala  
 405 410 415

1189804\_1.TXT

Gly Cys Ile His Gly Leu Ser Asn Val Lys Leu Asn Glu His Phe Leu  
420 425 430

Asn Thr Thr Asp Phe Leu Asp Thr Ile Lys Ser Asn Leu Asp Arg Ala  
435 440 445

Leu Gly Arg Gln  
450

<210> 11  
<211> 1359  
<212> DNA  
<213> Homo sapiens

<400> 11  
atggccggct acctgccgggt cgtgcgctcg ctctgcagag cctcaggctc gcggccggcc 60  
tgggcgccgg cggccctgac agccccacc tcgcaagagc agccgcggcg ccactatgcc 120  
gacaaaagga tcaaggtggc gaagcccgtg gtggagatgg atggtgatga gatgaccctg 180  
attatctggc agttcatcaa ggagaagctc atcctgcccc acgtggacat ccagctaaag 240  
tattttgacc tcgggctccc aaaccgtgac cagactgatg accagggtcac cattgactct 300  
gcactggcca cccagaagta cagtgtggct gtcaagtgtg ccaccatcac ccctgatgag 360  
gcccgtgtgg aagagttcaa gctgaagaag atgtggaaaa gtccaatgg aactatccgg 420  
aacatcctgg gggggactgt cttccgggag cccatcatct gcaaaaacat cccacgccta 480  
gtccctggct ggaccaagcc catcaccatt ggcaggcacg cccatggcga ccagtacaag 540  
gccacagact ttgtggcaga cggggccggc actttcaaaa tggctttcac cccaaaagat 600  
ggcagtggtg tcaaggagtg ggaagtgtac aacttccccg caggcggcgt gggcatgggc 660  
atgtacaaca ccgacgagtc catctcaggt tttgcgaca gctgcttcca gtatgccatc 720  
cagaagaaat ggccgctgta catgagcacc aagaacacca tactgaaagc ctacgatggg 780  
cgtttcaagg acatcttcca ggagatcttt gacaagcact ataagaccga cttcgacaag 840  
aataagatct ggtatgagca cgggctcatt gatgacatgg tggctcaggt cctcaagtct 900  
tcgggtggct ttgtgtgggc ctgcaagaac tatgacggag atgtgcagtc agacatcctg 960  
gcccagggct ttggctccct tggcctgatg acgtccgtcc tggctctgcc tgatgggaag 1020  
acgattgagg ctgaggccgc tcatgggacc gtcacccgcc actatcggga gcaccagaag 1080  
ggccggccca ccagcaccaa ccccatcgcc agcatctttg cctggacacg tggcctggag 1140  
caccggggga agctggatgg gaaccaagac ctcatcaggt ttgccagat gctggagaag 1200  
gtgtgctgg agacggtgga gagtggagcc atgaccaagg acctggcggg ctgcattcac 1260  
ggcctcagca atgtgaagct gaacgagcac ttctgaaca ccacggactt cctcgacacc 1320

atcaagagca acctggacag agccctgggc aggcagtag 1359

<210> 12  
 <211> 1740  
 <212> DNA  
 <213> Homo sapiens

<400> 12  
 ccagcgtag cccgcggcca ggcagccggg aggagcggcg cgcgctcggc cctctcccgc 60  
 cctgctcgtt cgctctccag cttgggatgg ccggctacct gcgggtcgtg cgctcgtct 120  
 gcagagcctc aggctcgcgg ccggcctggg cgccggcggc cctgacagcc cccacctcgc 180  
 aagagcagcc gcggcgccac tatgccgaca aaaggatcaa ggtggcgaag cccgtggtgg 240  
 agatggatgg tgatgagatg acccgtatta tctggcagtt catcaaggag aagctcatcc 300  
 tgccccacgt ggacatccag ctaaagtatt ttgacctcgg gctcccaaac cgtgaccaga 360  
 ctgatgacca ggtcaccatt gactctgcac tggccacca gaagtacagt gtggctgtca 420  
 agtgtgccac catcacccct gatgaggccc gtgtggaaga gttcaagctg aagaagatgt 480  
 ggaaaagtcc caatggaact atccggaaca tcctgggggg gactgtcttc cgggagccca 540  
 tcatctgcaa aaacatccca cgcctagtcc ctggctggac caagcccatc accattggca 600  
 ggcacgcca tggcgaccag tacaaggcca cagactttgt ggagaccgg gccggcactt 660  
 tcaaaatggt cttcaccca aaagatggca gtggtgtcaa ggagtgggaa gtgtacaact 720  
 tccccgagg cggcgtggc atgggcatgt acaacaccga cgagtccatc tcaggttttg 780  
 cgcacagctg cttccagtat gccatccaga agaaatggcc gctgtacatg agcaccaaga 840  
 acaccatact gaaagcctac gatgggcggt tcaaggacat cttccaggag atctttgaca 900  
 agcactataa gaccgacttc gacaagaata agatctggta tgagcaccgg ctattgatg 960  
 acatggtggc tcaggtcctc aagtcttcgg gtggctttgt gtggcctgc aagaactatg 1020  
 acggagatgt gcagtcagac atcctggccc agggcttttg ctcccttggc ctgatgacgt 1080  
 ccgtcctggt ctgccctgat ggaagacga ttgaggctga ggccgctcat gggaccgtca 1140  
 cccgccacta tcgggagcac cagaagggcc ggcccaccag caccaacccc atcgccagca 1200  
 tctttgctg gacacgtggc ctggagcacc gggggaagct ggatgggaac caagacctca 1260  
 tcaggtttgc ccagatgctg gagaagggtg gcgtggagac ggtggagagt ggagccatga 1320  
 ccaaggacct ggcgggctgc attcacggcc tcagcaatgt gaagctgaac gagcacttcc 1380  
 tgaacaccac ggacttcctc gacacatca agagcaacct ggacagagcc ctgggcaggc 1440  
 agtaggggga ggcgccacc atggctgcag tggaggggcc agggctgagc cggcgggtcc 1500  
 tcctgagcgc ggcagagggt gacctcaca gccctctct ggaggcctt ctaggggatg 1560  
 ttttttata agccagatgt ttttaaaagc atatgtgtgt ttcccctcat ggtgacgtga 1620

1189804\_1.TXT

ggcaggagca gtgcgtttta cctcagccag tcagtatggt ttgcatactg taatttatat 1680  
tgcccttggga acacatgggtg ccatatttag ctactaaaaa gctcttcaca aaaaaaaaaa 1740

<210> 13  
<211> 414  
<212> PRT  
<213> Homo sapiens

<400> 13  
Met Ser Lys Lys Ile Ser Gly Gly Ser Val Val Glu Met Gln Gly Asp  
1 5 10 15

Glu Met Thr Arg Ile Ile Trp Glu Leu Ile Lys Glu Lys Leu Ile Phe  
20 25 30

Pro Tyr Val Glu Leu Asp Leu His Ser Tyr Asp Leu Gly Ile Glu Asn  
35 40 45

Arg Asp Ala Thr Asn Asp Gln Val Thr Lys Asp Ala Ala Glu Ala Ile  
50 55 60

Lys Lys His Asn Val Gly Val Lys Cys Ala Thr Ile Thr Pro Asp Glu  
65 70 75 80

Lys Arg Val Glu Glu Phe Lys Leu Lys Gln Met Trp Lys Ser Pro Asn  
85 90 95

Gly Thr Ile Arg Asn Ile Leu Gly Gly Thr Val Phe Arg Glu Ala Ile  
100 105 110

Ile Cys Lys Asn Ile Pro Arg Leu Val Ser Gly Trp Val Lys Pro Ile  
115 120 125

Ile Ile Gly Arg His Ala Tyr Gly Asp Gln Tyr Arg Ala Thr Asp Phe  
130 135 140

Val Val Pro Gly Pro Gly Lys Val Glu Ile Thr Tyr Thr Pro Ser Asp  
145 150 155 160

Gly Thr Gln Lys Val Thr Tyr Leu Val His Asn Phe Glu Glu Gly Gly  
165 170 175

Gly Val Ala Met Gly Met Tyr Asn Gln Asp Lys Ser Ile Glu Asp Phe  
180 185 190

Ala His Ser Ser Phe Gln Met Ala Leu Ser Lys Gly Trp Pro Leu Tyr  
195 200 205

1189804\_1.TXT

Leu Ser Thr Lys Asn Thr Ile Leu Lys Lys Tyr Asp Gly Arg Phe Lys  
 210 215 220

Asp Ile Phe Gln Glu Ile Tyr Asp Lys Gln Tyr Lys Ser Gln Phe Glu  
 225 230 235 240

Ala Gln Lys Ile Trp Tyr Glu His Arg Leu Ile Asp Asp Met Val Ala  
 245 250 255

Gln Ala Met Lys Ser Glu Gly Gly Phe Ile Trp Ala Cys Lys Asn Tyr  
 260 265 270

Asp Gly Asp Val Gln Ser Asp Ser Val Ala Gln Gly Tyr Gly Ser Leu  
 275 280 285

Gly Met Met Thr Ser Val Leu Val Cys Pro Asp Gly Lys Thr Val Glu  
 290 295 300

Ala Glu Ala Ala His Gly Thr Val Thr Arg His Tyr Arg Met Tyr Gln  
 305 310 315 320

Lys Gly Gln Glu Thr Ser Thr Asn Pro Ile Ala Ser Ile Phe Ala Trp  
 325 330 335

Thr Arg Gly Leu Ala His Arg Ala Lys Leu Asp Asn Asn Lys Glu Leu  
 340 345 350

Ala Phe Phe Ala Asn Ala Leu Glu Glu Val Ser Ile Glu Thr Ile Glu  
 355 360 365

Ala Gly Phe Met Thr Lys Asp Leu Ala Ala Cys Ile Lys Gly Leu Pro  
 370 375 380

Asn Val Gln Arg Ser Asp Tyr Leu Asn Thr Phe Glu Phe Met Asp Lys  
 385 390 395 400

Leu Gly Glu Asn Leu Lys Ile Lys Leu Ala Gln Ala Lys Leu  
 405 410

<210> 14  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 14  
 gguuucugca gagucuacu

<210> 15  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 15  
aguagacucu gcagaaacc 19

<210> 16  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 16  
cucuucgcca gcauaucau 19

<210> 17  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 17  
augauaugcu ggcgaagag 19

<210> 18  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 18  
ggcaggcgau aaacuacau 19

<210> 19  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

1189804\_1.TXT

<400> 19  
auguaguuuu ucgccugcc 19

<210> 20  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 20  
gcgauaaaacu acauucagu 19

<210> 21  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 21  
acugaaugua guuuaucgc 19

<210> 22  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 22  
gaaaucuauu cacugucac 19

<210> 23  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 23  
uugacaguga auagauuuc 19

<210> 24  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

oligonucleotide

<400> 24  
guucuguggu agaugca 19

<210> 25  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 25  
ugcaucucua ccacagaac 19

<210> 26  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 26  
gcaaggagau gaaugaca 19

<210> 27  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 27  
ugucauuuca ucuccuugc 19

<210> 28  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 28  
ggagaugaaa ugacacgaa 19

<210> 29  
<211> 19  
<212> RNA  
<213> Artificial Sequence



1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 29  
uucgugucau uucaucc 19

<210> 30  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 30  
gagaugaaau gacacgaau 19

<210> 31  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 31  
auucguguca uucaucuc 19

<210> 32  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 32  
gaugaauga cacgaauca 19

<210> 33  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 33  
ugauucgugu cauucauc 19

<210> 34  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 34

cgaaucauuu gggaaauga

19

<210> 35

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 35

ucaauuccca aaugauucg

19

<210> 36

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 36

gggaauugau uaaagagaa

19

<210> 37

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 37

uucucuuaa ucaauuccc

19

<210> 38

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 38

ccuacgugga auuggaucu

19

<210> 39

<211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 39  
 agauccaau ccacguagg 19

<210> 40  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 40  
 cuacguggaa uuggaucua 19

<210> 41  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 41  
 uagaucacau uccacguag 19

<210> 42  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 42  
 ggaucuacau agcuaugau 19

<210> 43  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 43  
 aucauagcua uguagaucc 19

1189804\_1.TXT

<210> 44  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 44  
gcuaugauuu aggcauaga 19

<210> 45  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 45  
ucuaugccua aaucauagc 19

<210> 46  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 46  
ggaugcugca gaagcuaua 19

<210> 47  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 47  
uauagcuucu gcagcaucc 19

<210> 48  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 48

	1189804_1.TXT	
cagaagcuau aaagaagca		19
<210> 49		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 49		
ugcuucuuua uagcuucug		19
<210> 50		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 50		
gaagcuauaa agaagcaua		19
<210> 51		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 51		
uaugcuucuu uauagcuuc		19
<210> 52		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 52		
gcuaaauuu ggcgucaaa		19
<210> 53		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		

1189804\_1.TXT

<400> 53  
uuugacgccca acauuauugc 19

<210> 54  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 54  
cugaugagaa gaggguuga 19

<210> 55  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 55  
ucaaccucucu ucucaucag 19

<210> 56  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 56  
guugaggagu ucaaguuga 19

<210> 57  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 57  
ucaacuugaa cuccaac 19

<210> 58  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 58  
gaguucaagu ugaacaaca 19

<210> 59  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 59  
uuuguuucaa cuugaacuc 19

<210> 60  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 60  
guugaaacaa auguggaaa 19

<210> 61  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 61  
uuuccacauu uguuugaac 19

<210> 62  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 62  
caaaugugga aaucaccaa 19

<210> 63  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 63  
uuggugauuu ccacauug 19

<210> 64  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 64  
ccaaauggca ccuacgaa 19

<210> 65  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 65  
uucguuggu gccauugg 19

<210> 66  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 66  
cauacgaaau auucuggu 19

<210> 67  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 67  
accagaaua uuucguug 19

<210> 68  
<211> 19



<212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 68  
 gagaagccau uaucugcaa 19

<210> 69  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 69  
 uugcagauaa uggcuucuc 19

<210> 70  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 70  
 cuaucaucau aggucguca 19

<210> 71  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 71  
 ugacgaccua ugaugauag 19

<210> 72  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 72  
 caucauaggu cgucaugcu 19

<210> 73  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 73  
 agcaugacga ccuaugaug 19

<210> 74  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 74  
 cauaggucgu caugcuuau 19

<210> 75  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 75  
 auaagcauga cgaccuaug 19

<210> 76  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 76  
 gagauaaccu acacaccaa 19

<210> 77  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 77  
 uuggugugua gguaucuc 19

1189804\_1.TXT

<210> 78  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 78  
ccugguacau aacuuugaa 19

<210> 79  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 79  
uucaaaguua uguaccagg 19

<210> 80  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 80  
cuuugaagaa ggugguggu 19

<210> 81  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 81  
accaccaccu ucuucaaag 19

<210> 82  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

1189804\_1.TXT

<400> 82  
gggauguaua aucaagaua 19

<210> 83  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 83  
uaucuugauu auacauccc 19

<210> 84  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 84  
gcacacaguu ccuuccaaa 19

<210> 85  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 85  
uuuggaagga acugugugc 19

<210> 86  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 86  
guuccuucca aauggcucu 19

<210> 87  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

## oligonucleotide

<400> 87  
agagccauuu ggaaggaac 19

<210> 88  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 88  
gguuggccuu uguaucuga 19

<210> 89  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 89  
ucagauaca aggccaacc 19

<210> 90  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 90  
cuuuguaucu gagcaccaa 19

<210> 91  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 91  
uuggugcuca gauacaaag 19

<210> 92  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 92  
gaagaaauau gaugggcu 19

<210> 93  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 93  
acgcccauca uuuuucuc 19

<210> 94  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 94  
guccaguuu gaagcuca 19

<210> 95  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 95  
uugagcuuca aacugggac 19

<210> 96  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 96  
gguaugagca uaggcucau 19

<210> 97  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 97

augagccuau gcucauacc

19

<210> 98

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 98

ggcccaagcu augaaauca

19

<210> 99

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 99

ugauuucaua gcuugggcc

19

<210> 100

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 100

cccaagcuau gaaaucaga

19

<210> 101

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 101

ucugauuuca uagcuuggg

19

<210> 102

<211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 102  
 cagauggcaa gacaguaga 19  
  
 <210> 103  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 103  
 ucuacugucu ugccaucug 19  
  
 <210> 104  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 104  
 gcaagacagu agaagcaga 19  
  
 <210> 105  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 105  
 ucugcuucua cugucuugc 19  
  
 <210> 106  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 106  
 gcauguacca gaaaggaca 19



1189804\_1.TXT

<210> 107  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 107  
uguccuuucu gguacaugc 19

<210> 108  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 108  
ccaaucccau ugcuccau 19

<210> 109  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 109  
auggaagcaa ugggauugg 19

<210> 110  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 110  
ccacagagca aagcuugau 19

<210> 111  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 111

1189804\_1.TXT  
aucaagcuuu gcucugugg 19

<210> 112  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 112  
cacagagcaa agcuugaua 19

<210> 113  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 113  
uaucaagcuu ugcucugug 19

<210> 114  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 114  
gagcaaagcu ugauaaca 19

<210> 115  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 115  
uuguuaucaa gcuuugcuc 19

<210> 116  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 116  
gagcuugccu ucuuugcaa 19

<210> 117  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 117  
uugcaaagaa ggcaagcuc 19

<210> 118  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 118  
cuuugcaaa u gcuuuggaa 19

<210> 119  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 119  
uuccaaagca uuugcaaag 19

<210> 120  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 120  
caaaugcuuu ggaagaagu 19

<210> 121  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 121  
 acuucuucca aagcauuug 19

<210> 122  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 122  
 cuuuggaaga agucucuau 19

<210> 123  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 123  
 auagagacuu cuuccaaag 19

<210> 124  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 124  
 gaagaagucu cuauugaga 19

<210> 125  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 125  
 ucucauaga gacuucuuc 19

<210> 126  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 126  
gaagucucua uugagacaa 19

<210> 127  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 127  
uugucucaau agagacuuc 19

<210> 128  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 128  
ggacuuggcu gcuugcauu 19

<210> 129  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 129  
aaugcaagca gccaagucc 19

<210> 130  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 130  
cuuggcugcu ugcauuaaa 19

<210> 131  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 131  
 uuuaaugcaa gcagccaag 19

<210> 132  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 132  
 cauuaaaggu uuacccaau 19

<210> 133  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 133  
 auuggguaaa ccuuuaug 19

<210> 134  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 134  
 ccaaugugca acguucuga 19

<210> 135  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 135  
 ucagaacguu gcacauugg 19

<210> 136  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 136  
 gugcaacguu cugacuacu 19

<210> 137  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 137  
 aguagucaga acguugcac 19

<210> 138  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 138  
 cguucugacu acuugaaua 19

<210> 139  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 139  
 uauucaagua gucagaacg 19

<210> 140  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 140  
 cauuugaguu cauggauaa 19

<210> 141  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 141  
 uuaucauga acucaaaug 19

<210> 142  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 142  
 guucauggau aaacuugga 19

<210> 143  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 143  
 uccaaguuaa uccaugaac 19

<210> 144  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 144  
 cauggauaaa cuuggagaa 19

<210> 145  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide



1189804\_1.TXT

<400> 145  
uucuccaagu uuauccaug 19

<210> 146  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 146  
caaacuagcu caggccaaa 19

<210> 147  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 147  
uuuggccuga gcuaguuug 19

<210> 148  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 148  
ccugagcuaa gaaggauaa 19

<210> 149  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 149  
uuaucuuucu uagcucagg 19

<210> 150  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

oligonucleotide

<400> 150  
cuaagaagga uaaugucu 19

<210> 151  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 151  
agacaauuau ccuucuug 19

<210> 152  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 152  
cuguguuaca cucaaggau 19

<210> 153  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 153  
auccuugagu guaacacag 19

<210> 154  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 154  
guguuacacu caaggauaa 19

<210> 155  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 155  
uuauccuuga guguaacac 19

<210> 156  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 156  
cacucaagga uaaaggcaa 19

<210> 157  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 157  
uugccuuuau ccuugagug 19

<210> 158  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 158  
guaauuuguu uagaagcca 19

<210> 159  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 159  
uggcuucuaa acaaaauac 19

<210> 160  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 160  
guuauugcca ccuuuguga 19

<210> 161  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 161  
ucacaaaggu ggcauaaac 19

<210> 162  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 162  
cagccuagga auucgguaa 19

<210> 163  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 163  
uaaccgaauu ccuaggcug 19

<210> 164  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 164  
gccuaggaau ucgguuagu 19

<210> 165

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 165  
acuaaccgaa uccuaggc 19

<210> 166  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 166  
ccuaggaauu cgguuagua 19

<210> 167  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 167  
uacuaaccga auccuagg 19

<210> 168  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 168  
ggaauucggu uaguacuca 19

<210> 169  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 169  
ugaguacuaa ccgaaucc 19

<210> 170  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 170  
 gaauucgguu aguacucau 19

<210> 171  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 171  
 augaguacua accgaauuc 19

<210> 172  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 172  
 gguuaguacu cauuguau 19

<210> 173  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 173  
 auacaaauga guacuaacc 19

<210> 174  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 174

1189804\_1.TXT  
guacucauuu guauucacu 19

<210> 175  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 175  
agugaauaca aaugaguac 19

<210> 176  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 176  
gguaaaugau agccacagu 19

<210> 177  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 177  
acuguggcua ucauuuacc 19

<210> 178  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 178  
guaaaugaua gccacagua 19

<210> 179  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 179  
uacuguggcu aucauuuac 19

<210> 180  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 180  
ccacaguauu gcucccuaa 19

<210> 181  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 181  
uuaggagca auacugugg 19

<210> 182  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 182  
gggaaguucu ggugucaua 19

<210> 183  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 183  
uaugacacca gaacuuccc 19

<210> 184  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>



<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 184  
guucuggugu cauagauau 19

<210> 185  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 185  
auaucuauga caccagaac 19

<210> 186  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 186  
gcugugcauu aaacuugca 19

<210> 187  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 187  
ugcaaguuaa augcacagc 19

<210> 188  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 188  
gugcauuaaa cuugcacau 19

<210> 189  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 189  
augugcaagu uaaaugcac 19

<210> 190  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 190  
gcuuuaaacu ugcacauga 19

<210> 191  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 191  
ucaugugcaa guuuuauugc 19

<210> 192  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 192  
caugacugga acgaaguau 19

<210> 193  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 193  
auacuucguu ccagucaug 19

<210> 194  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 194  
 ggaacgaagu augagugca 19

<210> 195  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 195  
 ugcacucaua cuucguucc 19

<210> 196  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 196  
 gaacgaagua ugagugcaa 19

<210> 197  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 197  
 uugcacucau acuucguuc 19

<210> 198  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 198  
 gagugcaacu caaaugugu 19

<210> 199  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 199  
 acacauuga guugcacuc 19

<210> 200  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 200  
 gcaacucaa uguguugaa 19

<210> 201  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 201  
 uucaacacau uugaguugc 19

<210> 202  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 202  
 caaauuguu gaagauacu 19

<210> 203  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 203  
 agauucuca acacauug 19

<210> 204  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 204  
 guguugaaga uacugcagu 19

<210> 205  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 205  
 acugcaguau cuucaacac 19

<210> 206  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 206  
 guugaagaua cugcaguca 19

<210> 207  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 207  
 ugacugcagu aucuucac 19

<210> 208  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 208  
ccuugcugaa uguuuccaa 19

<210> 209  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 209  
uuggaaacau ucagcaagg 19

<210> 210  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 210  
cuugcugaau guuuccaau 19

<210> 211  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 211  
auuggaaaca uucagcaag 19

<210> 212  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 212  
gcugaauguu uccaauaga 19

<210> 213  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 53

oligonucleotide

<400> 213  
ucuaauuggaa acauucagc 19

<210> 214  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 214  
ccaauagacu aaauacugu 19

<210> 215  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 215  
acaguauuuu gucuauugg 19

<210> 216  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 216  
gaguauuggaa uccggaaua 19

<210> 217  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 217  
uauuccggau uccaaacuc 19

<210> 218  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 218  
ggaauccgga auaaaauacu 19

<210> 219  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 219  
aguauuuauu ccggauucc 19

<210> 220  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 220  
gaaucggaa uaaauacua 19

<210> 221  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 221  
uaguauuuau uccggauuc 19

<210> 222  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 222  
ggaauaaaaa cuaccugga 19

<210> 223  
<211> 19  
<212> RNA



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 223

uccagguagu auuuauucc

19

<210> 224

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 224

ggccuggccu gaauuuau

19

<210> 225

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 225

auaaauuca ggccaggcc

19

<210> 226

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 226

gccugaauau uauacuacu

19

<210> 227

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 227

aguaguauaa uauucaggc

19

<210> 228

<211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 228  
 cuggccugaa uauuauacu 19

<210> 229  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 229  
 aguauaaauau ucaggccag 19

<210> 230  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 230  
 cauauuucau ccaagugca 19

<210> 231  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 231  
 ugcacuugga ugaaauaug 19

<210> 232  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 232  
 gugcaauaau guaagcuga 19

1189804\_1.TXT

<210> 233  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 233  
ucagcuuaca uuauugcac 19

<210> 234  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 234  
gcaauaauugu aagcugaau 19

<210> 235  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 235  
auucagcuua cauuaugc 19

<210> 236  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 236  
cacuaucua ucuucuccu 19

<210> 237  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 237

	1189804_1.TXT	
aggagaagau aagauagug		19
<210> 238		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 238		
cuucuccuga acuguugau		19
<210> 239		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 239		
aucaacaguu caggagaag		19
<210> 240		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 240		
aaccuaucau cauaggucg		19
<210> 241		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		
<400> 241		
cgaccuauga ugauagguu		19
<210> 242		
<211> 19		
<212> RNA		
<213> Artificial Sequence		
<220>		
<223> Description of Artificial Sequence: Synthetic oligonucleotide		

1189804\_1.TXT

<400> 242  
accuaucauc auaggucgu 19

<210> 243  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 243  
acgaccuaug augauaggu 19

<210> 244  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 244  
ccuaucauca uaggucguc 19

<210> 245  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 245  
gacgaccuau gaugauagg 19

<210> 246  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 246  
cuaucacau aggucguca 19

<210> 247  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 247  
ugacgaccua ugaugauag 19

<210> 248  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 248  
uaucaucaua ggucgucau 19

<210> 249  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 249  
augacgaccu augaugaua 19

<210> 250  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 250  
aucaucauag gucgucaug 19

<210> 251  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 251  
caugacgacc uaugaugau 19

<210> 252  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 252  
ucaucauagg ucgucaugc 19

<210> 253  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 253  
gcaugacgac cuaugauga 19

<210> 254  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 254  
caucauaggu cgucaugcu 19

<210> 255  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 255  
agcaugacga ccuaugaug 19

<210> 256  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 256  
aucauagguc gucaugcu 19

<210> 257  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 257  
 aagcaugacg accuaugau 19

<210> 258  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 258  
 ucauaggucg ucaugcuua 19

<210> 259  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 259  
 uaagcaugac gaccuauga 19

<210> 260  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 260  
 cauaggucgu caugcuuau 19

<210> 261  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 261  
 auaagcauga cgaccuaug 19



<210> 262  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 262  
auaggucguc augcuuau 19

<210> 263  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 263  
cauaagcaug acgaccuau 19

<210> 264  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 264  
uaggucguca ugcuuaugg 19

<210> 265  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 265  
ccauaagcau gacgaccua 19

<210> 266  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 266  
aggucgucau gcuuaugg 19

<210> 267  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 267  
cccauaagca ugacgaccu 19

<210> 268  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 268  
ggucgucaug cuuaugggg 19

<210> 269  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 269  
ccccauaagc augacgacc 19

<210> 270  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 270  
gucgucaugc uuaugggga 19

<210> 271  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 271  
ucccauaagc augacgacc 19

<210> 272  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 272  
ucgucaugcu uauggggau 19

<210> 273  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 273  
aucccauaag caugacgac 19

<210> 274  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 274  
aaccuaucau cauagguca 19

<210> 275  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 275  
ugaccuauga ugauagguu 19

<210> 276  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 66

oligonucleotide

<400> 276  
accuaucauc auaggucau 19

<210> 277  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 277  
augaccuaug augauaggu 19

<210> 278  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 278  
ccuaucauca uaggucauc 19

<210> 279  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 279  
gaugaccuau gaugauagg 19

<210> 280  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 280  
cuaucaucau aggucauca 19

<210> 281  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 281  
ugaugaccua ugaugauag 19

<210> 282  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 282  
uaucaucaua ggucaucau 19

<210> 283  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 283  
augaugaccu augaugaua 19

<210> 284  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 284  
aucaucauag gucaucaug 19

<210> 285  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 285  
caugaugacc uaugaugau 19

<210> 286  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 286

ucaucauagg ucaucaugc

19

<210> 287

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 287

gcaugaugac cuaugauga

19

<210> 288

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 288

caucauaggu caucaugcu

19

<210> 289

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 289

agcaugauga ccuaugaug

19

<210> 290

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 290

aucauagguc aucaugcuu

19

<210> 291

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 291  
aagcaugaug accuaugau 19

<210> 292  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 292  
ucauagguca ucaugcuua 19

<210> 293  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 293  
uaagcaugau gaccuauga 19

<210> 294  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 294  
cauaggucan cauagcuau 19

<210> 295  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 295  
auaagcauga ugaccuau 19

1189804\_1.TXT

<210> 296  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 296  
auaggucauc augcuuaug 19  
  
<210> 297  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 297  
cauaagcaug augaccuau 19  
  
<210> 298  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 298  
uaggucauca ugcuuauagg 19  
  
<210> 299  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 299  
ccaauagcau gaugaccua 19  
  
<210> 300  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 300



1189804\_1.TXT  
aggucaucau gcuuauggg 19

<210> 301  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 301  
cccuaaagca ugaugaccu 19

<210> 302  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 302  
ggucaucaug cuuaugggg 19

<210> 303  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 303  
ccccauaagc augaugacc 19

<210> 304  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 304  
gucaucaugc uuaugggga 19

<210> 305  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 305  
uccccauaag caugaugac 19

<210> 306  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 306  
ucaucaugcu uauggggau 19

<210> 307  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 307  
auccccaaua gcaugauga 19

<210> 308  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 308  
aaccuaucau cauagguag 19

<210> 309  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 309  
cuaccuauga ugauagguu 19

<210> 310  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 310  
accuaucauc auagguagu 19

<210> 311  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 311  
acuaccuau augauaggu 19

<210> 312  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 312  
ccuaucauca uagguaguc 19

<210> 313  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 313  
gacuaccuau gaugauagg 19

<210> 314  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 314  
cuaucaucau agguaguca 19

<210> 315  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 315  
ugacuaccua ugaugauag 19

<210> 316  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 316  
uaucaucaua gguagucau 19

<210> 317  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 317  
augacuaccu augaugaua 19

<210> 318  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 318  
aucaucauag guagucaug 19

<210> 319  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 319  
caugacuacc uaugaugau 19

<210> 320  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 320  
 ucaucauagg uagucaugc 19

<210> 321  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 321  
 gcaugacuac cuaugauga 19

<210> 322  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 322  
 caucauaggu agucaugcu 19

<210> 323  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 323  
 agcaugacua ccuaugaug 19

<210> 324  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 324  
 aucauaggua gucaugcu 19

<210> 325  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 325  
aagcaugacu accuaugau 19

<210> 326  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 326  
ucauagguag ucaugcuua 19

<210> 327  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 327  
uaagcaugac uaccuauga 19

<210> 328  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 328  
cauagguagu caugcuuau 19

<210> 329  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 329  
auaagcauga cuaccuaug 19

1189804\_1.TXT

<210> 330  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 330  
auagguaguc augcuuau 19

<210> 331  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 331  
cauaagcaug acuaccuau 19

<210> 332  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 332  
uagguaguca ugcuuau 19

<210> 333  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 333  
ccaauagcau gacuaccua 19

<210> 334  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 334  
agguagucau gcuuauggg 19

<210> 335  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 335  
cccauaagca ugacuaccu 19

<210> 336  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 336  
gguagucaug cuuaugggg 19

<210> 337  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 337  
ccccauaagc augacuacc 19

<210> 338  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 338  
guagucaugc uuaugggga 19

<210> 339  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide



oligonucleotide

<400> 339  
uccccauaag caugacuac 19

<210> 340  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 340  
uagucaugcu uauggggau 19

<210> 341  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 341  
auccccaaua gcaugacua 19

<210> 342  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 342  
aaccuaucau cauagguug 19

<210> 343  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 343  
caaccuauga ugauagguu 19

<210> 344  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 344  
accuaucauc auagguugu 19

<210> 345  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 345  
acaaccuaug augauaggu 19

<210> 346  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 346  
ccuaucauca uagguuguc 19

<210> 347  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 347  
gacaaccuau gaugauagg 19

<210> 348  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 348  
cuaucaucau agguuguca 19

<210> 349  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 349

ugacaaccua ugaugauag

19

<210> 350

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 350

uaucaucaua gguugucau

19

<210> 351

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 351

augacaaccu augaugaua

19

<210> 352

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 352

aucaucauag guugucaug

19

<210> 353

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 353

caugacaacc uaugaugau

19

<210> 354

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 354  
ucaucauagg uugucaugc 19

<210> 355  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 355  
gcaugacaac cuaugauga 19

<210> 356  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 356  
caucauaggu ugucaugcu 19

<210> 357  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 357  
agcaugacaa ccuaugaug 19

<210> 358  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 358  
aucauagguu gucaugcuu 19

1189804\_1.TXT

<210> 359  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 359  
aagcaugaca accuaugau 19

<210> 360  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 360  
ucauagguug ucaugcuua 19

<210> 361  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 361  
uaagcaugac aaccuauga 19

<210> 362  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 362  
cauagguugu caugcuuau 19

<210> 363  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 363

1189804\_1.TXT  
auaagcauga caaccuau 19

<210> 364  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 364  
auagguuguc augcuuau 19

<210> 365  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 365  
cauaagcaug acaaccuau 19

<210> 366  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 366  
uagguuguca ugcuuauagg 19

<210> 367  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 367  
ccaauagcau gacaaccua 19

<210> 368  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 368  
agguugucau gcuauggg 19

<210> 369  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 369  
cccuaagca ugacaaccu 19

<210> 370  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 370  
gguugucaug cuuauggg 19

<210> 371  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 371  
ccccauaagc augacaacc 19

<210> 372  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 372  
guugucaugc uuaugggga 19

<210> 373  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 373  
uccccauaag caugacaac 19

<210> 374  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 374  
uugucaugcu uauggggau 19

<210> 375  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 375  
auccccaaua gcaugacaa 19

<210> 376  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 376  
aaccuaucau cauaggugg 19

<210> 377  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 377  
ccaccuauga ugauagguu 19

<210> 378  
<211> 19  
<212> RNA  
<213> Artificial Sequence



1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 378  
accuaucauc auagguggu 19

<210> 379  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 379  
accaccuau augauaggu 19

<210> 380  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 380  
ccuaucauca uaggugguc 19

<210> 381  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 381  
gaccacuau gaugauagg 19

<210> 382  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 382  
cuaucaucau aggugguca 19

<210> 383  
<211> 19

<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 383  
ugaccaccua ugaugauag 19

<210> 384  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 384  
uaucaucaua gguggucau 19

<210> 385  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 385  
augaccaccu augaugaua 19

<210> 386  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 386  
aucaucauag guggucaug 19

<210> 387  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 387  
caugaccacc uaugaugau 19

<210> 388  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 388  
ucaucauagg uggucaugc 19

<210> 389  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 389  
gcaugaccac cuaugauga 19

<210> 390  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 390  
caucauaggu ggucaugcu 19

<210> 391  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 391  
agcaugacca ccuaugaug 19

<210> 392  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 392  
aucauaggug gucaugcu 19

1189804\_1.TXT

<210> 393  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 393  
aagcaugacc accuaugau 19

<210> 394  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 394  
ucauaggugg ucaugcuua 19

<210> 395  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 395  
uaagcaugac cacuauga 19

<210> 396  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 396  
cauagguggu caugcuuau 19

<210> 397  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 397  
auaagcauga ccaccuau 19

<210> 398  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 398  
auaggguguc augcuuau 19

<210> 399  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 399  
cauaagcaug accaccuau 19

<210> 400  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 400  
uaggguguca ugcuuau 19

<210> 401  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 401  
ccaauagcau gaccaccua 19

<210> 402  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 92

oligonucleotide

<400> 402  
 agguugucau gcuuauggg 19

<210> 403  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 403  
 cccauaagca ugaccaccu 19

<210> 404  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 404  
 gguugucaug cuuaugggg 19

<210> 405  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 405  
 ccccauaagc augaccacc 19

<210> 406  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 406  
 guugucaugc uuaugggga 19

<210> 407  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 407  
uccccauaag caugaccac 19

<210> 408  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 408  
uugucaugcu uauggggau 19

<210> 409  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 409  
auccccaaua gcaugacca 19

<210> 410  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 410  
aaccuaucau cauaggucg 19

<210> 411  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 411  
cgaccuauga ugauagguu 19

<210> 412  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 412

accuaucauc auaggucgu

19

<210> 413

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 413

acgaccuaug augauaggu

19

<210> 414

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 414

ccuaucauca uaggucguc

19

<210> 415

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 415

gacgaccuau gaugauagg

19

<210> 416

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 416

cuaucacau aggucguca

19

<210> 417



<211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 417  
 ugacgaccua ugaugauag 19

<210> 418  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 418  
 uaucaucaua ggucgucau 19

<210> 419  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 419  
 augacgaccu augaugaua 19

<210> 420  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 420  
 aucaucauag gucgucaug 19

<210> 421  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 421  
 caugacgacc uaugaugau 19

1189804\_1.TXT

<210> 422  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 422  
ucaucauagg ucgucaugc 19

<210> 423  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 423  
gcaugacgac cuaugauga 19

<210> 424  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 424  
caucauaggu cgucaugcu 19

<210> 425  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 425  
agcaugacga ccuaugaug 19

<210> 426  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 426

1189804\_1.TXT  
aucauagguc gucaugcuu 19

<210> 427  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 427  
aagcaugacg accaugau 19

<210> 428  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 428  
ucauaggucg ucaugcuua 19

<210> 429  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 429  
uaagcaugac gaccuauga 19

<210> 430  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 430  
cauaggucgu caugcuuau 19

<210> 431  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 431  
auaagcauga cgaccuaug 19

<210> 432  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 432  
auaggucguc augcuuaug 19

<210> 433  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 433  
cauaagcaug acgaccuau 19

<210> 434  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 434  
uaggucguca ugcuaugg 19

<210> 435  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 435  
ccaauagcau gacgaccua 19

<210> 436  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 436  
aggucgucau gcuuauggg 19

<210> 437  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 437  
cccuaaagca ugacgaccu 19

<210> 438  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 438  
ggucgucaug cuuaugggg 19

<210> 439  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 439  
ccccauaagc augacgacc 19

<210> 440  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 440  
gucgucaugc uuaugggga 19

<210> 441  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 441  
uccccauaag caugacgac 19

<210> 442  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 442  
ucgucaugcu uauggggau 19

<210> 443  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 443  
auccccaaua gcaugacga 19

<210> 444  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 444  
aaccuaucau cauaggucu 19

<210> 445  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 445  
agaccuauga ugauagguu 19

<210> 446  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 446  
 accuaucauc auaggucuu 19

<210> 447  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 447  
 aagaccuau augauaggu 19

<210> 448  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 448  
 ccuaucauca uaggucuuc 19

<210> 449  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 449  
 gaagaccuau gaugauagg 19

<210> 450  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 450  
 cuaucaucau aggucuuca 19

<210> 451  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 451  
 ugaagaccua ugaugauag 19

<210> 452  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 452  
 uaucaucaua ggucuucau 19

<210> 453  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 453  
 augaagaccu augaugaua 19

<210> 454  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 454  
 aucaucauag gucucaug 19

<210> 455  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 455  
 caugaagacc uaugaugau 19



<210> 456  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 456  
ucaucauagg ucuucaugc 19

<210> 457  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 457  
gcaugaagac cuaugauga 19

<210> 458  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 458  
caucauaggu cuucaugcu 19

<210> 459  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 459  
agcaugaaga ccuaugaug 19

<210> 460  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 460  
aucauagguc uucaugcuu 19

<210> 461  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 461  
aagcaugaag accuaugau 19

<210> 462  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 462  
ucauaggucu ucaugcuua 19

<210> 463  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 463  
uaagcaugaa gaccuauga 19

<210> 464  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 464  
cauaggucuu caugcuuau 19

<210> 465  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 105

oligonucleotide

<400> 465  
 auaagcauga agaccuaug 19

<210> 466  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 466  
 auaggucuuc augcuuaug 19

<210> 467  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 467  
 cauaagcaug aagaccuau 19

<210> 468  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 468  
 uaggucuuca ugcuuaugg 19

<210> 469  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 469  
 ccauaagcau gaagaccua 19

<210> 470  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 470  
aggucuucau gcuuauggg 19

<210> 471  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 471  
cccuaagca ugaagaccu 19

<210> 472  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 472  
ggucuucaug cuuaugggg 19

<210> 473  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 473  
ccccauaagc augaagacc 19

<210> 474  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 474  
gucuucaugc uuaugggga 19

<210> 475  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 475

uccccauaag caugaagac

19

<210> 476

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 476

ucuucaugcu uauggggau

19

<210> 477

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 477

auccccaaua gcaugaaga

19

<210> 478

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 478

gugaugagau gacccguau

19

<210> 479

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 479

auacggguca ucucaucac

19

<210> 480

<211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 480  
 gaugagauga cccguauua 19

<210> 481  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 481  
 uauuacgggu caucucauc 19

<210> 482  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 482  
 cguauuaucau ggcaguuca 19

<210> 483  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 483  
 ugaacugcca gauauuacg 19

<210> 484  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 484  
 ggcaguucau caaggagaa 19

1189804\_1.TXT

<210> 485  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 485  
uucuccuuga ugaacugcc 19

<210> 486  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 486  
guguggaaga guucaagcu 19

<210> 487  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 487  
agcuugaacu cuuccacac 19

<210> 488  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 488  
guggaagagu ucaagcuga 19

<210> 489  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 489

ucagcuugaa cucuuccac 1189804\_1.TXT 19

<210> 490  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 490  
gaagaguuca agcugaaga 19

<210> 491  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 491  
ucuucagcuu gaacucuuc 19

<210> 492  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 492  
caguaugcca uccagaaga 19

<210> 493  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 493  
ucuucuggau ggcauacug 19

<210> 494  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide



1189804\_1.TXT

<400> 494  
cuguacauga gcaccaaga 19

<210> 495  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 495  
ucuuggugcu cauguacag 19

<210> 496  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 496  
gcaccaagaa caccauacu 19

<210> 497  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 497  
aguauggugu ucuuggugc 19

<210> 498  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 498  
ccaucugaa agccuacga 19

<210> 499  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 499  
ucguaggcuu ucaguaugg 19

<210> 500  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 500  
cauacugaaa gccuacgau 19

<210> 501  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 501  
aucguaggcu uucaguaug 19

<210> 502  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 502  
guuuaagga caucuucca 19

<210> 503  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 503  
uggaagaugu ccuugaaac 19

<210> 504  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 504  
ccgacuucga caagaauaa 19

<210> 505  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 505  
uuauucuugu cgaagucgg 19

<210> 506  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 506  
gacuucgaca agaauaaga 19

<210> 507  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 507  
ucuuauucuu gucgaaguc 19

<210> 508  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 508  
gacaagaaua agaucuggu 19

<210> 509  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 509  
 accagauuu auucuuguc 19

<210> 510  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 510  
 ggcucuuga ugacauggu 19

<210> 511  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 511  
 accaugucau caaugagcc 19

<210> 512  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 512  
 gcaagaacua ugacggaga 19

<210> 513  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
 <400> 513  
 ucuccucau aguucuugc 19

<210> 514  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 514  
caagaacuau gacggagau 19

<210> 515  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 515  
aucuccguca uaguucuug 19

<210> 516  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 516  
gagaugugca gucagacau 19

<210> 517  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 517  
augucugacu gcacaucuc 19

<210> 518  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 518  
cugaugggaa gacgauuga 19

<210> 519  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 519  
 ucaaucgucu ucccaucag 19

<210> 520  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 520  
 gcaaugugaa gcugaacga 19

<210> 521  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 521  
 ucguucagcu ucacauugc 19

<210> 522  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 522  
 cuguaauuuu uauugcccu 19

<210> 523  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

1189804\_1.TXT

<400> 523  
agggcaauau aaauacag 19

<210> 524  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 524  
cauggugcca uauuuagcu 19

<210> 525  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 525  
agcuauauuau ggcaccaug 19

<210> 526  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 526  
ggugccauau uuagcuacu 19

<210> 527  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 527  
aguagcuaaa uauggcacc 19

<210> 528  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

oligonucleotide

<400> 528  
gugccauauu uagcuacua 19

<210> 529  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 529  
uaguagcuaa auauggcac 19

<210> 530  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 530  
gccauuuua gcuacuaaa 19

<210> 531  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 531  
uuuaguagcu aaauaggc 19

<210> 532  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 532  
gcccacacc auuggcagg 19

<210> 533  
<211> 19  
<212> RNA  
<213> Artificial Sequence



1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 533  
ccugccaug gugauggc 19

<210> 534  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 534  
cccaucacca uggcaggc 19

<210> 535  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 535  
gccugccaau ggugaugg 19

<210> 536  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 536  
ccaucaccau uggcaggca 19

<210> 537  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 537  
ugccugccaa uggugaugg 19

<210> 538  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 538

caucaccauu ggcaggcac

19

<210> 539

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 539

gugccugcca auggugaug

19

<210> 540

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 540

aucaccauug gcaggcacg

19

<210> 541

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 541

cgugccugcc aauggugau

19

<210> 542

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 542

ucaccauugg caggcacgc

19

<210> 543

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 543  
gcgugccugc caaugguga 19

<210> 544  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 544  
caccauuggc aggcacgcc 19

<210> 545  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 545  
ggcgugccug ccaauggug 19

<210> 546  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 546  
accuauuggca ggcacgccc 19

<210> 547  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 547  
gggugugccu gccaauggu 19

<210> 548  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 548  
 ccauuggcag gcacgccca 19

<210> 549  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 549  
 ugggcgugcc ugccaugg 19

<210> 550  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 550  
 cauuggcagg cacgcccau 19

<210> 551  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 551  
 augggcgugc cugccaug 19

<210> 552  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 552

1189804\_1.TXT  
auuggcaggc acgccaug 19

<210> 553  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 553  
caugggcgug ccugccaau 19

<210> 554  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 554  
uuggcaggca cgccaugg 19

<210> 555  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 555  
ccaugggcgu gccugccaa 19

<210> 556  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 556  
uggcaggcac gcccauggc 19

<210> 557  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 557  
gccaugggcg ugccugcca 19

<210> 558  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 558  
ggcaggcacg cccauggcg 19

<210> 559  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 559  
cgccaugggc gugccugcc 19

<210> 560  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 560  
gcaggcacgc ccauggcga 19

<210> 561  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 561  
ucgccauggg cgugccugc 19

<210> 562  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 562  
 caggcacgcc cauggcgac 19

<210> 563  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 563  
 gucgccaugg gcgugccug 19

<210> 564  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 564  
 aggcacgccc auggcgacc 19

<210> 565  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 565  
 ggucgccaug ggcgugccu 19

<210> 566  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 566  
 gcccaucacc auuggcggg 19

<210> 567  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 567  
cccgccaaug gugaugggc 19

<210> 568  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 568  
cccaucacca uuggcgggc 19

<210> 569  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 569  
gcccgccaa uggugauggg 19

<210> 570  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 570  
ccaucaccau uggcgggca 19

<210> 571  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 571  
ugcccgcaa uggugaugg 19

<210> 572  
<211> 19



<212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
         oligonucleotide  
  
 <400> 572  
 caucaccauu ggcgggcac 19

<210> 573  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
         oligonucleotide  
  
 <400> 573  
 gugcccgcca auggugaug 19

<210> 574  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
         oligonucleotide  
  
 <400> 574  
 aucaccauug gcgggcacg 19

<210> 575  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
         oligonucleotide  
  
 <400> 575  
 cgugcccgcc auggugau 19

<210> 576  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
         oligonucleotide  
  
 <400> 576  
 ucaccauugg cgggcacgc 19

<210> 577  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 577  
gcgugcccgc caaugguga 19

<210> 578  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 578  
caccauuggc gggcacgcc 19

<210> 579  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 579  
ggcgugcccg ccaauggug 19

<210> 580  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 580  
accauuggcg ggcacgccc 19

<210> 581  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 581  
ggcgugccc gccaauggu 19

<210> 582  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 582  
ccauggcg gcacgccca 19

<210> 583  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 583  
ugggcgugcc cgccaugg 19

<210> 584  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 584  
cauggcggg cacgcccau 19

<210> 585  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 585  
augggcgugc ccgccaug 19

<210> 586  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 586  
auuggcgggc acgccaug 19

<210> 587  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 587  
cauggcgug cccgccaau 19

<210> 588  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 588  
uuggcgggca cgccaugg 19

<210> 589  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 589  
ccauggcggu gcccgcaa 19

<210> 590  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 590  
uggcgggcac gcccauggc 19

<210> 591  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 131

oligonucleotide

<400> 591  
gccaugggcg ugcccgcca 19

<210> 592  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 592  
ggcgggcacg cccauggcg 19

<210> 593  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 593  
cgccaugggc gugcccgcc 19

<210> 594  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 594  
gcgggcacgc ccauggcga 19

<210> 595  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 595  
ucgccauggg cgugcccgc 19

<210> 596  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 596  
cgggcacgcc cauggcgac 19

<210> 597  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 597  
gucgccaugg gcgugccc 19

<210> 598  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 598  
gggcacgccc auggcgacc 19

<210> 599  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 599  
ggucgccaug ggcgugccc 19

<210> 600  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 600  
gcccacacc auuggcugg 19

<210> 601  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 601

ccagccaug gugaugggc

19

<210> 602

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 602

cccaucacca uggcuggc

19

<210> 603

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 603

gccagccaau ggugauggg

19

<210> 604

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 604

ccaucaccau uggcuggca

19

<210> 605

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 605

ugccagccaa uggugaugg

19

<210> 606

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 606  
cauccauu ggcuggcac 19

<210> 607  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 607  
gugccagcca auggugaug 19

<210> 608  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 608  
aucaccauug gcuggcacg 19

<210> 609  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 609  
cgugccagcc auggugau 19

<210> 610  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 610  
ucaccauugg cuggcacgc 19



<210> 611  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 611  
 gcgugccagc caaugguga 19

<210> 612  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 612  
 caccauuggc uggcacgcc 19

<210> 613  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 613  
 ggcgugccag ccaauggug 19

<210> 614  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 614  
 accauuggcu ggcacgccc 19

<210> 615  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 615

1189804\_1.TXT  
gggcgugcca gccaauggu 19

<210> 616  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 616  
ccauggcug gcacgccca 19

<210> 617  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 617  
ugggcgugcc agccaaugg 19

<210> 618  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 618  
cauggcugg cacgcccau 19

<210> 619  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 619  
augggcgugc cagccaug 19

<210> 620  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

1189804\_1.TXT

<400> 620  
auuggcuggc acgccaug 19

<210> 621  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 621  
caugggcgug ccagccaau 19

<210> 622  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 622  
uuggcuggca cgccaugg 19

<210> 623  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 623  
ccaugggcgu gccagccaa 19

<210> 624  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 624  
uggcuggcac gcccauggc 19

<210> 625  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 625  
gccaugggcg ugccagcca 19

<210> 626  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 626  
ggcuggcacg cccauggcg 19

<210> 627  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 627  
cgccaugggc gugccagcc 19

<210> 628  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 628  
gcuggcacgc ccauggcga 19

<210> 629  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 629  
ucgccauggg cgugccagc 19

<210> 630  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 630  
cuggcacgcc cauggcgac 19

<210> 631  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 631  
gucgccaugg gcgugccag 19

<210> 632  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 632  
uggcacgccc auggcgacc 19

<210> 633  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 633  
ggucgccaug ggcgugcca 19

<210> 634  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 634  
gcccaucacc auuggcaag 19

<210> 635  
<211> 19

<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 635  
cuugccaug gugauggc 19

<210> 636  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 636  
cccaucacca uuggcaagc 19

<210> 637  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 637  
gcuugccaau ggugaugg 19

<210> 638  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 638  
ccaucaccau uggcaagca 19

<210> 639  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 639  
ugcuugccaa uggugaugg 19

<210> 640  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 640  
caucaccauu ggcaagcac 19

<210> 641  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 641  
gugcuugcca auggugaug 19

<210> 642  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 642  
aucaccauug gcaagcacg 19

<210> 643  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 643  
cgugcuugcc auggugau 19

<210> 644  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 644  
ucaccauugg caagcacg 19

<210> 645  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 645  
gcgugcuugc caaugguga 19

<210> 646  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 646  
caccauuggc aagcacgcc 19

<210> 647  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 647  
ggcgugcuug ccaauggug 19

<210> 648  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 648  
accuauuggca agcacgccc 19

<210> 649  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide



<400> 649  
gggcgugcuu gccaauggu 19

<210> 650  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 650  
ccauggcaa gcacgccca 19

<210> 651  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 651  
ugggcgugcu ugccaugg 19

<210> 652  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 652  
cauggcaag cacgcccau 19

<210> 653  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 653  
augggcgugc uugccaug 19

<210> 654  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 144

oligonucleotide

<400> 654  
 auuggcaagc acgccaug 19

<210> 655  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 655  
 cauggcgug cuugccaau 19

<210> 656  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 656  
 uuggcaagca cgccaugg 19

<210> 657  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 657  
 ccauggcggu gcuugccaa 19

<210> 658  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 658  
 uggcaagcac gcccauggc 19

<210> 659  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 659  
gccaugggcg ugcuugcca 19

<210> 660  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 660  
ggcaagcacg cccauggcg 19

<210> 661  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 661  
cgccaugggc gugcuugcc 19

<210> 662  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 662  
gcaagcacgc ccauggcga 19

<210> 663  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 663  
ucgccauggg cgugcuugc 19

<210> 664  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 664

caagcacgcc cauggcgac

19

<210> 665

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 665

gucgccaugg gcgugcuug

19

<210> 666

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 666

aagcacgcc auggcgacc

19

<210> 667

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 667

ggucgccaug ggcgugcuu

19

<210> 668

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 668

gcccacacc auuggcacg

19

<210> 669

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 669  
cgugccaug gugauggc 19

<210> 670  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 670  
cccaucacca uggcagc 19

<210> 671  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 671  
gcgugccaau ggugaugg 19

<210> 672  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 672  
ccaucaccau uggcagca 19

<210> 673  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 673  
ugcgugccaa uggugaugg 19

1189804\_1.TXT

<210> 674  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 674  
caucaccauu ggcacgcac 19

<210> 675  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 675  
gugcgugcca auggugaug 19

<210> 676  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 676  
aucaccauug gcacgcacg 19

<210> 677  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 677  
cgugcgugcc auggugau 19

<210> 678  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 678

ucaccauugg cacgcacgc 1189804\_1.TXT 19

<210> 679  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 679  
gcgugcgugc caaugguga 19

<210> 680  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 680  
caccauuggc acgcacgcc 19

<210> 681  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 681  
ggcgugcgug ccaauuggug 19

<210> 682  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 682  
accauuggca cgcacgccc 19

<210> 683  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 683  
gggcgugcgu gccaauggu 19

<210> 684  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 684  
ccauggcac gcacgccca 19

<210> 685  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 685  
ugggcgugcg ugccaugg 19

<210> 686  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 686  
cauggcacg cacgcccau 19

<210> 687  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 687  
augggcgugc gugccaug 19

<210> 688  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>



<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 688  
auuggcacgc acgcccaug 19

<210> 689  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 689  
caugggcgug cgugccaau 19

<210> 690  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 690  
uuggcacgca cgcccaugg 19

<210> 691  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 691  
ccaugggcgu gcgugccaa 19

<210> 692  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 692  
uggcacgcac gcccauggc 19

<210> 693  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 693  
gccaugggcg ugcgugcca 19

<210> 694  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 694  
ggcacgcacg cccauggcg 19

<210> 695  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 695  
cgccaugggc gugcgugcc 19

<210> 696  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 696  
gcacgcacgc ccauggcga 19

<210> 697  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 697  
ucgccauggg cgugcgugc 19

<210> 698  
<211> 19

<212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 698  
 cacgcacgcc cauggcgac 19

<210> 699  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 699  
 gucgccaugg gcgugcgug 19

<210> 700  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 700  
 acgcacgccc auggcgacc 19

<210> 701  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 701  
 ggucgccaug ggcgugcgu 19

<210> 702  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 702  
 gcccaucacc auuggcaug 19

<210> 703  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 703  
caugccaug gugauggc 19

<210> 704  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 704  
cccaucacca uuggcaugc 19

<210> 705  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 705  
gcaugccaau ggugaugg 19

<210> 706  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 706  
ccaucaccau uggcaugca 19

<210> 707  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide  
  
<400> 707  
ugcaugccaa uggugaugg 19

<210> 708  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 708  
caucaccauu ggcaugcac 19

<210> 709  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 709  
gugcaugcca auggugaug 19

<210> 710  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 710  
aucaccauug gcaugcacg 19

<210> 711  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 711  
cgugcaugcc aauggugau 19

<210> 712  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 712  
ucaccauugg caugcacgc 19

<210> 713  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 713  
gcgugcaugc caaugguga 19

<210> 714  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 714  
caccauuggc augcacgcc 19

<210> 715  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 715  
ggcgugcaug ccaauggug 19

<210> 716  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 716  
accauuggca ugcacgccc 19

<210> 717  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 157

oligonucleotide

<400> 717  
 gggcgugcau gccaauggu 19

<210> 718  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 718  
 ccauggcau gcacgccca 19

<210> 719  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 719  
 uggcgugca ugccaugg 19

<210> 720  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 720  
 cauggcaug cacgcccau 19

<210> 721  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide

<400> 721  
 auggcgugc augccaug 19

<210> 722  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 722  
auuggcaugc acgccaug 19

<210> 723  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 723  
cauggcgug caugccaau 19

<210> 724  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 724  
uuggcaugca cgccaugg 19

<210> 725  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 725  
ccauggcggu gcaugccaa 19

<210> 726  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 726  
uggcaugcac gcccauggc 19

<210> 727  
<211> 19  
<212> RNA



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 727

gcccaugggcg ugcaugcca

19

<210> 728

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 728

ggcaugcacg cccauggcg

19

<210> 729

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 729

cgccaugggc gugcaugcc

19

<210> 730

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 730

gcaugcacgc ccauggcga

19

<210> 731

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 731

ucgccauggg cgugcaugc

19

<210> 732

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 732  
caugcacgcc cauggcgac 19

<210> 733  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 733  
gucgccaugg gcgugcaug 19

<210> 734  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 734  
augcacgccc auggcgacc 19

<210> 735  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 735  
ggucgccaug ggcgugcau 19

<210> 736  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 736  
gcccacacc auuggcagc 19

1189804\_1.TXT

<210> 737  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 737  
gcugccaug gugauggc 19

<210> 738  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 738  
cccaucacca uggcagcc 19

<210> 739  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 739  
ggcugccaau ggugaugg 19

<210> 740  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 740  
ccaucaccau uggcagcca 19

<210> 741  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 741

1189804\_1.TXT  
uggcugccaa uggugaugg 19

<210> 742  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 742  
caucaccauu ggcagccac 19

<210> 743  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 743  
guggcugcca auggugaug 19

<210> 744  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 744  
aucaccauug gcagccacg 19

<210> 745  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 745  
cguggcugcc auggugau 19

<210> 746  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

1189804\_1.TXT

<400> 746  
ucaccaugg cagccacgc 19

<210> 747  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 747  
gcgggcugc caaugguga 19

<210> 748  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 748  
caccauggc agccacgcc 19

<210> 749  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 749  
ggcgggcug ccaauggug 19

<210> 750  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 750  
accauggca gccacgccc 19

<210> 751  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 751  
gggcguggcu gccaauggu 19

<210> 752  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 752  
ccauggcag ccacgccca 19

<210> 753  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 753  
ugggcguggc ugccaugg 19

<210> 754  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 754  
cauggcagc cacgcccau 19

<210> 755  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 755  
augggcgugg cugccaug 19

<210> 756  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 756  
auuggcagcc acgccaug 19

<210> 757  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 757  
caugggcgug gcugccaau 19

<210> 758  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 758  
uuggcagcca cgccaugg 19

<210> 759  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 759  
ccaugggcgu ggugccaa 19

<210> 760  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 760  
uggcagccac gcccauggc 19

<210> 761  
<211> 19

<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 761  
gccaugggcg uggcugcca 19

<210> 762  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 762  
ggcagccacg cccauggcg 19

<210> 763  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 763  
cgccaugggc guggcugcc 19

<210> 764  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 764  
gcagccacgc ccauggcga 19

<210> 765  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 765  
ucgccauggg cguggcugc 19



<210> 766  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide  
  
<400> 766  
cagccacgcc cauggcgac 19

<210> 767  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide  
  
<400> 767  
gucgccaugg gcguggcug 19

<210> 768  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide  
  
<400> 768  
agccacgccc auggcgacc 19

<210> 769  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide  
  
<400> 769  
ggucgccaug ggcguggcu 19

<210> 770  
<211> 19  
<212> RNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide  
  
<400> 770  
gcccaucacc auuggcagu 19

<210> 771  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 771  
acugccaaug gugaugggc 19

<210> 772  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 772  
cccaucacca uggcaguc 19

<210> 773  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 773  
gacugccaau ggugauggg 19

<210> 774  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 774  
ccaucaccau uggcaguca 19

<210> 775  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 775  
ugacugccaa uggugaugg 19

<210> 776  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 776  
caucaccauu ggcagucac 19

<210> 777  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 777  
gugacugcca auggugaug 19

<210> 778  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 778  
aucaccauug gcagucacg 19

<210> 779  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
oligonucleotide

<400> 779  
cgugacugcc auggugau 19

<210> 780  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Page 170

oligonucleotide

<400> 780  
ucaccauugg cagucacgc 19

<210> 781  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 781  
gcgugacugc caaugguga 19

<210> 782  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 782  
caccuuggc agucacgcc 19

<210> 783  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 783  
ggcgugacug ccauggug 19

<210> 784  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 784  
accuuggca gucacgccc 19

<210> 785  
<211> 19  
<212> RNA  
<213> Artificial Sequence

1189804\_1.TXT

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 785  
gggcgugacu gccaauggu 19

<210> 786  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 786  
ccauggcag ucacgccca 19

<210> 787  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 787  
ugggcgugac ugccaugg 19

<210> 788  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 788  
cauggcagu cacgcccau 19

<210> 789  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 789  
augggcguga cugccaug 19

<210> 790  
<211> 19  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 790

auuggcaguc acgccaug

19

<210> 791

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 791

cauggcgug acugccaau

19

<210> 792

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 792

uuggcaguca cgccaugg

19

<210> 793

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 793

ccauggcgug gacugccaa

19

<210> 794

<211> 19

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 794

uggcagucac gcccauggc

19

<210> 795

<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 795  
gccaugggcg ugacugcca 19

<210> 796  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 796  
ggcagucacg cccauggcg 19

<210> 797  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 797  
cgccaugggc gugacugcc 19

<210> 798  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 798  
gcagucacgc ccauggcga 19

<210> 799  
<211> 19  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 799  
ucgccauggg cgugacugc 19

1189804\_1.TXT

<210> 800  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 800  
 cagucacgcc cauggcgac 19  
  
 <210> 801  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 801  
 gucgccaugg gcgugacug 19  
  
 <210> 802  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 802  
 agucacgccc auggcgacc 19  
  
 <210> 803  
 <211> 19  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
 oligonucleotide  
  
 <400> 803  
 ggucgccaug ggcgugacu 19  
  
 <210> 804  
 <211> 489  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 804  
 gcataatgag ctctatatgc catcactgca gttgtaggtt ataactatcc attgtctga 60  
 aaaactttgc ttctaatttt tctctttcaa gctatgattt aggcatagag aatcgtgatg 120



1189804\_1.TXT

ccaccaacga ccaagtcacc aaggatgctg cagaagctat aaagaagcat aatgttggcg	180
tcaaatgtgc cactatcact cctgatgaga agagggttga ggagttcaag ttgaaacaaa	240
tgtggaaatc accaaatggc accatacgaa atattctggg tggcacggtc ttcagagaag	300
ccattatctg caaaaatata ccccggttg tgagtggatg ggtaaacct atcatcatag	360
gtcgtcatgc ttatgggat caagtaagtc atgttgcaa taatgtgatt ttgcatgbtg	420
gccagaaat ttccaacttg tatgtgtttt attcttatct tttggtatct acaccatta	480
agcaaggta	489



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 4 columns: APPLICATION NUMBER (15/589,615), FILING OR 371(C) DATE (05/08/2017), FIRST NAMED APPLICANT (Leonard Luan C), ATTY. DOCKET NO./TITLE (AGS-013C2)

148106
GOODWIN PROCTER LLP
PATENT ADMINISTRATOR
100 Northern Avenue
BOSTON, MA 02210

CONFIRMATION NO. 7017
FORMALITIES LETTER



Date Mailed: 05/24/2017

NOTICE TO FILE MISSING PARTS OF NONPROVISIONAL APPLICATION

FILED UNDER 37 CFR 1.53(b)

Filing Date Granted

Items Required To Avoid Abandonment:

An application number and filing date have been accorded to this application. The item(s) indicated below, however, are missing. Applicant is given TWO MONTHS from the date of this Notice within which to file all required items below to avoid abandonment. Extensions of time may be obtained by filing a petition accompanied by the extension fee under the provisions of 37 CFR 1.136(a).

The application is informal since it does not comply with the regulations for the reason(s) indicated below.

The required item(s) identified below must be timely submitted to avoid abandonment:

- A substitute specification in compliance with 37 CFR 1.52, 1.121(b)(3), and 1.125, is required. The substitute specification must be submitted with markings and be accompanied by a clean version (without markings) as set forth in 37 CFR 1.125(c) and a statement that the substitute specification contains no new matter (see 37 CFR 1.125(b)). The specification, claims, and/or abstract page(s) submitted is not acceptable and cannot be scanned or properly stored because:
- The application papers (including any electronically submitted papers) are not in compliance with 37 CFR 1.52 because pages 179-183 contain text that is unreadable or of insufficient clarity. Application papers (including any electronically submitted papers) must be presented in a form having sufficient clarity and contrast between the background of the paper and the writing thereon to permit the Office to electronically reproduce the papers by use of digital imaging and optical character recognition. See 37 CFR 1.52(a)(1)(v).
- This application does not contain, as a separate part of the disclosure on paper copy, a "Sequence Listing" as required by 37 CFR 1.821(c) Applicant must provide an initial paper or compact disc copy of the "Sequence Listing", as well as an amendment specifically directing its entry into the application and a statement that the content of the sequence listing information recorded in computer readable form is identical to the written (on paper or compact disc) sequence listing and, where applicable, includes no new matter, as required by 37 CFR 1.821(e), 1.821(f), 1.821(g), 1.825(b), or 1.825(d). If the effective filing date is on or after September 8, 2000, see the final rulemaking notice published in the Federal Register at 65 FR 54604 (September 8, 2000) and 1238 OG 145 (September 19, 2000).
- A copy of the "Sequence Listing" in computer readable form has not been submitted as required by 37 CFR 1.821(e). If the effective filing date is on or after September 8, 2000, see the final rulemaking notice

published in the Federal Register at 65 FR 54604 (September 8, 2000) and 1238 OG 145 (September 19, 2000). Applicant must provide an initial computer readable form (CRF) copy of the "Sequence Listing" and a statement that the content of the sequence listing information recorded in computer readable form is identical to the written (on paper or compact disc) sequence listing and, where applicable, includes no new matter, as required by 37 CFR 1.821(e), 1.821(f), 1.821(g), 1.825(b), or 1.825(d). If applicant desires the sequence listing in the instant application to be identical with that of another application on file in the U.S. Patent and Trademark Office, such request in accordance with 37 CFR 1.821(e) may be submitted in lieu of a new CRF.

**For questions regarding compliance to these requirements, please contact:**

- **For Rules interpretation, call (571) 272-2510**
- **For CRF submission help, call (571) 272-2510**
- **For PatentIn software program support technical assistance, call (866)217-9197.**
- **Send e-mail correspondence for PatentIn software program help to [ebc@uspto.gov](mailto:ebc@uspto.gov)**
- **Patent software is available at <http://www.uspto.gov/patents/resources/tools/checker/patentinrel.jsp> .**

Applicant is cautioned that correction of the above items may cause the specification and drawings page count to exceed 100 pages. If the specification and drawings exceed 100 pages, applicant will need to submit the required application size fee.

**SUMMARY OF FEES DUE:**

The fee(s) required within **TWO MONTHS** from the date of this Notice to avoid abandonment is/are itemized below. No entity status discount is in effect. If applicant is qualified for small entity status, a written assertion of small entity status must be submitted to establish small entity status. (See 37 CFR 1.27). If applicant is qualified for micro entity status, an acceptable Certification of Micro Entity Status must be submitted to establish micro entity status. (See 37 CFR 1.29 and forms PTO/SB/15A and 15B.)

- **\$ 800 for 75 electronically equivalent pages in excess of 100 application size fee.**
- **\$( 0) previous unapplied payment amount.**
- **\$ 800 TOTAL FEE BALANCE DUE.**

**Items Required To Avoid Processing Delays:**

Applicant is notified that the above-identified application contains the deficiencies noted below. No period for reply is set forth in this notice for correction of these deficiencies. However, if a deficiency relates to the inventor's oath or declaration, the applicant must file an oath or declaration in compliance with 37 CFR 1.63, or a substitute statement in compliance with 37 CFR 1.64, executed by or with respect to each actual inventor no later than the expiration of the time period set in the "Notice of Allowability" to avoid abandonment. See 37 CFR 1.53(f).

- A properly executed inventor's oath or declaration has not been received for the following inventor(s):  
Leonard Luan C

Replies must be received in the USPTO within the set time period or must include a proper Certificate of Mailing or Transmission under 37 CFR 1.8 with a mailing or transmission date within the set time period. For more information and a suggested format, see Form PTO/SB/92 and MPEP 512.

Replies should be mailed to:

Mail Stop Missing Parts  
Commissioner for Patents  
P.O. Box 1450  
Alexandria VA 22313-1450

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web, including a copy of this Notice and selecting the document description "Applicant response to Pre-Exam Formalities Notice".  
<https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at 1-866-217-9197 or visit our website at <http://www.uspto.gov/ebc>.

If you are not using EFS-Web to submit your reply, you must include a copy of this notice.

Questions about the contents of this notice and the requirements it sets forth should be directed to the Office of Data Management, Application Assistance Unit, at **(571) 272-4000** or **(571) 272-4200** or **1-888-786-0101**.

/sareebuddin/

---

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

---

In re Patent Application of:  
Lenny Dang et al.

Application No.: 15/589,615

Confirmation No.: 7017

Filed: May 8, 2017

Art Unit: N/A

For: METHODS AND COMPOSITIONS FOR  
CELL-PROLIFERATION-RELATED  
DISORDERS

---

Examiner: Not Yet Assigned

**REQUEST TO CORRECT INVENTORSHIP UNDER 37 C.F.R. § 1.48(d) AND FOR  
CORRECTED FILING RECEIPT**

Office of Initial Patent Examination's Filing Receipt Corrections  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Dear Sir:

Applicant hereby requests that a corrected Filing Receipt be issued in the above-identified patent application. Filed concurrently herewith is (1) a marked up Application Data Sheet specifying the inventorship and Applicant of the application, and (2) a marked up Filing Receipt. The error in inventorship occurred without any deceptive intent on the part of the inventors.

Changes are shown by underlining for added matter and ~~striketrough~~ for deleted matter. Please update the following information on the Filing Receipt.

Inventor(s)

Lenny Dang ~~Leonard Luan C~~, Boston, MA;

Valeria Fantin, Buringame, CA;

Stefan Gross, Brookline, MA;

Hyun ~~G.~~ Gyung Jang, Waltham, MA;

Shengfang Jin, Newton, MA;

Francesco G. Salituro, Marlborough, MA;

Jeffrey O. Saunders, Lincoln, MA;

Shin-San M. Su, Boston, MA;

Katharine Yen, Wellesley, MA;

Applicant(s)

Agios Pharmaceuticals, Inc., Cambridge, MA

~~Leonard Luan C, Boston, MA;~~

~~Valeria Fantin, Buringame, CA;~~

~~Stefan Gross, Brookline, MA;~~

~~Hyun G. Jang, Waltham, MA;~~

~~Shengfang Jin, Newton, MA;~~

~~Francesco G. Salituro, Marlborough, MA;~~

~~Jeffrey O. Saunders, Lincoln, MA;~~

~~Shin-San M. Su, Boston, MA;~~

~~Katharine Yen, Wellesley, MA;~~

Thus the Corrected Filing Receipt should read as follows:

Inventor(s)

Lenny Dang, Boston, MA;

Valeria Fantin, Buringame, CA;

Stefan Gross, Brookline, MA;

Hyun Gyung Jang, Waltham, MA;

Shengfang Jin, Newton, MA;

Francesco G. Salituro, Marlborough, MA;

Jeffrey O. Saunders, Lincoln, MA;

Shin-San M. Su, Boston, MA;

Katharine Yen, Wellesley, MA;

Applicant(s)

Agios Pharmaceuticals, Inc., Cambridge, MA

Furthermore, Applicant additionally requests that all pertinent U.S. Patent and Trademark Office records relating to the subject application be changed to reflect this correction.

The processing fee of \$140.00 as set forth in 37 C.F.R. § 1.17(i) is submitted herewith. However, if a further fee is due, please charge our Deposit Account No. 07-1700, under Order No. AGS-013C2 from which the undersigned is authorized to draw.

Dated: July 20, 2017

Respectfully submitted,

Electronic signature: /Ioana Davies/  
Ioana Davies

Registration No.: 75,817  
GOODWIN PROCTER LLP  
100 Northern Avenue  
Boston, Massachusetts 02210  
(617) 570-3920

**Corrected Application Data Sheet**

**Inventor Information**

Inventor Number:: 1  
Given Name:: ~~Leonard Luan~~ Lenny  
Family Name:: ~~G. Dang~~  
Suffix:: Dang  
City of Residence:: Boston  
State or Province of Residence:: MA  
Country of Residence:: US  
Street of mailing address:: 30 Union Street, #201  
City of mailing address:: Boston  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02118

Inventor Number:: 2  
Given Name:: Valeria  
Family Name:: Fantin  
City of Residence:: Burlingame  
State or Province of Residence:: CA  
Country of Residence:: US  
Street of mailing address:: 3133 Frontera Way, Apt. 217  
City of mailing address:: Burlingame



State or Province of mailing address:: CA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 94010

Inventor Number:: 3  
Given Name:: Stefan  
Family Name:: Gross  
City of Residence:: Brookline  
State or Province of Residence:: MA  
Country of Residence:: US  
Street of mailing address:: 118 Addington Road #2  
City of mailing address:: Brookline  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02445

Inventor Number:: 4  
Given Name:: Hyun  
Middle Name:: ~~G.~~Gyung  
Family Name:: Jang  
City of Residence:: Waltham  
State or Province of Residence:: MA  
Country of Residence:: US

Street of mailing address:: 37 Albemarle Road  
City of mailing address:: Waltham  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02452

Inventor Number:: 5  
Given Name:: Shengfang  
Family Name:: Jin  
City of Residence:: Newton  
State or Province of Residence:: MA  
Country of Residence:: US

Street of mailing address:: 6 Audubon Drive  
City of mailing address:: Newton  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02467

Inventor Number:: 6  
Given Name:: Francesco  
Middle Name:: G.  
Family Name:: Salituro  
City of Residence:: Marlborough

State or Province of Residence:: MA  
Country of Residence:: US  
Street of mailing address:: 25 Baker Drive  
City of mailing address:: Marlborough  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 01752

Inventor Number:: 7  
Given Name:: Jeffrey  
Middle Name:: O.  
Family Name:: Saunders  
City of Residence:: Lincoln  
State or Province of Residence:: MA  
Country of Residence:: US  
Street of mailing address:: 188 Tower Road  
City of mailing address:: Lincoln  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 01773

Inventor Number:: 8  
Given Name:: Shin-San

Middle Name:: M- Michael  
Family Name:: Su  
City of Residence:: Boston  
State or Province of Residence:: MA  
Country of Residence:: US  
Street of mailing address:: 128 Beacon Street, Unit F  
City of mailing address:: Boston  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02116

Inventor Number:: 9  
Given Name:: Katharine  
Family Name:: Yen  
City of Residence:: Wellesley  
State or Province of Residence:: MA  
Country of Residence:: US  
Street of mailing address:: 7 Norwich Rd.  
City of mailing address:: Wellesley  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02481

## Application Information

Application Number::	<u>15/589,615</u>
Filing Date::	<u>05/08/17</u>
Application Type::	Regular
Subject Matter::	Utility
CD-ROM or CD-R?::	None
Sequence submission?::	None
Computer Readable Form (CRF)?::	No
Title::	METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION- RELATED DISORDERS
Attorney Docket Number::	AGS-013C2
Request for Early Publication?::	No
Request for Non-Publication?::	No
Small Entity?::	No
Petition included?::	No
Portions or all of the application associated with this Application Data Sheet may fall under a Secrecy Order pursuant to 37 CFR 5.2::	No
This application (1) claims priority to or the benefit of an application filed before March 16, 2013 and (2) also contains, or contained at any time, a claim to a claimed invention that has an effective filing date on or after March 16, 2013::	No

**Representative Information**

Representative Customer Number:: 148106

**Applicant Information**

Applicant Number:: 1  
Applicant Type:: Assignee  
Organization Name:: Agios Pharmaceuticals, Inc.  
Street of mailing address:: 88 Sidney Street  
City of mailing address:: Cambridge  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02139

Applicant Number:: 4  
Given Name:: Leonard Luan Lenny  
Family Name:: C. Dang  
Suffix:: Dang  
Applicant Type:: Joint Inventor  
Street of mailing address:: ~~30 Union Street, #201~~  
City of mailing address:: Boston  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02118

Applicant Number:: 2  
Given Name:: Valeria  
Family Name:: Fantin  
Applicant Type:: Joint Inventor  
Street of mailing address:: ~~3133 Frontera Way, Apt. 217~~  
City of mailing address:: Burlingame  
State or Province of mailing address:: CA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 94010

Applicant Number:: 3  
Given Name:: Stefan  
Family Name:: Gross  
Applicant Type:: Joint Inventor  
Street of mailing address:: ~~118 Addington Road #2~~  
City of mailing address:: Brookline  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02445

Applicant Number:: 4  
Given Name:: Hyun  
Middle Name:: G. Gyung

Family Name:: Jang  
Applicant Type:: Joint Inventor  
Street of mailing address:: ~~37 Albemarle Road~~  
City of mailing address:: ~~Waltham~~  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02452

Applicant Number:: 5  
Given Name:: Shengfang  
Family Name:: Jin  
Applicant Type:: Joint Inventor  
Street of mailing address:: ~~6 Audubon Drive~~  
City of mailing address:: ~~Newton~~  
State or Province of mailing address:: MA  
Country of mailing address:: US  
Postal or Zip Code of mailing address:: 02467

Applicant Number:: 6  
Given Name:: Francesco  
Middle Name:: G.  
Family Name:: Salituro  
Applicant Type:: Joint Inventor



Street of mailing address:: ~~25 Baker Drive~~  
City of mailing address:: ~~Marlborough~~  
State or Province of mailing address:: ~~MA~~  
Country of mailing address:: ~~US~~  
Postal or Zip Code of mailing address:: ~~01752~~

Applicant Number:: ~~7~~  
Given Name:: ~~Jeffrey~~  
Middle Name:: ~~O.~~  
Family Name:: ~~Saunders~~  
Applicant Type:: ~~Joint Inventor~~

Street of mailing address:: ~~188 Tower Road~~  
City of mailing address:: ~~Lincoln~~  
State or Province of mailing address:: ~~MA~~  
Country of mailing address:: ~~US~~  
Postal or Zip Code of mailing address:: ~~01773~~

Applicant Number:: ~~8~~  
Given Name:: ~~Shin-San~~  
Middle Name:: ~~M. Michael~~  
Family Name:: ~~Su~~  
Applicant Type:: ~~Joint Inventor~~  
Street of mailing address:: ~~128 Beacon Street, Unit F~~

City of mailing address:: Boston  
 State or Province of mailing address:: MA  
 Country of mailing address:: US  
 Postal or Zip Code of mailing address:: 02116

Applicant Number:: 9  
 Given Name:: Katharine  
 Family Name:: Yen  
 Applicant Type:: Joint Inventor  
 Street of mailing address:: ~~7 Norwich Rd.~~  
 City of mailing address:: Wellesley  
 State or Province of mailing address:: MA  
 Country of mailing address:: US  
 Postal or Zip Code of mailing address:: 02481

**NOTE:** This Application Data Sheet must be signed in accordance with 37 CFR 1.33(b). **However, if this Application Data Sheet is submitted with the INITIAL filing of the application and either box A or B is not checked in subsection 2 of the "Authorization or Opt-Out of Authorization to Permit Access" section, then this form must also be signed in accordance with 37 CFR 1.14(c).**

This Application Data Sheet **must** be signed by a patent practitioner if one or more of the applicants is a **juristic entity** (e.g., corporation or association). If the applicant is two or more joint inventors, this form must be signed by a patent practitioner, **all** joint inventors who are the applicant, or one or more joint inventor-applicants who have been given power of attorney (e.g., see USPTO Form PTO/AIA/81) on behalf of **all** joint inventor-applicants.

See 37 CFR 1.4(d) for the manner of making signatures and certifications.

Signature	/Ioana Davies/	Date (YYYY-MM-DD)	2017-07-20
Name	Ioana Davies	Registration Number	75,817



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 7 columns: APPLICATION NUMBER, FILING or 371(c) DATE, GRP ART UNIT, FIL FEE REC'D, ATTY DOCKET NO, TOT CLAIMS, IND CLAIMS. Row 1: 15/589,615, 05/08/2017, 1740, AGS-013C2, 18, 1

CONFIRMATION NO. 7017

FILING RECEIPT

148106
GOODWIN PROCTER LLP
PATENT ADMINISTRATOR
100 Northern Avenue
BOSTON, MA 02210



Date Mailed: 05/24/2017

Receipt is acknowledged of this non-provisional patent application. The application will be taken up for examination in due course. Applicant will be notified as to the results of the examination. Any correspondence concerning the application must include the following identification information: the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please submit a written request for a Filing Receipt Correction. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections

Inventor(s)

- Lenny Dang Leonard Luan C, Boston, MA;
Valeria Fantin, Burlingame, CA;
Stefan Gross, Brookline, MA;
Hyun Gyung Hyun G. Jang, Waltham, MA;
Shengfang Jin, Newton, MA;
Francesco G. Salituro, Marlborough, MA;
Jeffrey O. Saunders, Lincoln, MA;
Shin-San M. Su, Boston, MA;
Katharine Yen, Wellesley, MA;

Applicant(s)

- ~~Leonard Luan C, Boston, MA;~~ Agios Pharmaceuticals, Inc., Cambridge, MA
~~Valeria Fantin, Burlingame, CA;~~
~~Stefan Gross, Brookline, MA;~~
~~Hyun G. Jang, Waltham, MA;~~
~~Shengfang Jin, Newton, MA;~~
~~Francesco G. Salituro, Marlborough, MA;~~
~~Jeffrey O. Saunders, Lincoln, MA;~~
~~Shin-San M. Su, Boston, MA;~~
~~Katharine Yen, Wellesley, MA;~~

Assignment For Published Patent Application

Agios Pharmaceuticals, Inc., Cambridge, MA

Power of Attorney: None

Domestic Priority data as claimed by applicant

This application is a CON of 13/939,519 07/11/2013

which is a CON of 13/256,396 11/29/2011 ABN  
which is a 371 of PCT/US10/27253 03/12/2010  
which claims benefit of 61/266,929 12/04/2009  
and claims benefit of 61/253,820 10/21/2009  
and claims benefit of 61/229,689 07/29/2009  
and claims benefit of 61/227,649 07/22/2009  
and claims benefit of 61/220,543 06/25/2009  
and claims benefit of 61/180,609 05/22/2009  
and claims benefit of 61/173,518 04/28/2009  
and claims benefit of 61/160,664 03/16/2009  
and claims benefit of 61/160,253 03/13/2009

**Foreign Applications** for which priority is claimed (You may be eligible to benefit from the **Patent Prosecution Highway** program at the USPTO. Please see <http://www.uspto.gov> for more information.) - None.

*Foreign application information must be provided in an Application Data Sheet in order to constitute a claim to foreign priority. See 37 CFR 1.55 and 1.76.*

**Permission to Access Application via Priority Document Exchange:** Yes

**Permission to Access Search Results:** Yes

Applicant may provide or rescind an authorization for access using Form PTO/SB/39 or Form PTO/SB/69 as appropriate.

**If Required, Foreign Filing License Granted:** 05/23/2017

The country code and number of your priority application, to be used for filing abroad under the Paris Convention, is **US 15/589,615**

**Projected Publication Date:** To Be Determined - pending completion of Missing Parts

**Non-Publication Request:** No

**Early Publication Request:** No

**Title**

METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS

**Preliminary Class**

**Statement under 37 CFR 1.55 or 1.78 for AIA (First Inventor to File) Transition Applications:** No

## **PROTECTING YOUR INVENTION OUTSIDE THE UNITED STATES**

Since the rights granted by a U.S. patent extend only throughout the territory of the United States and have no effect in a foreign country, an inventor who wishes patent protection in another country must apply for a patent in a specific country or in regional patent offices. Applicants may wish to consider the filing of an international application under the Patent Cooperation Treaty (PCT). An international (PCT) application generally has the same effect as a regular national patent application in each PCT-member country. The PCT process **simplifies** the filing of patent applications on the same invention in member countries, but **does not result** in a grant of "an international patent" and does not eliminate the need of applicants to file additional documents and fees in countries where patent protection is desired.

Almost every country has its own patent law, and a person desiring a patent in a particular country must make an application for patent in that country in accordance with its particular laws. Since the laws of many countries differ in various respects from the patent law of the United States, applicants are advised to seek guidance from specific foreign countries to ensure that patent rights are not lost prematurely.

Applicants also are advised that in the case of inventions made in the United States, the Director of the USPTO must issue a license before applicants can apply for a patent in a foreign country. The filing of a U.S. patent application serves as a request for a foreign filing license. The application's filing receipt contains further information and guidance as to the status of applicant's license for foreign filing.

Applicants may wish to consult the USPTO booklet, "General Information Concerning Patents" (specifically, the section entitled "Treaties and Foreign Patents") for more information on timeframes and deadlines for filing foreign patent applications. The guide is available either by contacting the USPTO Contact Center at 800-786-9199, or it can be viewed on the USPTO website at <http://www.uspto.gov/web/offices/pac/doc/general/index.html>.

For information on preventing theft of your intellectual property (patents, trademarks and copyrights), you may wish to consult the U.S. Government website, <http://www.stopfakes.gov>. Part of a Department of Commerce initiative, this website includes self-help "toolkits" giving innovators guidance on how to protect intellectual property in specific countries such as China, Korea and Mexico. For questions regarding patent enforcement issues, applicants may call the U.S. Government hotline at 1-866-999-HALT (1-866-999-4258).

## **LICENSE FOR FOREIGN FILING UNDER**

### **Title 35, United States Code, Section 184**

### **Title 37, Code of Federal Regulations, 5.11 & 5.15**

#### **GRANTED**

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 37 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Bureau of Industry and Security, Department of Commerce (15 CFR parts 730-774); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

**NOT GRANTED**

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15(b).

---

***SelectUSA***

The United States represents the largest, most dynamic marketplace in the world and is an unparalleled location for business investment, innovation, and commercialization of new technologies. The U.S. offers tremendous resources and advantages for those who invest and manufacture goods here. Through SelectUSA, our nation works to promote and facilitate business investment. SelectUSA provides information assistance to the international investor community; serves as an ombudsman for existing and potential investors; advocates on behalf of U.S. cities, states, and regions competing for global investment; and counsels U.S. economic development organizations on investment attraction best practices. To learn more about why the United States is the best country in the world to develop technology, manufacture products, deliver services, and grow your business, visit <http://www.SelectUSA.gov> or call +1-202-482-6800.

## Electronic Patent Application Fee Transmittal

<b>Application Number:</b>	15589615			
<b>Filing Date:</b>	08-May-2017			
<b>Title of Invention:</b>	METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS			
<b>First Named Inventor/Applicant Name:</b>	Leonard Luan C			
<b>Filer:</b>	Ioana Davies			
<b>Attorney Docket Number:</b>	AGS-013C2			
Filed as Large Entity				
<b>Filing Fees for Utility under 35 USC 111(a)</b>				
<b>Description</b>	<b>Fee Code</b>	<b>Quantity</b>	<b>Amount</b>	<b>Sub-Total in USD(\$)</b>
<b>Basic Filing:</b>				
<b>Pages:</b>				
UTILITY APPL SIZE FEE PER 50 SHEETS >100	1081	2	400	800
<b>Claims:</b>				
<b>Miscellaneous-Filing:</b>				
PROCESSING FEE, EXCEPT PROV. APPLS.	1830	1	140	140
<b>Petition:</b>				
<b>Patent-Appeals-and-Interference:</b>				

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
<b>Post-Allowance-and-Post-Issuance:</b>				
<b>Extension-of-Time:</b>				
<b>Miscellaneous:</b>				
<b>Total in USD (\$)</b>				<b>940</b>



## Electronic Acknowledgement Receipt

<b>EFS ID:</b>	29832415
<b>Application Number:</b>	15589615
<b>International Application Number:</b>	
<b>Confirmation Number:</b>	7017
<b>Title of Invention:</b>	METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED DISORDERS
<b>First Named Inventor/Applicant Name:</b>	Leonard Luan C
<b>Customer Number:</b>	148106
<b>Filer:</b>	Ioana Davies
<b>Filer Authorized By:</b>	
<b>Attorney Docket Number:</b>	AGS-013C2
<b>Receipt Date:</b>	20-JUL-2017
<b>Filing Date:</b>	08-MAY-2017
<b>Time Stamp:</b>	10:40:33
<b>Application Type:</b>	Utility under 35 USC 111(a)

### Payment information:

Submitted with Payment	yes
Payment Type	CARD
Payment was successfully received in RAM	\$940
RAM confirmation Number	072017INTEFSW10412300
Deposit Account	
Authorized User	

The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows:

<b>File Listing:</b>					
<b>Document Number</b>	<b>Document Description</b>	<b>File Name</b>	<b>File Size(Bytes)/ Message Digest</b>	<b>Multi Part /.zip</b>	<b>Pages (if appl.)</b>
1		Response_to_Notice_to_File_Missing_Parts_of_Application.pdf	22686 b36e8823991e1c35a3f4fd187d347fa76f9818f	yes	3
<b>Multipart Description/PDF files in .zip description</b>					
	<b>Document Description</b>		<b>Start</b>	<b>End</b>	
	Applicant Response to Pre-Exam Formalities Notice		1	1	
	Specification		2	2	
	Applicant Response to Pre-Exam Formalities Notice		3	3	
<b>Warnings:</b>					
<b>Information:</b>					
2	Specification	Continuation_Application_with_updated_pages_179-183_clean.pdf	1315437 3c7058509e6763b902d569129ab18c75ebcb987a	no	186
<b>Warnings:</b>					
<b>Information:</b>					
3	Specification	Continuation_Application_with_updated_pages_179-183_marked.pdf	1338561 67f9f635e5b9bb9c9fb784602308765671808/a8	no	188
<b>Warnings:</b>					
<b>Information:</b>					
4	Request for Transfer of a Computer Readable Form	Request_For_Transfer_of_Computer_Readable_Form.pdf	25239 7c5d415b541d4daa40d7eb1827a47f92eb12a458	no	1
<b>Warnings:</b>					
<b>Information:</b>					
5	Sequence Listing	C2081-701320_Sequence_Listing_v.PDF	3202208 ea3bf368e1f13ed1c89a98b25b7b991130be2d10	no	176

<b>Warnings:</b>					
<b>Information:</b>					
6	Applicant Response to Pre-Exam Formalities Notice	NTFMP.PDF	144690 3133ba09d1e43e912215ba0832db13ebe08ddb68	no	3
<b>Warnings:</b>					
<b>Information:</b>					
7	Request for Corrected Filing Receipt	Request_for_Corrected_Filing_Receipt.pdf	24729 9b0b9252a6c5290789e3d54be7663fe7970d8f60	no	3
<b>Warnings:</b>					
<b>Information:</b>					
8	Application Data Sheet	Application_Data_Sheet.pdf	31750 5a0976f94e9837bd759ec145b30ba72f37e64828	no	11
<b>Warnings:</b>					
<b>Information:</b>					
This is not an USPTO supplied ADS fillable form					
9	Request for Corrected Filing Receipt	Marked_up_Filing_Receipt.pdf	522263 ca80d721e088132b80e196e8e7024a3869f1c6ee	no	4
<b>Warnings:</b>					
<b>Information:</b>					
10	Fee Worksheet (SB06)	fee-info.pdf	32120 e23ff4561d30ea605be13dd37f3006405696aea2	no	2
<b>Warnings:</b>					
<b>Information:</b>					
<b>Total Files Size (in bytes):</b>				6659683	

**This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.**

**New Applications Under 35 U.S.C. 111**

**If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.**

**National Stage of an International Application under 35 U.S.C. 371**

**If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.**

**New International Application Filed with the USPTO as a Receiving Office**

**If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.**

Docket No.: AGS-013C2  
(PATENT)

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

---

In re Patent Application of:  
Lenny Dang et al.

Application No.: 15/589,615

Confirmation No.: 7017

Filed: May 8, 2017

Art Unit: N/A

For: METHODS AND COMPOSITIONS FOR  
CELL-PROLIFERATION-RELATED  
DISORDERS

---

Examiner: Not Yet Assigned

**AMENDMENT AND RESPONSE TO NOTICE TO FILE MISSING PARTS OF  
APPLICATION**

MS Missing Parts  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Dear Sir:

In response to the Notice to File Missing Parts of Application mailed May 24, 2017,  
Applicant respectfully submits:

An **Amendment to the Specification**, which begins on page 2 of this paper;

\$800.00 for pages in excess of 100 application size fee; and

Part 2 Copy of Notice.

**AMENDMENTS TO THE SPECIFICATION**

Please amend the above-identified U.S. patent application specification with the following:

A substitute specification in compliance with 37 CFR 1.52, 1.121 (b)(3), and 1.125, submitted with markings and accompanied by a clean version without markings;

A paper copy of a "Sequence Listing" as required by 37 CFR 1.821 (c);

A request for transfer of a "Sequence Listing" in computer readable form as required by 37 CFR 1.821 (e);

The substitute specification and the Sequence Listing contain no new matter.

**REMARKS**

Applicant believes no further fee is due with this response. However, if such a fee is due, please charge our Deposit Account No. 07-1700, under Order No. AGS-013C2 from which the undersigned is authorized to draw.

Dated: July 20, 2017

Respectfully submitted,

Electronic signature: /Ioana Davies/

Ioana Davies

Registration No.: 75,817

GOODWIN PROCTER LLP

100 Northern Avenue

Boston, Massachusetts 02210

(617) 570-3920

**METHODS AND COMPOSITIONS FOR CELL-PROLIFERATION-RELATED  
DISORDERS**

**CLAIM OF PRIORITY**

This application is a continuation of U.S.S.N. 13/939,519, filed July 11, 2013, which is a continuation of U.S.S.N. 13/256,396, filed November 29, 2011, which is a national stage application under 35 U.S.C. §371 of International Application No. PCT/US2010/027253, filed March 12, 2010, published as International Publication No. WO 2010/105243 on September 16, 2010, which claims priority to U.S.S.N. 61/160,253, filed March 13, 2009; U.S.S.N. 61/160,664, filed March 16, 2009; U.S.S.N. 61/173,518, filed April 28, 2009; U.S.S.N. 61/180,609, filed May 22, 2009; U.S.S.N. 61/220,543, filed June 25, 2009; U.S.S.N. 61/227,649, filed July 22, 2009; U.S.S.N. 61/229,689, filed July 29, 2009; U.S.S.N. 61/253,820, filed October 21, 2009; and U.S.S.N. 61/266,929, filed December 4, 2009, the contents of each of which are incorporated herein by reference.

**FIELD OF THE INVENTION**

The invention relates to methods and compositions for evaluating and treating cell proliferation-related disorders, *e.g.*, proliferative disorders such as cancer.

**BACKGROUND**

Isocitrate dehydrogenase, also known as IDH, is an enzyme which participates in the citric acid cycle. It catalyzes the third step of the cycle: the oxidative decarboxylation of isocitrate, producing alpha-ketoglutarate ( $\alpha$ -ketoglutarate or  $\alpha$ -KG) and CO<sub>2</sub> while converting NAD<sup>+</sup> to NADH. This is a two-step process, which involves oxidation of isocitrate (a secondary alcohol) to oxalosuccinate (a ketone), followed by the decarboxylation of the carboxyl group beta to the ketone, forming alpha-ketoglutarate. Another isoform of the enzyme catalyzes the same reaction; however this reaction is unrelated to the citric acid cycle, is carried out in the cytosol as well as the mitochondrion and peroxisome, and uses NADP<sup>+</sup> as a cofactor instead of NAD<sup>+</sup>.



### SUMMARY OF THE INVENTION

Methods and compositions disclosed herein relate to the role played in disease by neoactive products produced by neoactive mutant enzymes, e.g., mutant metabolic pathway enzymes. The inventors have discovered, *inter alia*, a neoactivity associated with IDH mutants and that the product of the neoactivity can be significantly elevated in cancer cells. Disclosed herein are methods and compositions for treating, and methods of evaluating, subjects having or at risk for a disorder, e.g., a cell proliferation-related disorder characterized by a neoactivity in a metabolic pathway enzyme, e.g., IDH neoactivity. Such disorders include e.g., proliferative disorders such as cancer. The inventors have discovered and disclosed herein novel therapeutic agents for the treatment of disorders, e.g., cancers, characterized by, e.g., by a neoactivity, neoactive protein, neoactive mRNA, or neoactive mutations. In embodiments a therapeutic agent reduces levels of neoactivity or neoactive product or ameliorates an effect of a neoactive product. Methods described herein also allow the identification of a subject, or identification of a treatment for the subject, on the basis of neoactivity genotype or phenotype. This evaluation can allow for optimal matching of subject with treatment, e.g., where the selection of subject, treatment, or both, is based on an analysis of neoactivity genotype or phenotype. E.g., methods describe herein can allow selection of a treatment regimen comprising administration of a novel compound, e.g., a novel compound disclosed herein, or a known compound, e.g., a known compound not previously recommended for a selected disorder. In embodiments the known compound reduces levels of neoactivity or neoactive product or ameliorates an effect of a neoactive product. Methods described herein can guide and provide a basis for selection and administration of a novel compound or a known compound, or combination of compounds, not previously recommended for subjects having a disorder characterized by a somatic neoactive mutation in a metabolic pathway enzyme. In embodiments the neoactive genotype or phenotype can act as a biomarker the presence of which indicates that a compound, either novel, or previously known, should be administered, to treat a disorder characterized by a somatic neoactive mutation in a metabolic pathway enzyme. Neoactive mutants of IDH1 having a neoactivity that results in the production of 2-hydroxyglutarate, e.g., R-2-hydroxyglutarate and associated disorders are discussed in detail herein. They are exemplary, but not limiting, examples of embodiments of the invention.

While not wishing to be bound by theory it is believed that the balance between the production and elimination of neoactive product, e.g., 2HG, e.g., R-2HG, is important in disease. Neoactive mutants, to varying degrees for varying mutations, increase the level of neoactive product, while other processes, e.g., in the case of 2HG, e.g., R-2HG, enzymatic degradation of 2HG, e.g., by 2HG dehydrogenase, reduce the level of neoactive product. An incorrect balance is associated with disease. In embodiments, the net result of a neoactive mutation at IDH1 or IDH2 result in increased levels, in affected cells, of neoactive product, 2HG, e.g., R-2HG,

Accordingly, in one aspect, the invention features, a method of treating a subject having a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder. The cell proliferation-related disorder is characterized by a somatic mutation in a metabolic pathway enzyme. The mutation is associated with a neoactivity that results in the production of a neoactivity product. The method comprises: administering to the subject a therapeutically effective amount of a therapeutic agent described herein, e.g., a therapeutic agent that decreases the level of neoactivity product encoded by a selected or mutant somatic allele, e.g., an inhibitor of a neoactivity of the metabolic pathway enzyme (the neoactive enzyme), a therapeutic agent that ameliorates an unwanted affect of the neoactivity product, or a nucleic acid based inhibitor, e.g., a dRNA which targets the neoactive enzyme mRNA, to thereby treat the subject.

In an embodiment the subject is a subject not having, or not diagnosed as having, 2-hydroxyglutaric aciduria.

In an embodiment the subject has a cell proliferation-related disorder, e.g., a cancer, characterized by the neoactivity of the metabolic pathway enzyme encoded by selected or mutant allele.

In an embodiment the subject has a cell proliferation-related disorder, e.g., a cancer, characterized by the product formed by the neoactivity of the metabolic pathway enzyme encoded by selected or mutant allele.

In one embodiment, the metabolic pathway is selected from a metabolic pathway leading to fatty acid biosynthesis, glycolysis, glutaminolysis, the pentose phosphate shunt, nucleotide biosynthetic pathways, or the fatty acid biosynthetic pathway.

In an embodiment the therapeutic agent is a therapeutic agent described herein.

In an embodiment the method comprises selecting a subject on the basis of having a cancer characterized by the selected or mutant allele, the neoactivity, or an elevated level of neoactivity product.

In an embodiment the method comprises selecting a subject on the basis of having a cancer characterized by the product formed by the neoactivity of the protein encoded by selected or mutant allele, *e.g.*, by the imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI (magnetic resonance imaging) and/or MRS (magnetic resonance spectroscopy), to determine the presence, distribution or level of the product of the neoactivity, *e.g.*, in the case of an IDH1 allele described herein, 2-hydroxyglutarate (sometimes referred to herein as 2HG), *e.g.*, R-2-hydroxyglutarate (sometimes referred to herein as R-2HG).

In an embodiment the method comprises confirming or determining, *e.g.*, by direct examination or evaluation of the subject, or sample *e.g.*, tissue, product (*e.g.*, feces, sweat, semen, exhalation, hair or nails), or bodily fluid (*e.g.*, blood (*e.g.*, blood plasma), urine, lymph, or cerebrospinal fluid or other sample sourced disclosed herein) therefrom, (*e.g.*, by DNA sequencing, immuno analysis, or assay for enzymatic activity), or receiving such information about the subject, that the cancer is characterized by the selected or mutant allele.

In an embodiment the method comprises confirming or determining, *e.g.*, by direct examination or evaluation of the subject, the level of neoactivity or the level of the product of the neoactivity, or receiving such information about the subject. In an embodiment the presence, distribution or level of the product of the neoactivity, *e.g.*, in the case of an IDH1 allele described herein, 2HG, *e.g.*, R-2HG, is determined non-invasively, *e.g.*, by imaging methods, *e.g.*, by magnetic resonance-based methods.

In an embodiment the method comprises administering a second anti-cancer agent or therapy to the subject, *e.g.*, surgical removal or administration of a chemotherapeutic.

In another aspect, the invention features, a method of treating a subject having a cell proliferation-related disorder, *e.g.*, a precancerous disorder, or cancer. In an embodiment the subject does not have, or has not been diagnosed as having, 2-hydroxyglutaric aciduria. The cell proliferation-related disorder is characterized by a somatic allele, *e.g.*, a preselected allele, or mutant allele, of an IDH, *e.g.*, IDH1 or IDH2, which encodes a mutant IDH, *e.g.*, IDH1 or IDH2, enzyme having a neoactivity.

In embodiments the neoactivity is alpha hydroxy neoactivity. As used herein, alpha hydroxy neoactivity refers to the ability to convert an alpha ketone to an alpha hydroxy. In embodiments alpha hydroxy neoactivity proceeds with a reductive cofactor, e.g., NADPH or NADH. In embodiments the alpha hydroxyl neoactivity is 2HG neoactivity. 2HG neoactivity, as used herein, refers to the ability to convert alpha ketoglutarate to 2-hydroxyglutarate (sometimes referred to herein as 2HG), e.g., R-2-hydroxyglutarate (sometimes referred to herein as R-2HG). In embodiments 2HG neoactivity proceeds with a reductive cofactor, e.g., NADPH or NADH. In an embodiment a neoactive enzyme, e.g., an alpha hydroxyl, e.g., a 2HG, neoactive enzyme, can act on more than one substrate, e.g., more than one alpha hydroxy substrate.

The method comprises administering to the subject an effective amount of a therapeutic agent of type described herein to thereby treat the subject.

In an embodiment the therapeutic agent: results in lowering the level of a neoactivity product, e.g., an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the method comprises administering a therapeutic agent that lowers neoactivity, e.g., 2HG neoactivity. In an embodiment the method comprises administering an inhibitor of a mutant IDH protein, e.g., a mutant IDH1 or mutant IDH2 protein, having a neoactivity, e.g., alpha hydroxy neoactivity, e.g., 2HG neoactivity.

In an embodiment the therapeutic agent comprises a compound from Table 24a or Table 24b or a compound having the structure of Formula (X) or (Formula (XI) described herein.

In an embodiment the therapeutic agent comprises nucleic acid-based therapeutic agent, e.g., a dsRNA, e.g., a dsRNA described herein.

In an embodiment the the therapeutic agent is an inhibitor, e.g., a polypeptide, peptide, or small molecule (e.g., a molecule of less than 1,000 daltons), or aptomer, that binds to an IDH1 mutant or wildtype subunit and inhibits neoactivity, e.g., by inhibiting formation of a dimer, e.g., a homodimer of mutant IDH1 subunits or a heterodimer of a mutant and a wildtype subunit. In an embodiment the inhibitor is a polypeptide. In an embodiment the polypeptide acts as a dominant negative with respect to the neoactivity of the mutant enzyme. The polypeptide can correspond to full length IDH1 or a fragment thereof. The polypeptide need not be identical with

the corresponding residues of wildtype IDH1, but in embodiments has at least 60, 70, 80, 90 or 95 % homology with wildtype IDH1.

In an embodiment the therapeutic agent decreases the affinity of an IDH, *e.g.*, IDH1 or IDH2 neoactive mutant protein for NADH, NADPH or a divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , or decreases the levels or availability of NADH, NADPH or divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , *e.g.*, by competing for binding to the mutant enzyme. In an embodiment the enzyme is inhibited by replacing  $Mg^{2+}$  or  $Mn^{2+}$  with  $Ca^{2+}$ .

In an embodiment the therapeutic agent is an inhibitor that reduces the level a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the level of the product of a mutant having a neoactivity of an IDH, *e.g.*, IDH1 or IDH2 mutant, *e.g.*, it reduces the level of 2HG, *e.g.*, R-2HG.

In an embodiment the therapeutic agent is an inhibitor that:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that is selected on the basis that it:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH1, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of a mutant IDH, *e.g.*, IDH1 or IDH2, protein or mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2, protein.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of neoactive enzyme activity, *e.g.*, by interacting with, *e.g.*, binding to, mutant IDH, *e.g.*, IDH1 or IDH2, protein. In an embodiment the inhibitor is other than an antibody.

In an embodiment the therapeutic agent is an inhibitor that is a small molecule

and interacts with, *e.g.*, binds, the mutant RNA, *e.g.*, mutant IDH1 or IDH2 mRNA (*e.g.*, mutant IDH1 mRNA).

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, binds, either the mutant IDH, *e.g.*, IDH1 or IDH2, protein or interacts directly with, *e.g.*, binds, the mutant IDH mRNA, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the IDH is IDH1 and the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. Mutations in IDH1 associated with 2HG neoactivity include mutations at residue 132, *e.g.*, R132H, R132C, R132S, R132G, R132L, or R132V (*e.g.*, R132H or R132C).

In an embodiment the IDH is IDH2 and the neoactivity of the IDH2 mutant is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. Mutations in IDH2 associated with 2HG neoactivity include mutations at residue 172, *e.g.*, R172K, R172M, R172S, R172G, or R172W.

Treatment methods described herein can comprise evaluating a neoactivity genotype or phenotype. Methods of obtaining and analyzing samples, and the *in vivo* analysis in subjects, described elsewhere herein, *e.g.*, in the section entitled, “Methods of evaluating samples and/or subjects,” can be combined with this method.

In an embodiment, prior to or after treatment, the method includes evaluating the growth, size, weight, invasiveness, stage or other phenotype of the cell proliferation-related disorder.

In an embodiment, prior to or after treatment, the method includes evaluating the IDH, *e.g.*, IDH1 or IDH2, alpha hydroxyl neoactivity genotype, *e.g.*, 2HG, genotype, or alpha hydroxy neoactivity phenotype, *e.g.*, 2HG, *e.g.*, R-2HG, phenotype. Evaluating the alpha hydroxyl, *e.g.*, 2HG, genotype can comprise determining if an IDH1 or IDH2 mutation having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, is present, *e.g.*, a mutation disclosed herein having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. Alpha hydroxy neoactivity phenotype, *e.g.*, 2HG, *e.g.*, R-2HG, phenotype, as used herein, refers to the level of alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, level of alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, or level of mutant enzyme having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity (or corresponding mRNA). The evaluation can be by a method described herein.

In an embodiment the subject can be evaluated, before or after treatment, to determine if the cell proliferation-related disorder is characterized by an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment a cancer, *e.g.*, a glioma or brain tumor in a subject, can be analyzed, *e.g.*, by imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS, *e.g.*, before or after treatment, to determine if it is characterized by presence of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method comprises evaluating, *e.g.*, by direct examination or evaluation of the subject, or a sample from the subject, or receiving such information about the subject, the IDH, *e.g.*, IDH1 or IDH2, genotype, or an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG phenotype of, the subject, *e.g.*, of a cell, *e.g.*, a cancer cell, characterized by the cell proliferation-related disorder. (As described in more detail elsewhere herein the evaluation can be, *e.g.*, by DNA sequencing, immuno analysis, evaluation of the presence, distribution or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, from spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS measurement, sample analysis such as serum or spinal cord fluid analysis, or by analysis of surgical material, *e.g.*, by mass-spectroscopy). In embodiments this information is used to determine or confirm that a proliferation-related disorder, *e.g.*, a cancer, is characterized by an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In embodiments this information is used to determine or confirm that a cell proliferation-related disorder, *e.g.*, a cancer, is characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having a mutation, *e.g.*, a His, Ser, Cys, Gly, Val, Pro or Leu (*e.g.*, His, Ser, Cys, Gly, Val, or Leu at residue 132, more specifically, His or Cys, or an IDH2 allele having a mutation at residue 172, *e.g.*, a K, M, S, G, or W).

In an embodiment, before and/or after treatment has begun, the subject is evaluated or monitored by a method described herein, *e.g.*, the analysis of the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, to select, diagnose or prognose the subject, to select an inhibitor, or to evaluate response to the treatment or progression of disease.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, a leukemia, *e.g.*, AML or ALL, *e.g.*, B-ALL or T-ALL, prostate cancer, fibrosarcoma, paraganglioma, or myelodysplasia or myelodysplastic syndrome (*e.g.*, B-ALL or T-ALL, prostate cancer, or myelodysplasia or myelodysplastic syndrome) and the evaluation is: evaluation of the presence, distribution, or level of an alpha

hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG; or evaluation of the presence, distribution, or level of a neoactivity, *e.g.*, an alpha hydroxy neoactivity, e.g., 2HG neoactivity, of an IDH1 or IDH2, mutant protein.

In an embodiment the disorder is other than a solid tumor. In an embodiment the disorder is a tumor that, at the time of diagnosis or treatment, does not have a necrotic portion. In an embodiment the disorder is a tumor in which at least 30, 40, 50, 60, 70, 80 or 90% of the tumor cells carry an IHD, e.g., IDH1 or IDH2, mutation having 2HG neoactivity, at the time of diagnosis or treatment.

In an embodiment the cell proliferation-related disorder is a cancer, e.g., a cancer described herein, characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by unwanted (i.e., increased) levels of an alpha hydroxy neoactivity, product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, *e.g.*, wherein the tumor is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, *e.g.*, a mutant described herein. Gliomas include astrocytic tumors, oligodendroglial tumors, oligoastrocytic tumors, anaplastic astrocytomas, and glioblastomas. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu (e.g., His, Ser, Cys, Gly, Val, or Leu), or any residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **Fig. 21**). In an embodiment the allele encodes an IDH1 having His at residue 132. In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the IDH1 allele has an A (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394A, a C394G, a C394T,



a G395C, a G395T or a G395A mutation; specifically a C394A or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8), more specifically His, Ser, Cys, Gly, Val, or Leu; or His or Cys.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8) , more specifically His, Ser, Cys, Gly, Val, or Leu; or His or Cys.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method comprises selecting a subject having a fibrosarcoma or paraganglioma wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having a fibrosarcoma or paraganglioma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having a fibrosarcoma or paraganglioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is localized or metastatic prostate cancer, *e.g.*, prostate adenocarcinoma, *e.g.*, wherein the cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

*E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al*, 2009, Int. J. Cancer, 125: 353-355 at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (*e.g.*, His, Ser, Cys,

Gly, Val, or Leu). In an embodiment the allele encodes an IDH1 having His or Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394T or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, wherein the cancer is characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a hematological cancer, *e.g.*, a leukemia, *e.g.*, AML, or ALL, wherein the hematological cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

In an embodiment the cell proliferation-related disorder is acute lymphoblastic leukemia (*e.g.*, an adult or pediatric form), *e.g.*, wherein the acute lymphoblastic leukemia (sometimes referred to herein as ALL) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. The ALL can be, *e.g.*, B-ALL or T-ALL. In an embodiment the cancer is characterized by increased levels of 2 an alpha hydroxy neoactivity product, *e.g.*, HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID

NO:8 (see also **FIG. 21**), more specifically His, Ser, Cys, Gly, Val, or Leu. In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, characterized by an IDH1 allele described herein, e.g., an IDH1 allele having Cys at residue 132 according to the sequence of SEQ ID NO:8.

In an embodiment the method comprises selecting a subject ALL, e.g., B-ALL or T-ALL, on the basis of cancer being characterized by having an IDH1 allele described herein, e.g., an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the cell proliferation-related disorder is acute myelogenous leukemia (e.g., an adult or pediatric form), e.g., wherein the acute myelogenous leukemia (sometimes referred to herein as AML) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, e.g., a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type. E.g., in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). E.g., the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**). In an embodiment the allele encodes an IDH1 having Cys, His or Gly at residue 132, more specifically, Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplasmic leukemia (AML) characterized by an IDH1 allele described herein, e.g., an IDH1 allele having Cys, His, or Gly at residue 132 according to the sequence of SEQ ID NO:8, more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML) on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His, or Gly at residue 132 (SEQ ID NO:8), more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML), on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method further comprises evaluating the subject for the presence of a mutation in the NRAS or NPMc gene.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, *e.g.*, wherein the myelodysplasia or myelodysplastic syndrome is characterized by having an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the disorder is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), more specifically His, Ser, Cys, Gly, Val, or Leu. In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His, or Gly at residue 132 according to the sequence of SEQ ID NO:8, more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His, or Gly at residue 132 (SEQ ID NO:8), more specifically, Cys at residue 132.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being

characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the cell proliferation-related disorder is a glioma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, e.g., 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH2 allele described herein, e.g., an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), more specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH2 allele described herein, e.g., an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), more specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment the cell proliferation-related disorder is a prostate cancer, e.g., prostate adenocarcinoma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, e.g., 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a prostate cancer, e.g., prostate adenocarcinoma, wherein the cancer is characterized by having

an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is ALL, *e.g.*, B-ALL or T-ALL, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**). In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is AML, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys,

Gly, Met, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172. In an embodiment the allele encodes an IDH2 having Gly at residue 172.

In an embodiment the method comprises selecting a subject having AML, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly or Met at residue 172 (SEQ ID NO:10), more specifically Lys or Met.

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, or Met at residue 172 (SEQ ID NO:10), more specifically Lys or Met.

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, characterized by a mutation, or preselected allele, of IDH2. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), more specifically Lys, Gly, Met, Trp or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172. In an embodiment the allele encodes an IDH2 having Gly at residue 172.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, or Met at residue 172 (SEQ ID NO:10), in specific embodiments, Lys or Met.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, or Met at residue 172 (SEQ ID NO:10), in specific embodiments, Lys or Met.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being

characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment a product of the neoactivity is 2HG (e.g., R-2HG) which acts as a metabolite. In another embodiment a product of the neoactivity is 2HG (e.g., R-2HG) which acts as a toxin, e.g., a carcinogen.

In some embodiments, the methods described herein can result in reduced side effects relative to other known methods of treating cancer.

Therapeutic agents and methods of subject evaluation described herein can be combined with other therapeutic modalities, e.g., with art-known treatments.

In an embodiment the method comprises providing a second treatment, to the subject, e.g., surgical removal, irradiation or administration of a chemotherapeutic agent, e.g., an administration of an alkylating agent. Administration (or the establishment of therapeutic levels) of the second treatment can: begin prior to the beginning or treatment with (or prior to the establishment of therapeutic levels of) the inhibitor; begin after the beginning or treatment with (or after the establishment of therapeutic levels of) the inhibitor, or can be administered concurrently with the inhibitor, e.g., to achieve therapeutic levels of both concurrently.

In an embodiment the cell proliferation-related disorder is a CNS tumor, e.g., a glioma, and the second therapy comprises administration of one or more of: radiation; an alkylating agent, e.g., temozolomide, e.g., Temoader®, or BCNU; or an inhibitor of HER1/EGFR tyrosine kinase, e.g., erlotinib, e.g., Tarceva®.

The second therapy, e.g., in the case of glioma, can comprise implantation of BCNU or carmustine in the brain, e.g., implantation of a Gliadel® wafer.

The second therapy, e.g., in the case of glioma, can comprise administration of imatinib, e.g., Gleevec®.

In an embodiment the cell proliferation-related disorder is prostate cancer and the second therapy comprises one or more of: androgen ablation; administration of a microtubule stabilizer, e.g., docetaxol, e.g., Taxotere®; or administration of a topoisomerase II inhibitor, e.g., mitoxantrone.

In an embodiment the cell proliferation-related disorder is ALL, e.g., B-ALL or T-ALL, and the second therapy comprises one or more of:

induction phase treatment comprising the administration of one or more of: a steroid; an inhibitor of microtubule assembly, e.g., vincristine; an agent that reduces



the availability of asparagine, e.g., asparaginase; an anthracycline; or an antimetabolite, e.g., methotrexate, e.g., intrathecal methotrexate, or 6-mercaptopurine;

consolidation phase treatment comprising the administration of one or more of: a drug listed above for the induction phase; an antimetabolite, e.g., a guanine analog, e.g., 6-thioguanine; an alkylating agent, e.g., cyclophosphamide; an anti-metabolite, e.g., AraC or cytarabine; or an inhibitor of topoisomerase I, e.g., etoposide; or

maintenance phase treatment comprising the administration of one or more of the drugs listed above for induction or consolidation phase treatment.

In an embodiment the cell proliferation-related disorder is AML and the second therapy comprises administration of one or more of: an inhibitor of topoisomerase II, e.g., daunorubicin, idarubicin, topotecan or mitoxantrone; an inhibitor of topoisomerase I, e.g., etoposide; or an anti-metabolite, e.g., AraC or cytarabine.

In another aspect, the invention features, a method of evaluating, e.g. diagnosing, a subject, e.g., a subject not having, or not diagnosed as having, 2-hydroxyglutaric aciduria. The method comprises analyzing a parameter related to the neoactivity genotype or phenotype of the subject, e.g., analyzing one or more of:

a) the presence, distribution, or level of a neoactive product, e.g., the product of an alpha hydroxy neoactivity, e.g., 2HG, e.g., R-2HG, e.g., an increased level of product, 2HG, e.g., R-2HG (as used herein, an increased level of a product of an alpha hydroxy neoactivity, e.g., 2HG, e.g., R-2HG, or similar term, e.g., an increased level of neoactive product or neoactivity product, means increased as compared with a reference, e.g., the level seen in an otherwise similar cell lacking the IDH mutation, e.g., IDH1 or IDH2 mutation, or in a tissue or product from a subject not having the mutation (the terms increased and elevated as referred to the level of a product of alpha hydroxyl neoactivity as used herein, are used interchangeably);

b) the presence, distribution, or level of a neoactivity, e.g., alpha hydroxy neoactivity, e.g., 2HG neoactivity, of an IDH1 or IDH2, mutant protein;

c) the presence, distribution, or level of a neoactive mutant protein, e.g., an IDH, e.g., an IDH1 or IDH2, mutant protein which has a neoactivity, e.g., alpha hydroxy neoactivity, e.g., 2HG neoactivity, or a corresponding RNA; or

d) the presence of a selected somatic allele or mutation conferring neoactivity, e.g., an IDH, e.g., IDH1 or IDH2, which encodes a protein with a neoactivity, e.g.,

alpha hydroxy neoactivity, e.g., 2HG neoactivity, e.g., an allele disclosed herein, in cells characterized by a cell proliferation-related disorder from the subject, thereby evaluating the subject.

In an embodiment analyzing comprises performing a procedure, e.g., a test, to provide data or information on one or more of a-d, e.g., performing a method which results in a physical change in a sample, in the subject, or in a device or reagent used in the analysis, or which results in the formation of an image representative of the data. Methods of obtaining and analyzing samples, and the in vivo analysis in subjects, described elsewhere herein, e.g., in the section entitled, "Methods of evaluating samples and/or subjects," can be combined with this method. In another embodiment analyzing comprises receiving data or information from such test from another party. In an embodiment the analyzing comprises receiving data or information from such test from another party and, the method comprises, responsive to that data or information, administering a treatment to the subject.

As described herein, the evaluation can be used in a number of applications, e.g., for diagnosis, prognosis, staging, determination of treatment efficacy, patent selection, or drug selection.

Thus, in an embodiment method further comprises, e.g., responsive to the analysis of one or more of a-d:

diagnosing the subject, e.g., diagnosing the subject as having a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder;

staging the subject, e.g., determining the stage of a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder;

providing a prognosis for the subject, e.g., providing a prognosis for a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a precancerous disorder;

determining the efficacy of a treatment, e.g., the efficacy of a chemotherapeutic agent, irradiation or surgery;

determining the efficacy of a treatment with a therapeutic agent, e.g., an inhibitor, described herein;

selecting the subject for a treatment for a cell proliferation-related disorder, e.g., a disorder characterized by unwanted cell proliferation, e.g., cancer, or a

precancerous disorder. The selection can be based on the need for a reduction in neoactivity or on the need for amelioration of a condition associated with or resulting from neoactivity. For example, if it is determined that the subject has a cell proliferation-related disorder, *e.g.*, *e.g.*, cancer, or a precancerous disorder characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, or by a mutant IDH1 or IDH2, having alpha hydroxyl neoactivity, *e.g.*, 2HG, neoactivity, selecting the subject for treatment with a therapeutic agent described herein, *e.g.*, an inhibitor (*e.g.*, a small molecule or a nucleic acid-based inhibitor) of the neoactivity of that mutant (*e.g.*, conversion of alpha-ketoglutarate to 2HG, *e.g.*, R-2HG);

correlating the analysis with an outcome or a prognosis;

providing a value for an analysis on which the evaluation is based, *e.g.*, the value for a parameter correlated to the presence, distribution, or level of an alpha hydroxyl neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG;

providing a recommendation for treatment of the subject; or

memorializing a result of, or output from, the method, *e.g.*, a measurement made in the course of performing the method, and optionally transmitting the memorialization to a party, *e.g.*, the subject, a healthcare provider, or an entity that pays for the subject's treatment, *e.g.*, a government, insurance company, or other third party payer.

As described herein, the evaluation can provide information on which a number of decisions or treatments can be based.

Thus, in an embodiment the result of the evaluation, *e.g.*, an increased level of an alpha hydroxyl neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, the presence of an IDH, *e.g.*, IDH1 or IDH2, neoactivity, *e.g.*, alpha hydroxyl neoactivity, *e.g.*, 2HG neoactivity, the presence of an IDH, *e.g.*, IDH1 or IDH2, mutant protein (or corresponding RNA) which has alpha hydroxyl neoactivity, *e.g.*, 2HG neoactivity, the presence of a mutant allele of IDH, *e.g.*, IDH1 or IDH2, having alpha hydroxyl neoactivity, 2HG neoactivity, *e.g.*, an allele disclosed herein, is indicative of:

a cell proliferation-related disorder, *e.g.*, cancer, *e.g.*, it is indicative of a primary or metastatic lesion;

the stage of a cell proliferation-related disorder;

a prognosis or outcome for a cell proliferation-related disorder, *e.g.*, it is indicative of a less aggressive form of the disorder, *e.g.*, cancer. *E.g.*, in the case of

glioma, presence of an alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG, can indicate a less aggressive form of the cancer;

the efficacy of a treatment, e.g., the efficacy of a chemotherapeutic agent, irradiation or surgery;

the need of of a therapy disclosed herein, e.g., inhibition a neoactivity of an IDH, e.g., IDH1 or IDH2, neoactive mutant described herein. In an embodiment relatively higher levels (or the presence of the mutant) is correlated with need of inhibition a neoactivity of an IDH, e.g., IDH1 or IDH2, mutant described herein; or

responsiveness to a treatment. The result can be used as a noninvasive biomarker for clinical response. E.g., elevated levels can be predictive on better outcome in glioma patients (e.g., longer life expectancy).

As described herein, the evaluation can provide for the selection of a subject.

Thus, in an embodiment the method comprises, e.g., responsive to the analysis of one or more of a-d, selecting a subject, e.g., for a treatment. The subject can be selected on a basis described herein, e.g., on the basis of:

said subject being at risk for, or having, higher than normal levels of an alpha hydroxy neoactivity product, e.g., 2-hydroxyglutarate (e.g., R-2HG) in cell having a cell proliferation-related disorder, e.g., a leukemia such as AML or ALL, e.g., B-ALL or T-ALL, or a tumor lesion, e.g., a glioma or a prostate tumor;

said subject having a proliferation-related disorder characterized by a selected IDH, e.g., IDH1 or IDH2 allele, e.g., an IDH1 or IDH2 mutation, having alpha hydroxyl neoactivity, e.g., 2HG neoactivity;

said subject having a selected IDH allele, e.g., a selected IDH1 or IDH2 allele; having alpha hydroxyl neoactivity, e.g., 2HG neoactivity;

said subject having a proliferation-related disorder;

said subject being in need of, or being able to benefit from, a therapeutic agent of a type described herein;

said subject being in need of, or being able to benefit from, a compound that inhibits alpha hydroxyl neoactivity, e.g., 2HG neoactivity;

said subject being in need of, or being able to benefit from, a compound that lowers the level of an alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG.

In an embodiment evaluation comprises selecting the subject, e.g., for treatment with an anti-neoplastic agent, on the establishment of, or determination that, the subject has increased alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG,

or increased alpha hydroxyl neoactivity, e.g., 2HG neoactivity, or that the subject is in need of inhibition of a neoactivity of an IDH, e.g., IDH1 or IDH2, mutant described herein.

As described herein, the evaluations provided for by methods described herein allow the selection of optimal treatment regimens.

Thus, in an embodiment the method comprises, e.g., responsive to the analysis of one or more of a-d, selecting a treatment for the subject, e.g., selecting a treatment on a basis disclosed herein. The treatment can be the administration of a therapeutic agent disclosed herein. The treatment can be selected on the basis that:

it is useful in treating a disorder characterized by one or more of alpha hydroxyl neoactivity, e.g., 2HG neoactivity, an IDH1 or IDH2, mutant protein having alpha hydroxyl neoactivity, e.g., 2HG neoactivity (or a corresponding RNA);

it is useful in treating a disorder characterized by a selected somatic allele or mutation of an IDH, e.g., IDH1 or IDH2, which encodes a protein with alpha hydroxyl neoactivity, e.g., 2HG neoactivity, e.g., an allele disclosed herein, in cells characterized by a cell proliferation-related disorder from the subject;

it reduces the level of an alpha hydroxyl neoactivity product, e.g., 2HG, e.g., R-2HG;

it reduces the level of alpha hydroxyl neoactivity, e.g., 2HG neoactivity.

In an embodiment evaluation comprises selecting the subject, e.g., for treatment.

In embodiments the treatment is the administration of a therapeutic agent described herein.

The methods can also include treating a subject, e.g., with a treatment selected in response to, or on the basis of, an evaluation made in the method.

Thus, in an embodiment the method comprises, e.g., responsive to the analysis of one or more of a-d, administering a treatment to the subject, e.g., the administration of a therapeutic agent of a type described herein.

In an embodiment the therapeutic agent comprises a compound from Table 24a or Table 24b or a compound having the structure of Formula (X) or (XI) described below.

In an embodiment the therapeutic agent comprises nucleic acid, e.g., dsRNA, e.g., a dsRNA described herein.

In an embodiment the the therapeutic agent is an inhibitor, *e.g.*, a polypeptide, peptide, or small molecule (*e.g.*, a molecule of less than 1,000 daltons), or aptomer, that binds to an IDH1 or IDH2 mutant (*e.g.*, an aptomer that binds to an IDH1 mutant) or wildtype subunit and inhibits neoactivity, *e.g.*, by inhibiting formation of a dimer, *e.g.*, a homodimer of mutant IDH1 or IDH2 subunits (*e.g.*, a homodimer of mutant IDH1 subunits) or a heterodimer of a mutant and a wildtype subunit. In an embodiment the inhibitor is a polypeptide. In an embodiment the polypeptide acts as a dominant negative with respect to the neoactivity of the mutant enzyme. The polypeptide can correspond to full length IDH1 or IDH2 or a fragment thereof (*e.g.*, the polypeptide correspondes to full length IDH1 or a fragment thereof). The polypeptide need not be indentical with the corresponding residues of wildtype IDH1 or IDH2 (*e.g.*, wildtype IDH1), but in embodiments has at least 60, 70, 80, 90 or 95 % homology with wildtype IDH1 or IDH2 (*e.g.*, wildtype IDH1).

In an embodiment the therapeutic agent decreases the affinity of an IDH, *e.g.*, IDH1 or IDH2 neoactive mutant protein for NADH, NADPH or a divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , or decreases the levels or availability of NADH, NADPH or divalent metal ion, *e.g.*,  $Mg^{2+}$  or  $Mn^{2+}$ , *e.g.*, by competing for binding to the mutant enzyme. In an embodiment the enzyme is inhibited by replacing  $Mg^{2+}$  or  $Mn^{2+}$  with  $Ca^{2+}$ .

In an embodiment the therapeutic agent is an inhibitor that reduces the level a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the level of the product of a mutant having a neoactivity of an IDH, *e.g.*, IDH1 or IDH2 mutant, *e.g.*, it reduces the level of 2HG, *e.g.*, R-2HG.

In an embodiment the therapeutic agent is an inhibitor that:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that is selected on the basis that it:

inhibits, *e.g.*, specifically, a neoactivity of an IDH, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein *e.g.*, 2HG neoactivity; or

inhibits both the wildtype activity and a neoactivity of an IDH1, *e.g.*, IDH1 or IDH2, *e.g.*, a neoactivity described herein, *e.g.*, 2HG neoactivity.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of a mutant IDH, *e.g.*, IDH1 or IDH2, protein or mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, it binds to, the mutant IDH, *e.g.*, IDH1 or IDH2, protein.

In an embodiment the therapeutic agent is an inhibitor that reduces the amount of neoactive enzyme activity, *e.g.*, by interacting with, *e.g.*, binding to, mutant IDH, *e.g.*, IDH1 or IDH2, protein. In an embodiment the inhibitor is other than an antibody.

In an embodiment the therapeutic agent is an inhibitor that is a small molecule and interacts with, *e.g.*, binds, the mutant RNA, *e.g.*, mutant IDH1 mRNA.

In an embodiment the therapeutic agent is an inhibitor that interacts directly with, *e.g.*, binds, either the mutant IDH, *e.g.*, IDH1 or IDH2, protein or interacts directly with, *e.g.*, binds, the mutant IDH mRNA, *e.g.*, IDH1 or IDH2 mRNA.

In an embodiment the therapeutic agent is administered.

In an embodiment the treatment: inhibits, *e.g.*, specifically, a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, a neoactivity described herein; or inhibits both the wildtype and activity and a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, a neoactivity described herein. In an embodiment, the subject is subsequently evaluated or monitored by a method described herein, *e.g.*, the analysis of the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, to evaluate response to the treatment or progression of disease.

In an embodiment the treatment is selected on the basis that it: inhibits, *e.g.*, specifically, a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity; or inhibits both the wildtype and activity and a neoactivity of IDH1 or IDH2 (*e.g.*, a neoactivity of IDH1), *e.g.*, a neoactivity described herein.

In an embodiment, the method comprises determining the possibility of a mutation other than a mutation in IDH1 or in IDH2. In embodiments a relatively high level of 2HG, *e.g.*, R-2HG is indicative of another mutation.

In an embodiment, which embodiment includes selecting or administering a treatment for the subject, the subject:

has not yet been treated for the subject the cell proliferation-related disorder and the selected or administered treatment is the initial or first line treatment;

has already been treated for the the cell proliferation-related and the selected or administered treatment results in an alteration of the existing treatment;

has already been treated for the the cell proliferation-related, and the selected treatment results in continuation of the existing treatment; or

has already been treated for the the cell proliferation-related disorder and the selected or administered treatment is different, *e.g.*, as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment, which embodiment includes selecting or administering a treatment for the subject, the selected or administered treatment can comprise:

a treatment which includes administration of a therapeutic agent at different, *e.g.*, a greater (or lesser) dosage (*e.g.*, different as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG);

a treatment which includes administration of a therapeutic agent at a different frequency, *e.g.*, more or less frequently, or not at all (*e.g.*, different as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG); or

a treatment which includes administration of a therapeutic agent in a different therapeutic setting (*e.g.*, adding or deleting a second treatment from the treatment regimen) (*e.g.*, different as compared to what was administered prior to the evaluation or to what would be administered in the absence of elevated levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG).

Methods of evaluating a subject described herein can comprise evaluating a neoactivity genotype or phenotype. Methods of obtaining and analyzing samples, and the in vivo analysis in subjects, described elsewhere herein, *e.g.*, in the section entitled, "Methods of evaluating samples and/or subjects," can be combined with this method.

In an embodiment the method comprises:



subjecting the subject (*e.g.*, a subject not having 2-hydroxyglutaric aciduria) to imaging and/or spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI and/or MRS *e.g.*, imaging analysis, to provide a determination of the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, *e.g.*, as associated with a tumor, *e.g.*, a glioma, in the subject;

optionally storing a parameter related to the determination, *e.g.*, the image or a value related to the image from the imaging analysis, in a tangible medium; and

responsive to the determination, performing one or more of: correlating the determination with outcome or with a prognosis; providing an indication of outcome or prognosis; providing a value for an analysis on which the evaluation is based, *e.g.*, the presence, distribution, or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG; providing a recommendation for treatment of the subject; selecting a course of treatment for the subject, *e.g.*, a course of treatment described herein, *e.g.*, selecting a course of treatment that includes inhibiting a neoactivity of a mutant IDH, *e.g.*, IDH1 or IDH2, allele, *e.g.*, a neoactivity described herein; administering a course of treatment to the subject, *e.g.*, a course of treatment described herein, *e.g.*, a course of treatment that includes inhibiting a neoactivity of a mutant IDH, *e.g.*, IDH1 or IDH2, allele, *e.g.*, a neoactivity described herein; and memorializing memorializing a result of the method or a measurement made in the course of the method, *e.g.*, one or more of the above and/or transmitting memorialization of one or more of the above to a party, *e.g.*, the subject, a healthcare provider, or an entity that pays for the subject's treatment, *e.g.*, a government, insurance company, or other third party payer.

In an embodiment the method comprises confirming or determining, *e.g.*, by direct examination or evaluation of the subject, or sample *e.g.*, tissue or bodily fluid (*e.g.*, blood (*e.g.*, blood plasma), urine, lymph, or cerebrospinal fluid) therefrom, (*e.g.*, by DNA sequencing or immuno analysis or evaluation of the presence, distribution or level of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG), or receiving such information about the subject, that the subject has a cancer characterized by an IDH, *e.g.*, IDH1 or IDH2, allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8), in specific embodiments, an IDH1 allele having His, Ser, Cys, Gly, Val, or Leu at residue 132 or an IDH1 allele having His or Cys at residue 132; or an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10).

In an embodiment, prior to or after treatment, the method includes evaluating the growth, size, weight, invasiveness, stage or other phenotype of the cell proliferation-related disorder.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, a leukemia, *e.g.*, AML or ALL, *e.g.*, B-ALL or T-ALL, prostate cancer, or myelodysplasia or myelodysplastic syndrome and the evaluation is a or b. In an embodiment the method comprises evaluating a sample, *e.g.*, a sample described herein, *e.g.*, a tissue, *e.g.*, a cancer sample, or a bodily fluid, *e.g.*, serum or blood, for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment, a subject is subjected to MRS and the evaluation comprises evaluating the presence or elevated amount of a peak correlated to or corresponding to 2HG, *e.g.*, R-2HG, as determined by magnetic resonance. For example, a subject can be analyzed for the presence and/or strength of a signal at about 2.5 ppm to determine the presence and/or amount of 2HG, *e.g.*, R-2HG in the subject.

In an embodiment the method comprises obtaining a sample from the subject and analyzing the sample, or analyzing the subject, *e.g.*, by imaging the subject and optionally forming a representation of the image on a computer.

In an embodiment the results of the analysis is compared to a reference.

In an embodiment a value for a parameter correlated to the presence, distribution, or level, *e.g.*, of 2HG, *e.g.*, R-2HG, is determined. It can be compared with a reference value, *e.g.*, the value for a reference subject not having abnormal presence, level, or distribution, *e.g.*, a reference subject cell not having a mutation in IDH, *e.g.*, IDH1 or IDH2, having a neoactivity described herein.

In an embodiment the method comprises determining if an IDH, *e.g.*, IDH1 or IDH2, mutant allele that is associated with 2HG neoactivity is present. *E.g.*, in the case of IDH1, the presence of a mutation at residue 132 associated with 2HG neoactivity can be determined. In the case of IDH2, the presence of a mutation at residue 172 associated with 2HG neoactivity can be determined. The determination can comprise sequencing a nucleic acid, *e.g.*, genomic DNA or cDNA, from an affected cell, which encodes the relevant amino acid(s). The mutation can be a deletion, insertion, rearrangement, or substitution. The mutation can involve a single nucleotide, *e.g.*, a single substitution, or more than one nucleotide, *e.g.*, a deletion of more than one nucleotides.

In an embodiment the method comprises determining the sequence at position 394 or 395 of the IDH1 gene, or determining the identity of amino acid residue 132 (SEQ ID NO:8) in the IDH1 gene in a cell characterized by the cell proliferation related disorder.

In an embodiment the method comprises determining the amino acid sequence, *e.g.*, by DNA sequencing, at position 172 of the IDH2 gene in a cell characterized by the cell proliferation related disorder.

In an embodiment a product of the neoactivity is 2-HG, *e.g.*, R-2HG, which acts as a metabolite. In another embodiment a product of the neoactivity is 2HG, *e.g.*, R-2HG, which acts as a toxin, *e.g.*, a carcinogen.

In an embodiment the disorder is other than a solid tumor. In an embodiment the disorder is a tumor that, at the time of diagnosis or treatment, does not have a necrotic portion. In an embodiment the disorder is a tumor in which at least 30, 40, 50, 60, 70, 80 or 90% of the tumor cells carry an IHD, *e.g.*, IDH1 or IDH2, mutation having 2HG neoactivity, at the time of diagnosis or treatment.

In an embodiment the cell proliferation-related disorder is a cancer, *e.g.*, a cancer described herein, characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a tumor of the CNS, *e.g.*, a glioma, *e.g.*, wherein the tumor is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. Gliomas include astrocytic tumors, oligodendroglial tumors, oligoastrocytic tumors, anaplastic astrocytomas, and glioblastomas. In an embodiment the tumor is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8

(see also **Fig. 21**). In an embodiment the allele encodes an IDH1 having His at residue 132. In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the IDH1 allele has an A (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394A, a C394G, a C394T, a G395C, a G395T or a G395A mutation, specifically C394A or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8) (*e.g.*, His, Ser, Cys, Gly, Val, or Leu; or His or Cys).

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His, Ser, Cys, Gly, Val, Pro or Leu at residue 132 (SEQ ID NO:8) (*e.g.*, His, Ser, Cys, Gly, Val, or Leu; or His or Cys).

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment, the cell proliferation disorder is fibrosarcoma or paraganglioma wherein the cancer is characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment, the cell proliferation disorder is fibrosarcoma or paraganglioma wherein the cancer is characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment, the cell proliferation disorder is fibrosarcoma or paraganglioma wherein the cancer is characterized by increased levels of an alpha hydroxy neoactivity, product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is localized or metastatic prostate cancer, *e.g.*, prostate adenocarcinoma, *e.g.*, wherein the cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type.

*E.g.*, in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al*, 2009, Int. J. Cancer, 125: 353-355 at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (*e.g.*, His, Ser, Cys, Gly, Val, or Leu). In an embodiment the allele encodes an IDH1 having His or Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394, or an A (or any other nucleotide other than G) at nucleotide position 395. In an embodiment the allele is a C394T or a G395A mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, wherein the cancer is characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having His or Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having prostate cancer, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a hematological cancer, *e.g.*, a leukemia, *e.g.*, AML, or ALL, wherein the hematological cancer is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is acute lymphoblastic leukemia (*e.g.*, an adult or pediatric form), *e.g.*, wherein the acute lymphoblastic leukemia (sometimes referred to herein as ALL) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. The ALL can be, *e.g.*, B-ALL or T-ALL. In an embodiment the

cancer is characterized by increased levels of 2 an alpha hydroxy neoactivity product, e.g., HG, e.g., R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (e.g., His, Ser, Cys, Gly, Val, or Leu). In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 according to the sequence of SEQ ID NO:8.

In an embodiment the method comprises selecting a subject ALL, e.g., B-ALL or T-ALL, on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having ALL, e.g., B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is acute myelogenous leukemia (*e.g.*, an adult or pediatric form), *e.g.*, wherein the acute myelogenous leukemia (sometimes referred to herein as AML) is characterized by an IDH1 somatic mutant having alpha hydroxy neoactivity, e.g., 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the cancer is characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, e.g., R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**) (e.g., His, Ser, Cys, Gly, Val or Leu). In an embodiment the allele encodes an IDH1 having Cys, His or Gly at residue 132, specifically, Cys.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML) characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His or Gly at residue 132 according to the sequence of SEQ ID NO:8, specifically, Cys.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML) on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys, His or Gly at residue 132 (SEQ ID NO:8), specifically, Cys.

In an embodiment the method comprises selecting a subject having acute myelogenous lymphoplastic leukemia (AML), on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the method further comprises evaluating the subject for the presence of a mutation in the NRAS or NPMc gene.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, *e.g.*, wherein the myelodysplasia or myelodysplastic syndrome is characterized by having an IDH1 somatic mutant having alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity, *e.g.*, a mutant described herein. In an embodiment the disorder is characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG, as compared to non-diseased cells of the same type. *E.g.*, in an embodiment, the IDH1 allele is an IDH1 having other than an Arg at residue 132 (SEQ ID NO:8). *E.g.*, the allele encodes His, Ser, Cys, Gly, Val, Pro or Leu, or any residue described in Kang *et al.*, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), specifically, His, Ser, Cys, Gly, Val, or Leu. In an embodiment the allele encodes an IDH1 having Cys at residue 132.

In an embodiment the IDH1 allele has a T (or any other nucleotide other than C) at nucleotide position 394. In an embodiment the allele is a C394T mutation according to the sequence of SEQ ID NO:5.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome characterized by an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 according to the sequence of SEQ ID NO:8.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome on the basis of cancer being characterized by having an IDH1 allele described herein, *e.g.*, an IDH1 allele having Cys at residue 132 (SEQ ID NO:8).

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a glioma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically, Lys, Gly, Met, Trp or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a glioma, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys, Gly, Met, Trp, Thr, or Ser at residue 172 (SEQ ID NO:10), specifically Lys, Gly, Met, Trp, or Ser; or Lys or Met.

In an embodiment the method comprises selecting a subject having a glioma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is a prostate cancer, *e.g.*, prostate adenocarcinoma, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in



described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having a prostate cancer, *e.g.*, prostate adenocarcinoma, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is ALL, *e.g.*, B-ALL or T-ALL, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having ALL, *e.g.*, B-ALL or T-ALL, on the basis of the cancer being characterized by increased levels

of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is AML, characterized by a mutation, or preselected allele, of IDH2 associated with an alpha hydroxy neoactivity, e.g., 2HG neoactivity. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having AML, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having AML, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, e.g., 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, e.g., 2HG, *e.g.*, R-2HG.

In an embodiment the cell proliferation-related disorder is myelodysplasia or myelodysplastic syndrome, characterized by a mutation, or preselected allele, of IDH2. *E.g.*, in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. *E.g.*, the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, wherein the cancer is characterized by having an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by an IDH2 allele described herein, *e.g.*, an IDH2 allele having Lys or Met at residue 172 (SEQ ID NO:10).

In an embodiment the method comprises selecting a subject having myelodysplasia or myelodysplastic syndrome, on the basis of the cancer being characterized by increased levels of an alpha hydroxy neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG. In an embodiment the method comprises evaluating a serum or blood sample for increased alpha neoactivity product, *e.g.*, 2HG, *e.g.*, R-2HG.

In another aspect the invention features a pharmaceutical composition of an inhibitor (*e.g.*, a small molecule or a nucleic acid-based inhibitor) described herein.

In an embodiment a mutant protein specific reagent, *e.g.*, an antibody that specifically binds an IDH mutant protein, *e.g.*, an antibody that specifically binds an IDH1-R132H mutant protein, can be used to detect neoactive mutant enzyme see, for example, that described by Y.Kato et al., "A monoclonal antibody IMab-1 specifically recognizes IDH1<sup>R132H</sup>, the most common glioma-derived mutation: (Kato, Biochem. Biophys. Res. Commun. (2009), which is hereby incorporated by reference in its entirety.

In another aspect, the invention features, a method of evaluating a candidate compound, *e.g.*, for the ability to inhibit a neoactivity of a mutant enzyme, *e.g.*, for use as an anti-proliferative or anti-cancer agent. In an embodiment the mutant enzyme is an IDH, *e.g.*, an IDH1 or IDH2 mutant, *e.g.*, a mutant described herein. In an embodiment the neoactivity is alpha hydroxy neoactivity, *e.g.*, 2HG neoactivity. The method comprises:

optionally supplying the candidate compound;

contacting the candidate compound with a mutant enzyme having a neoactivity, or with another enzyme, a referred to herein as a proxy enzyme, having an activity, referred to herein as a proxy activity, which is the same as the neoactivity (or with a cell or cell lysate comprising the same); and

evaluating the ability of the candidate compound to modulate, *e.g.*, inhibit or promote, the neoactivity or the proxy activity, thereby evaluating the candidate compound.

In an embodiment the mutant enzyme is a mutant IDH1, *e.g.*, an IDH1 mutant described herein, and the neoactivity is an alpha hydroxy neoactivity, *e.g.*, 2HG

neoactivity. Mutations associated with 2HG neoactivity in IDH1 include mutations at residue 132, e.g., R132H, R132C, R132S, R132G, R132L, or R132V, more specifically, R132H or R132C.

In an embodiment the mutant enzyme is a mutant IDH2, e.g., an IDH2 mutant described herein, and the neoactivity is an alpha hydroxy neoactivity, e.g., 2HG neoactivity. Mutations associated with 2HG neoactivity in IDH2 include mutations at residue 172, e.g., R172K, R172M, R172S, R172G, or R172W.

In an embodiment the method includes evaluating the ability of the candidate compound to inhibit the neoactivity or the proxy activity.

In an embodiment the method further comprises evaluating the ability of the candidate compound to inhibit the forward reaction of non-mutant or wild type enzyme activity, e.g., in the case of IDH, e.g., IDH1 or IDH2, the conversion of isocitrate to  $\alpha$ -ketoglutarate (or an intermediate thereof, including the reduced hydroxyl intermediate).

In an embodiment, the contacting step comprises contacting the candidate compound with a cell, or a cell lysate thereof, wherein the cell comprises a mutant enzyme having the neoactivity or an enzyme having the activity.

In an embodiment, the cell comprises a mutation, or preselected allele, of a mutant IDH1 gene. E.g., in an embodiment, the IDH1 allele encodes an IDH1 having other than an Arg at residue 132. E.g., the allele can encode His, Ser, Cys, Gly, Val, Pro or Leu, or any other residue described in Yan *et al.*, at residue 132, according to the sequence of SEQ ID NO:8 (see also **FIG. 21**), specifically His, Ser, Cys, Gly, Val, or Leu.

In an embodiment the allele encodes an IDH1 having His at residue 132.

In an embodiment the allele encodes an IDH1 having Ser at residue 132.

In an embodiment the allele is an Arg132His mutation, or an Arg132Ser mutation, according to the sequence of SEQ ID NO:8 (see **FIGs. 2** and **21**).

In an embodiment, the cell comprises a mutation, or preselected allele, of a mutant IDH2 gene. E.g., in an embodiment, the IDH2 allele encodes an IDH2 having other than an Arg at residue 172. E.g., the allele encodes Lys, Gly, Met, Trp, Thr, Ser, or any residue described in described in Yan *et al.*, at residue 172, according to the sequence of SEQ ID NO:10(see also **Fig. 22**), specifically, Lys, Gly, Met, Trp, or Ser. In an embodiment the allele encodes an IDH2 having Lys at residue 172. In an embodiment the allele encodes an IDH2 having Met at residue 172.