

Catechin gallates are NADP⁺-competitive inhibitors of glucose-6-phosphate dehydrogenase and other enzymes that employ NADP⁺ as a coenzyme

Eui Seok Shin,^a Jiyoung Park,^b Jae-Min Shin,^{c,d} Dooho Cho,^d Si Young Cho,^a Dong Wook Shin,^a Mira Ham,^b Jae Bum Kim^b and Tae Ryong Lee^{a,*}

^aR&D Center, AmorePacific Corporation, 314-1 Bora-Dong, Giheung-Gu, Yongin, Gyeonggi 446-729, Republic of Korea

^bSchool of Biological Sciences, Seoul National University, San 56-1, Sillim-Dong, Kwanak-Gu, Seoul 151-742, Republic of Korea

^cDepartment of Bioinformatics, Soongsil University, 1-1 Sangdo-Dong, Dongjak-Gu, Seoul 156-743, Republic of Korea

^dSBscience Inc., 3rd Fl. Sungok-Building, 4-1 Sunae-Dong, Bundang-Gu, Gyeonggi 463-825, Republic of Korea

Received 9 January 2008; revised 1 February 2008; accepted 5 February 2008

Available online 14 February 2008

Abstract—Recent studies have shown that glucose-6-phosphate dehydrogenase (G6PD) is an effectual therapeutic target for metabolic disorders, including obesity and diabetes. In this study, we used *in silico* and conventional screening approaches to identify putative inhibitors of G6PD and found that galled catechins (EGCG, GCG, ECG, CG), but not ungalled catechins (ECG, GC, EC, C), were NADP⁺-competitive inhibitors of G6PD and other enzymes that employ NADP⁺ as a coenzyme, such as IDH and 6PGD.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

Oxidative stress and associated inflammatory processes are believed to play important roles in the pathogenesis of metabolic syndromes as well as major age-related diseases.^{1–3} NADPH is an essential coenzyme for several enzymes that generate oxygen-free radicals, including NADPH oxidase, nitric oxide synthase, and the cytochrome P450 monooxygenases.⁴ Thus, a reduction in NADPH production could result in a significant change in cellular oxidative stress. In addition, NADPH is an essential element in lipogenesis^{5,6} and contributes to fatty acid and cholesterol synthesis by supplying reducing power. Therefore, NADPH-producing enzymes might be closely associated with oxidative stress, chronic inflammatory signals, and lipid metabolism disorders.

NADPH is produced by reduction of NADP⁺ in biochemical reactions catalyzed by several enzymes, including malic enzyme (ME), isocitrate dehydrogenase

(IDH), and glucose-6-phosphate dehydrogenase (G6PD) and 6-phosphogluconate dehydrogenase (6PGD), which are the first two enzymes of the pentose phosphate pathway (PPP).⁶ Among the four NADPH-producing enzymes, G6PD is the rate-limiting enzyme of PPP, and is highly conserved in most mammalian species.⁷ G6PD, which is expressed ubiquitously, is implicated in various cell functions, including cell growth, survival, and redox regulation, and its deficiency causes hemolytic anemia and neonatal jaundice.⁸ Interestingly, hormonal or nutritional regulation of G6PD was restricted to liver and adipose tissues.⁶ Hepatic G6PD is regulated by nutritional signals, including a high-carbohydrate diet, polyunsaturated fatty acids, and hormonal signals such as insulin, glucagon, thyroid hormone, and glucocorticoids.^{6,7} Furthermore, G6PD-deficient patients show a decrease in lipogenic rate and serum lipoprotein concentrations, implying the importance of G6PD in fatty acid synthesis.^{9,10} Recent studies have elucidated novel roles of adipose tissue G6PD in the etiology of metabolic disorders. G6PD expression is highly increased in obese subjects including *ob/ob*, *db/db*, and diet-induced obese mice, and high expression of G6PD in adipocytes is tightly associated with lipid dysregulation, oxidative stress, and the chronic inflammation found in obese or

Keywords: Catechin gallate; Glucose-6-phosphate dehydrogenase; Isocitrate dehydrogenase; 6-Phosphogluconate dehydrogenase; NADP.

* Corresponding author. Tel: +82 31 280 5961; fax: +82 31 899 2595; e-mail: trlee@amorepacific.com

diabetic subjects.^{11,12} These observations indicate that G6PD is a potential therapeutic target for obesity and/or diabetes-related diseases.

Dehydroepiandrosterone (DHEA) is a well-known, uncompetitive inhibitor of G6PD.^{13–15} It has anti-oxidative, anti-carcinogenic, anti-obesity, and anti-aging properties.^{16,17} Use of DHEA as an anti-obesity drug is hampered by the requirement of high oral dosage and its easy conversion into various active androgens. Thus, it is expected that finding more efficient inhibitors of G6PD could lead to potent therapeutic drugs against obesity and/or diabetes.

In this study, both in silico and conventional screening approaches targeting the coenzyme (NADP⁺) and substrate (glucose-6-phosphate, G6P) binding sites of G6PD were performed in an effort to discover candidate G6PD inhibitors, and catechin gallates were identified as potent NADP⁺-competitive inhibitors of G6PD.

2. Results and discussion




2.1. Identification of EGCG as a potent inhibitor of G6PD through virtual screening approach

From a virtual screening experiment targeting the coenzyme (NADP⁺) and substrate (G6P) binding sites of G6PD, 250 candidate compounds were selected from a collection of three million commercially available compounds, and purchased from several chemical library distribution companies. These candidate compounds were used in an in vitro G6PD inhibition assay and eight compounds showed more than 50% inhibition at less than 100 $\mu\text{mol/L}$. Among eight compounds, (–)-epigallocatechin gallate (EGCG) was identified as the most potent inhibitor of G6PD (Table 1 and Fig. 1). EGCG is the most abundant polyphenolic catechin isolated from green tea, which exhibits profound pharmacological activities including anti-oxidant activity, inhibition of cell proliferation, inhibition of ultraviolet B (UVB)-induced inflammatory responses, modulation of cell cycle regulation, anti-cholesterolemic activity, suppression of angiogenesis, and anti-carcinogenic effect.^{18–21}

2.2. Galloyl moiety of catechins is an essential structural feature in the inhibition of G6PD

To elucidate the structure–activity relationship of the inhibitory effects of EGCG on G6PD, the inhibition kinetics of EGCG and other green tea catechins were investigated. Interestingly, the inhibitory activity of green tea catechins on G6PD was restricted to gallated catechins, such as EGCG, (–)-gallocatechin gallate (GCG), (–)-epicatechin gallate (ECG), and (–)-catechin gallate (CG) when compared to ungallated catechins, such as (–)-epigallocatechin (EGC), (–)-gallocatechin (GC), (–)-epicatechin (EC), and (–)-catechin (C) (Fig. 1 and Table 2). All gallated catechins exhibited similar IC₅₀ values (0.18–0.25 $\mu\text{mol/L}$) for the inhibition of G6PD (Table 2). Therefore, the galloyl moiety of cat-

Table 1. IC₅₀ values of G6PD inhibitors

Structure	IC ₅₀ ($\mu\text{mol/L}$)
	0.25
	56.37
	21.76





Figure 1. Dose–response curves for EGCG and EGC on the rate of G6PD catalysis. The activity of G6PD was measured in the presence of various concentrations of EGCG (■) and EGC (●).

of G6PD, while the 5'-hydroxyl group on the B ring and the stereochemistry of the 2-position of the catechin skeleton are not necessary for inhibition of G6PD. Several supportive studies have shown that the galloyl moiety has active biological features. Tian and co-workers

Table 2. G6PD, 6PGD, and IDH IC₅₀ values for gallated catechins

Compound	General structure	R1	R2	IC ₅₀ (μmol/L)		
				G6PD	6PGD	IDH
EC	A	H	H	≥1000	≥1000	≥1000
EGC	A	OH	H	≥1000	≥1000	≥1000
ECG	A	H	3,4,5-Trihydroxybenzoyl	0.18 ± 0.01	1.21 ± 0.13	10.8 ± 1.66
EGCG	A	OH	3,4,5-Trihydroxybenzoyl	0.25 ± 0.02	0.72 ± 0.07	6.44 ± 1.12
CG	B	H	3,4,5-Trihydroxybenzoyl	0.24 ± 0.01	1.28 ± 0.08	6.62 ± 0.63
GCG	B	OH	3,4,5-Trihydroxybenzoyl	0.23 ± 0.02	1.45 ± 0.08	2.72 ± 0.21
GC	B	OH	H	≥1000	≥1000	≥1000
C	B	H	H	≥1000	≥1000	≥1000

synthase (FAS), and that the galloyl moiety is the critical structural feature in the inhibition of the β -ketoacyl reductase activity of FAS via a reversible association with the NADPH-binding site or with an adjacent area of the β -ketoacyl reductase of FAS.^{22,23} In addition, EGCG and ECG, but not EGC and EC, are potent inhibitors of glutamate dehydrogenase (GDH) with EC₅₀s in the nanomolar range. EGCG is a non-competitive inhibitor of both GDH substrates (NADH and 2-oxoglutarate), but acts in an allosteric manner.²⁴

2.3. Catechin gallates are NADP⁺-competitive inhibitors of G6PD and other enzymes that employ NADP⁺ as a coenzyme

To examine the manner in which gallated catechins inhibit G6PD activity, various concentrations of EGCG and CG were added to reactions containing various concentrations of NADP⁺ and G6P. Both EGCG and CG are competitive inhibitors of NADP⁺, but are uncompetitive inhibitors of G6P (Fig. 2). Based on the NADP⁺-competitive inhibition patterns of gallated catechins, it was proposed that gallated catechins could act as general inhibitors of enzymes that employ NADP⁺ as a coenzyme. Gallated catechins were indeed potent inhibitors of 6PGD and IDH, enzymes which both employ NADP⁺ as a coenzyme (Table 2).

It is well known that green tea catechins affect the reduction of body weight and prevent obesity-related metabolic disorders such as diabetes, hyperlipidemia, and hypertension in various animal models and in humans.^{25,26} Although catechins have been shown to be effective inhibitors of G6PD, it has not been clearly determined whether the therapeutic effects of catechins on metabolic disorders are directly associated with G6PD inhibition. Gallated catechins are not only inhibitors of G6PD, but are also inhibitors of 6PGD and IDH, which use NADP⁺ as a coenzyme. In addition, gallated catechins are good inhibitors of FAS and

and insulin secretion, respectively. Green tea catechins are also reported to be inhibitors of pancreatic phospholipase A2 (PLA2), and were found to inhibit the intestinal absorption of lipids in ovariectomized rats.²⁷ Thus, although gallated catechins are effective inhibitors of G6PD in vitro and show good anti-obesity effects in vivo, the extent of the effects that are directly associated with the inhibition of G6PD by catechins remains unclear.

2.4. Effects of EGCG on endogenous dehydrogenase activity in 3T3-L1 adipocytes lysates

To further investigate the inhibitory effects of EGCG, ECG, GCG, and CG on the production of NADPH by G6PD and 6PGD in adipocytes, measurement of NADPH production was performed using cell lysates from differentiated 3T3-L1 adipocytes. As shown in Figure 3A, EGCG, ECG, GCG, and CG effectively suppressed NADPH production in 3T3-L1 adipocytes with IC₅₀ values near 25 μmol/L. However, EGC, EC, GC, and C did not suppress NADPH production in 3T3-L1 adipocytes (Fig. 3B). We also compared the inhibitory effect of EGCG with that of DHEA, a well-known uncompetitive inhibitor of G6PD activity.^{13–16} As shown in Figure 3C, both DHEA and EGCG inhibited NADPH production in a dose-dependent manner, but DHEA showed only 40% maximum inhibition at concentrations above 100 μmol/L. This difference in inhibition patterns between EGCG and DHEA represents a difference in inhibition mechanisms. Consistent with the observation of differing inhibition mechanisms, EGCG also inhibited 6PGD activity in a dose-dependent manner, while DHEA did not inhibit 6PGD activity (Fig. 3D).

EGCG, ECG, GCG, and CG were significantly better inhibitors of endogenous NADPH production when compared to DHEA. However, considering the very low IC₅₀ values (0.18–0.25 μmol/L) in activity assay




Figure 2. Inhibition kinetics of EGCG and CG with respect to G6P and NADP^+ . G6PD was analyzed with respect to NADP^+ at three concentrations of EGCG and CG. (A) EGCG: 0 $\mu\text{mol/L}$ (\square), 0.5 $\mu\text{mol/L}$ (\triangle), and 1.0 $\mu\text{mol/L}$ (\circ). (B) CG: 0 $\mu\text{mol/L}$ (\square), 5.0 $\mu\text{mol/L}$ (\triangle), and 10.0 $\mu\text{mol/L}$ (\circ). G6PD was analyzed with respect to G6P at three concentrations of EGCG and CG. (C) EGCG: 0 $\mu\text{mol/L}$ (\square), 1.0 $\mu\text{mol/L}$ (\triangle), and 2.0 $\mu\text{mol/L}$ (\circ). (D) CG: 0 $\mu\text{mol/L}$ (\square), 5.0 $\mu\text{mol/L}$ (\triangle), and 10.0 $\mu\text{mol/L}$ (\circ).

ing that catechin gallates showed 100-fold higher IC_{50} values in the NADPH production assay using cell lysates of differentiated 3T3-L1 adipocytes. This difference in IC_{50} values may be partially due to the presence of other proteins, such as IDH, FAS, GDH, and others, which bind with EGCG and consequently reduce the amount of available free EGCG in the 3T3-L1 adipocyte cell lysate. The total plasma concentration after 50 mg EGCG oral intake (estimated amount of EGCG in a cup of green tea) in human is approximately 0.3 $\mu\text{mol/L}$.²⁸ Catechin gallate IC_{50} values of approximately 25 $\mu\text{mol/L}$ seem too high to suggest any physiological relevance to known *in vivo* activities of green tea catechins. However, unlike DHEA, catechin gallates are NADP^+ -competitive inhibitors of enzymes that employ NADP^+ as a coenzyme. Therefore, the *in vivo* EC_{50} values of catechin gallates will vary depending on the concentration of free NADP^+ in the target organ. The concentration of free NADP^+ in the target organ is difficult to measure. However, a reasonable estimation can be made based on several previous reports. Total concentration of NADP^+ and NADPH in rat liver is

ratio is about 0.005. A significant portion of the NADP^+ and NADPH is bound to protein; in the case of NADH , over 80% of NADH is protein bound. Therefore, the concentration of free NADP^+ would be much lower.^{29–31} The estimated free NADP^+ concentration in a target organ is approximately 0.1 $\mu\text{mol/L}$, which is more than 1000-fold lower than the concentration of NADP^+ used in the *in vitro* experiment. The high concentration of NADP^+ in the *in vitro* experiment is necessary to obtain a reasonable signal size for measurement. Therefore, EC_{50} values of catechin gallates could be much lower than 0.3 $\mu\text{mol/L}$. In addition, it was recently reported that multiple treatments with catechins showed synergistic effects.³² Thus, multiple treatments will further decrease their EC_{50} values and our results may, in fact, be physiologically relevant to known *in vivo* activities of green tea catechins in humans.

3. Conclusion

In this study, utilizing both *in silico* and conventional




Figure 3. Effects of catechins and DHEA on NADPH production. The production of NADPH by 3T3-L1 cell lysates was measured in the presence of various concentrations of catechins and DHEA. (A) EGCG (●), ECG (□), GCG (▲) and CG (×). (B) EGC (●), EC (□), GC (▲) and C (×). (C) EGCG (●) and DHEA (○). (D) The production of NADPH by 6PGD in 3T3-L1 cell lysates was measured in the presence of various concentrations of EGCG (●) and DHEA (○). Results are represented as means \pm SD of three-independent experiments.

chins, but not ungalated catechins, were NADP^+ -competitive inhibitors of G6PD and other enzymes that employ NADP^+ as a coenzyme. Although the extent of the effects that are directly attributable to the inhibition of each enzyme remains unclear, these results along with previous reports concerning the inhibition effects of green tea catechins on FAS, PLA2 and GDH explain how green tea catechins can display such broad in vivo activities against obesity and oxidative stress-related disorders. Catechin gallates showed somewhat high IC_{50} values in the NADPH production assay using cell lysates of differentiated 3T3-L1 adipocytes. However, these activities may still be physiologically relevant to known in vivo activities of green tea catechins in humans, due to NADP^+ -competitive inhibition of catechin gallates and low in vivo concentrations of free NADP^+ .

4. Experimental

4.1. Materials

Glucose-6-phosphate, sodium chloride, magnesium chloride, β -NADP, (–)-epigallocatechin gallate (EGCG), (–)-epigallocatechin (EGC), (–)-epicatechin gallate (ECG), (–)-epicatechin (EC), (–)-gallocatechin gallate (GCG), (–)-gallocatechin (GC), (–)-catechin gallate (CG), (–)-catechin (C), Glucose-6-phosphate dehydrogenase (EC 1.1.1.49) from bakers yeast, 6-phosphogluconic dehydrogenase from *Saccharomyces cerevisiae* (EC 1.1.1.44), 6-phosphogluconic dehydrogenase (EC 1.1.1.44) from porcine heart were purchased from Sigma–Aldrich.

(EC 1.1.1.42) from porcine heart were purchased from Sigma–Aldrich.

4.2. Modeling of the binding site of G6PD

Two binary complex structures of the G6PD human deletion mutant of Kotaka et al. with glucose-6-phosphate (G6P, PDB code: 2bhl), and NADP^+ (PDB code: 2bh9) were selected for virtual screening.³³ The superimposed structure of the two binary complexes is shown in Fig. S-1. For convenience in structure-based modeling, a combined structure was constructed by copying G6P from the PDB structure 2bhl into the PDB structure 2bh9, in a manner similar to that used by Kotaka et al.³³ (Figure S-2). The software program IDPharmo version 2.0 (Equispharm Inc., Seoul, Korea)³⁴ was used for virtual screening to search approximately three million commercially available library compounds in a period of eight hours, using a 3GHz, four-CPU Linux PC. IDPharmo is fingerprint-based virtual screening software, and three essential physicochemical features comprise its construction of reliable fingerprints: the hydrogen bond donor, the hydrogen bond acceptor, and the hydrophobic core. After several cycles of pharmacophore generation and refinement, 9 protein–ligand-binding features for the substrate- and coenzyme-binding sites of G6PD were obtained, as shown in Figure S-2. Based on the combination of these 9 ligand-binding features, 12 different pharmacophore models, termed PharmoMaps, were selected. Based on the 12 PharmoMaps, 1000 virtual compounds were

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.