



We're building a better ClinicalTrials.gov. Check it out and tell us what you think!

U.S. National Library of Medicine

ClinicalTrials.gov

[Find Studies](#) ▾

[About Studies](#) ▾

[Submit Studies](#) ▾

[Resources](#) ▾

[About Site](#) ▾

[PRS Login](#)

[Home](#) > [Search Results](#) > Study Record Detail

[Save this study](#)

Post-Gastric Bypass Hypoglycemia

The safety and scientific validity of this study is the responsibility of the study sponsor and investigators.

⚠ Listing a study does not mean it has been evaluated by the U.S. Federal Government. Read our [disclaimer](#) for details.

ClinicalTrials.gov Identifier: NCT01933490

[Recruitment Status](#) : Completed

[First Posted](#) : September 2, 2013

[Last Update Posted](#) : November 1, 2019

[View this study on
Beta.ClinicalTrials.gov](#)

Sponsor:

University of Minnesota

Information provided by (Responsible Party):

University of Minnesota

[Study Details](#)

[Tabular View](#)

[No Results Posted](#)

[Disclaimer](#)

[? How to Read a Study Record](#)

Study Description

Go to ▾

Brief Summary:

Post-gastric bypass hyperinsulinemic hypoglycemia is a recently described disorder occurring in some patients after gastric bypass surgery for obesity. The pathogenesis is incompletely understood but involves a robust insulin response to ingested carbohydrate. The resultant hyperinsulinemia sometimes produces hypoglycemia with neuroglycopenia,

confusion and even loss of consciousness. Various treatments have been recommended including low carbohydrate diets, coingestion of the medication acarbose with carbohydrate containing meals, partial pancreatectomy and even total pancreatectomy. None is completely satisfactory. We propose to test two new potential treatments. Using a design with random assignment of three conditions we plan to compare, in 10 patients with post-gastric bypass hyperinsulinemic hypoglycemia, a high carbohydrate test meal (control condition), a high carbohydrate test meal after pre-treatment with rapid acting aspart insulin (insulin condition), and a high fructose, low glucose test meal with carbohydrate and caloric content similar to the control meal (fructose condition).

<u>Condition or disease</u> <input type="checkbox"/>	<u>Intervention/treatment</u> <input type="checkbox"/>	<u>Phase</u> <input type="checkbox"/>
Hyperinsulinemic Hypoglycemia	Other: high carbohydrate test meal Other: high carbohydrate test meal after pre-treatment with rapid acting aspart insulin Other: high fructose , low glucose test meal with carbohydrate and caloric content similar to the control meal	Not Applicable

Detailed Description:

Post-gastric bypass hyperinsulinemic hypoglycemia is a recently described disorder occurring in some patients after gastric bypass surgery for obesity. The pathogenesis is incompletely understood but involves a robust insulin response to ingested carbohydrate. The resultant hyperinsulinemia sometimes produces hypoglycemia with neuroglycopenia, confusion and even loss of consciousness. Various treatments have been recommended including low carbohydrate diets, coingestion of the medication acarbose with carbohydrate containing meals, partial pancreatectomy and even total pancreatectomy. None is completely satisfactory. We propose to test two new potential treatments. Using a design with random assignment of three conditions we plan to compare, in 10 patients with post-gastric bypass hyperinsulinemic hypoglycemia, a high carbohydrate test meal (control condition), a high carbohydrate test meal after pre-treatment with rapid acting aspart insulin (insulin condition), and a high fructose, low glucose test meal with carbohydrate and caloric content similar to the control meal (fructose condition). The hypothesis to be tested are 1) pretreatment with aspart insulin will prevent, or at least reduce, the occurrence of hypoglycemia and 2) substitution of fructose for glucose in the test meal will prevent, or at least reduce, the occurrence of hypoglycemia. Plasma glucose and serum insulin will be sampled before and for four hours after the three test conditions. The primary study endpoint will be the occurrence or not of plasma glucose < 60 mg/dL after the test meals. The control meal will be compared to the insulin pre-treated test meal and, in a separate comparison, to the fructose test meal. Secondary endpoints will be comparisons between the control and active treatments in peak postprandial serum insulin, peak postprandial plasma glucose, nadir postprandial plasma glucose, and the 4-hr longitudinal course of plasma glucose measurements.

Study Design

Go to

Study Type : Interventional (Clinical Trial)

Actual Enrollment : 10 participants

Allocation: Randomized

Intervention Model: Parallel Assignment

Masking: None (Open Label)

Primary Purpose: Health Services Research

Official Title: Prevention of Hypoglycemia in Patients With Post-Gastric Bypass
Hyperinsulinemic Hypoglycemia

Study Start Date : August 2013

Actual Primary Completion Date : August 2014

Actual Study Completion Date : August 2014

Resource links provided by the National Library of Medicine



[MedlinePlus Genetics](#) related topics: [Congenital hyperinsulinism](#)

[MedlinePlus](#) related topics: [Hypoglycemia](#)

[Genetic and Rare Diseases Information Center](#) resources:

[Congenital Hyperinsulinism](#)

[U.S. FDA Resources](#)

Arms and Interventions

Go to

Arm <input type="checkbox"/>	Intervention/treatment <input type="checkbox"/>
a high carbohydrate test meal (control condition) a high carbohydrate test meal (control condition)	Other: high carbohydrate test meal Other: high carbohydrate test meal after pre-treatment with rapid acting aspart insulin Other: high fructose , low glucose test meal with carbohydrate and caloric content similar to the control meal
Active Comparator: high carbohydrate test meal after pre-treatment a high carbohydrate test meal after pre-treatment with rapid acting aspart insulin (insulin condition)	Other: high carbohydrate test meal Other: high carbohydrate test meal after pre-treatment with rapid acting aspart insulin Other: high fructose , low glucose test meal with carbohydrate and caloric content similar to the control meal
Active Comparator: high fructose low glucose test meal high fructose , low glucose test meal with carbohydrate and caloric content similar to the control meal (fructose condition)	Other: high carbohydrate test meal Other: high carbohydrate test meal after pre-treatment with rapid acting aspart insulin Other: high fructose , low glucose test meal with carbohydrate and caloric content similar to the control meal

Outcome Measures

Go to ▼

Primary Outcome Measures □ :

1. The primary study endpoint will be occurrence or not of plasma glucose < 60 mg/dL during the 4 hours after the test meal (binary endpoint). [Time Frame: 4 hours after meal]

The primary study endpoint will be occurrence or not of plasma glucose < 60 mg/dL during the 4 hours after the test meal (binary endpoint).

Eligibility Criteria

Go to ▼

Information from the National Library of Medicine



Choosing to participate in a study is an important personal decision. Talk with your doctor and family members or friends about deciding to join a study. To learn more about this study, you or your doctor may contact the study research staff using the contacts provided below. For general information, [Learn About Clinical Studies](#).

Ages Eligible for Study: 21 Years and older (Adult, Older Adult)

Sexes Eligible for Study: All

Accepts Healthy Volunteers: No

Criteria

Inclusion Criteria:

- Participants must be at least 21 years of age
- History of postprandial hypoglycemia with neuroglycopenia occurring one year or more after gastric bypass surgery
- History of spontaneous correction of hypoglycemia
- Normal fasting plasma glucose and serum insulin after a carbohydrate containing mixed meal, demonstration of serum insulin > 50u/UL and plasma glucose < 50mg/dL

Exclusion Criteria:

- Under 21 years of age

Contacts and Locations

Go to ▼

Information from the National Library of Medicine

To learn more about this study, you or your doctor may contact the study research staff using the contact information provided by the sponsor.

Please refer to this study by its ClinicalTrials.gov identifier (NCT number):

NCT01933490

Locations**United States, Minnesota**

University of Minnesota

Minneapolis, Minnesota, United States, 55455

Sponsors and Collaborators

University of Minnesota

Investigators

Principal Investigator: John Bantle, MD University of Minnesota

More Information

Go to ▼

Publications automatically indexed to this study by ClinicalTrials.gov Identifier (NCT Number):

[Bantle AE, Wang Q, Bantle JP. Post-Gastric Bypass Hyperinsulinemic Hypoglycemia: Fructose is a Carbohydrate Which Can Be Safely Consumed. J Clin Endocrinol Metab. 2015 Aug;100\(8\):3097-102. doi: 10.1210/jc.2015-1283. Epub 2015 Jun 2.](#)

Responsible Party: University of Minnesota
ClinicalTrials.gov Identifier: [NCT01933490](#) [History of Changes](#)
Other Study ID Numbers: 1306M37181
First Posted: September 2, 2013 [Key Record Dates](#)
Last Update Posted: November 1, 2019
Last Verified: October 2019

Keywords provided by University of Minnesota:

hyperinsulinemic hypoglycemia

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.