

Stool DNA Testing For Colon Cancer

Date: December 13, 2006

Source: The Mount Sinai Hospital / Mount Sinai School of Medicine

Summary: A prospective, multi-center study of stool DNA testing has found that a new test demonstrated an 88 percent sensitivity for colorectal cancer, and with equal detection across all stages of cancer, regardless of the cancer's location in the colon. The study was published online in the American Gastroenterological Association's journal, Clinical Gastroenterology and Hepatology.

FULL STORY

Mount Sinai School of Medicine and EXACT Sciences Corporation have announced the publication of results from a prospective, multi-center study of stool DNA testing. The study found that the test demonstrated an 88% sensitivity for colorectal cancer, and with equal detection across all stages of cancer, regardless of the cancer's location in the colon.

The study was published online in the American Gastroenterological Association's journal, Clinical Gastroenterology and Hepatology, and will appear in the January 2007 print issue.

"This study confirms that stool-based DNA technologies can achieve high sensitivities for detecting colorectal cancer," stated Steven Itzkowitz, M.D., principal investigator and Professor and Associate Director of Gastroenterology at Mount Sinai School of Medicine.

"For those individuals who are unwilling or unable to undergo colonoscopy, stool DNA testing offers a valuable and patient-friendly screening option. These results also underscore that as new markers and technologies are developed and validated, they can readily be incorporated into existing stool DNA tests to improve cancer detection and, ultimately, patient outcomes."

The published study, entitled "Improved Fecal DNA Test for Colorectal Cancer Screening," evaluated 162 patients, 40 individuals with cancer and 122 individuals with normal colonoscopies.

An enhanced marker panel, using a refined DNA capture and stabilization process, detected 88% of cancers with a specificity of 82%.

"This publication is further validation that stool DNA technology offers a powerful tool for physicians and patients in detecting colorectal cancer," said Don Hardison, EXACT Sciences' President and

"Without new, non-invasive approaches such as stool DNA testing, it will be difficult to increase current colorectal screening and decrease mortality rates, a major goal for our company as well as a mandate of the American Cancer Society."

Colorectal cancer is the second leading cause of cancer deaths in the U.S. and more than half of the over 80 million people over the age of 50 have never been screened. A recent NCI study published in the American Cancer Society's journal, *Cancer*, projected that, using traditional screening approaches, U.S. screening and mortality reduction goals cannot be achieved even under the most optimistic of scenarios. The report concluded that consideration of new screening technologies, such as stool DNA testing, is warranted.

Story Source:

Materials provided by **The Mount Sinai Hospital / Mount Sinai School of Medicine**. *Note: Content may be edited for style and length.*

Cite This Page:

MLA	APA	Chicago
-----	-----	---------

The Mount Sinai Hospital / Mount Sinai School of Medicine. "Stool DNA Testing For Colon Cancer." ScienceDaily. ScienceDaily, 13 December 2006. <www.sciencedaily.com/releases/2006/12/061213104115.htm>.

Explore More

from ScienceDaily

RELATED STORIES

New Study Provides Evidence for Three-Year Interval for Multi-Target Stool DNA Screening for Those at Average Risk of Colon Cancer

Nov. 29, 2022 — A scientific study exploring the appropriate interval for colorectal cancer screening via non-invasive multi-target stool DNA testing for individuals with average risk for the disease reported ...

Study Suggests Blood Test Could Guide Precision Treatment in Bladder Cancer

June 21, 2021 — A blood test that can detect tiny amounts of circulating cancer DNA may be able to identify risk of cancer recurrence and guide precision treatment in bladder cancer following surgery, according to a ...

Average-Risk Individuals May Prefer Stool-Based Test Over Colonoscopy for Cancer Screening

Apr. 22, 2021 — When given a choice, most individuals with an average risk of colorectal cancer said they would prefer a stool-based screening test for colorectal cancer over colonoscopy, the method most often ...

Gut Microbiota Imbalance Promotes the Onset of Colorectal Cancer

Nov. 18, 2019 — Researchers have demonstrated that an imbalance in the gut microbiota, also known as 'dysbiosis', promotes the onset of colorectal cancer. The teams demonstrated that transplanting fecal flora from ...