

Look to Paddock for All of Your Compounding Needs

TOPICALS
INJECTABLES
ORAL LIQUIDS
ORAL SOLIDS
SUPPOSITORIES
COMPOUNDING

ORA-SWEET[®]

Flavored Syrup Vehicle

ORA-SWEET[®]

Paddock ORA-SWEET

Size	NDC
473 ML (One Pint)	0574-0304-16

Ora-Sweet is a syrup vehicle used to simplify the process of flavoring and sweetening extemporaneous compounded oral preparations. Ora-Sweet allows the pharmacist to formulate elegant, sweetened products with minimum time and maximum dependability. Ora-Sweet is the modern version of simple syrup flavored with a citrus-berry blend for a highly palatable taste.



Applications

Ora-Sweet may be used alone or in combination with other agents. Ora-Sweet will retain its flavoring properties when diluted up to 50% with water or suspending agents. Its versatility makes it ideal for the flavoring of:

- Pediatric preparations
- Geriatric preparations

How to Use

Ora-Sweet is the ideal flavoring and sweetening agent for many suspensions, but it is specially formulated to complement Paddock's suspending vehicle Ora-Plus. Ora-Sweet and Ora-Plus can be combined in a 50/50 ratio to produce a pleasant tasting elegant suspension.

Properties

Ora-Sweet contains sucrose which acts as a sweetening agent. Small amounts of glycerin and sorbitol are added to prevent "cap lock" problems common to most syrups. Its flavoring agents help increase palatability. Ora-Sweet is buffered to a slightly acidic pH to help diminish degradation of medicinal agents through oxidation.

Ingredients

Purified water, sucrose, glycerin, sorbitol, and flavoring. Buffered with citric acid and sodium phosphate. Preserved with methylparaben and potassium sorbate.

Specifications

Appearance: Clear liquid with a slight tint

pH: Approximately 4.2

Taste: Sweet citrus-berry flavor

Osmolality: 3240 mOsm/Kg

Order through your wholesaler. For additional information please contact Paddock Laboratories at 1-800-328-5113 or www.paddocklabs.com

ORA-SWEET Formulating and Compounding Examples

CHLOROQUINE PHOSPHATE 15 mg/mL, 100 mL²

Chloroquine Phosphate 500 mg/tablet	3 tablets
Ora-Plus	50 mL
Ora-Sweet*	q.s. 100 mL

- Expiration 60 days
- Protect from light
- Shake well before using

METOLAZONE 1 mg/mL, 120 mL¹

Metolazone 10 mg/tablet	12 tablets
Ora-Plus	60 mL
Ora-Sweet*	q.s. 120 mL

- Expiration 60 days
- Protect from light
- Shake well before using

PROCAINAMIDE HCl 50 mg/mL, 120 mL¹

Procainamide HCl 250 mg/capsule	24 capsules
Ora-Plus	60 mL
Ora-Sweet*	q.s. 120 mL

- Expiration 60 days
- Protect from light
- Shake well before using

KETOCONAZOLE 20 mg/mL, 120 mL¹

Ketoconazole 200 mg/tablet	12 tablets
Ora-Plus	60 mL
Ora-Sweet*	q.s. 120 mL

- Expiration 60 days
- Protect from light
- Shake well before using

METRONIDAZOLE 50 mg/mL, 120 mL¹

Metronidazole powder	6 grams
Ora-Plus	60 mL
Ora-Sweet*	q.s. 120 mL

- Expiration 60 days
- Protect from light
- Shake well before using

SPIRONOLACTONE 25 mg/mL, 120 mL¹

Spirolactone 25 mg/tablet	120 tablets
Ora-Plus	60 mL
Ora-Sweet*	q.s. 120 mL

- Expiration 60 days
- Protect from light
- Shake well before using



General Directions For Compounding ORA-SWEET*/ORA-PLUS Suspensions

ORA-SWEET and ORA-PLUS should be combined in a 50/50 ratio to achieve optimal formulation.

- Calculate the total amount of active ingredient and volume of solution needed to create a suspension of the proper concentration. (Calculations are provided in the examples above.) *It may be wise to add approximately 10% overage for compounding losses.*
- Crush tablets with a mortar and pestle to a fine powder. If the compound is already in the powder form, use the mortar and pestle to smooth out the powder.
- Add a small amount of **Ora-Plus** and triturate to a thick, smooth paste. Add the remainder of **Ora-Plus** by geometric dilution. *The amount of Ora-Plus should be 50% of the total solution.*
- Bring the suspension to a final volume using **Ora-Sweet***. Mix briefly with a mortar and pestle until a uniform suspension is formed.
- Dispense in a tight, light resistant amber bottle with appropriate labeling.
- Label with an expiration date. *If the stability of the medication in an oral suspension or syrup is unknown, conservative dating is suggested.*

- Depending on the medicinal agent used, label containers "Shake Well Before Using," "Protect From Light," and "Keep Refrigerated."

*Ora-Sweet may be substituted with Ora-Sweet SF. Both products may be used in a 1:1 ratio with Ora-Plus.

¹American Journal of Health-System Pharmacists 1996; 53:2073-8.

²American Journal of Health-System Pharmacists 1998; 55:1915-20.

Contraindications

Ora-Sweet is contraindicated in persons who have shown hypersensitivity to any of the listed ingredients.

All information presented is intended to demonstrate the application of Paddock vehicles for compounding. Paddock products are sold on the understanding that purchasers will make their own determinations as to the suitability, safety and effectiveness of their applications. The uses presented by Paddock are only to assist our customers in exploring possible applications. All information and data presented are believed to be accurate and reliable, but are presented without the assumption of any liability by Paddock Laboratories. Paddock Laboratories does not warrant against infringements of patents of third parties by reason of any uses made of the vehicles in combination with other material or in the operation of any process, and purchasers assume all risks of patent infringement by reason of any such use, combination or operation.