
December 2013 Volume 119, Number 6 ISSN 0003-3022



ANESTHESIOLOGY The Journal of the American Society of Anesthesiologists, Inc. • anesthesiology.org




Including 11 Articles and 1 Editorials from or about

Find authenticated court documents without watermarks at docketalarm.com.

DOCKE

ANESTHESIOLOGY





ON THE COVER:

We continue our end-of-the-year special focus on an institution which has developed and maintained research excellence in the specialty. This year a majority of the December issue comes from the department of anesthesiology at the University of Michigan (Ann Arbor, Michigan). (The heli-photo of the University of Michigan medical center was provided by Dale Fisher.)

1233 1236
1236
1239
1241
1244
1247
1249

Altered States: Psychedelics and Anesthetics Eduardo E. Icaza and George A. Mashour 1255

- ♦ Refers to This Month in Anesthesiology
- Refers to Editorial Views

RM

DOCKE

Δ

 This article is from or about the University of Michigan

Find authenticated court documents without watermarks at docketalarm.com.

See Supplemental Digital Content

CME Article

CONTENTS

An Analysis of Methodologies That Can Be Used to Validate if a Perioperative Surgical Home Improves the Patient-centeredness, Evidence-based Practice, Quality, Safety, and Value of Patient Care 1261

Thomas R. Vetter, Nataliya V. Ivankova, Lee A. Goeddel, Gerald McGwin, Jr., and Jean-Francois Pittet for the UAB Perioperative Surgical Home Group

PERIOPERATIVE MEDICINE

♦ M Increased Risk of Intraoperative Awareness in Patients with a History of Awareness Amrita Aranake, Stephen Gradwohl, Arbi Ben-Abdallah, Nan Lin, Amy Shanks, Daniel L. Helsten, David B. Glick, Eric Jacobsohn, Alex J. Villafranca, Alex S. Evers, Michael S. Avidan, and George A. Mashour In a matched cohort analysis of patients drawn from three trials including over 25,000 patients, those with a history of intraoperative awareness had a five-fold increased incidence of awareness compared with propensity-matched controls who did not have a history of awareness. Anesthetic management did not differ between the cohorts; in view of the likely increased risk of awareness, clinicians should consider modifying anesthetic management in patients with a history of awareness. ♦ Failure of the Laryngeal Mask Airway Unique[™] and Classic[™] in the Pediatric Surgical Patient: A Study of Clinical Predictors and Outcomes 1284

Michael R. Mathis, Bishr Haydar, Emma L. Taylor, Michelle Morris, Shobha V. Malviya, Robert E. Christensen, Satya-Krishna Ramachandran, and Sachin Kheterpal

This study evidenced that laryngeal mask airway failures occur in 0.86% of 11,910 pediatric anesthesia cases with its planned use, and are independently associated with ear/nose/throat procedures, admission status, prolonged surgical duration, airway abnormalities, and patient transport.

Six Percent Hydroxyethyl Starch 130/0.4 (Voluven[®]) versus 5% Human Serum Albumin for Volume Replacement Therapy during Elective Open-heart Surgery in Pediatric Patients 1296 Philippe Van der Linden, Andrée De Villé, Anna Hofer, Martina Heschl, and Hans Gombotz

This randomized clinical trial demonstrated equivalent efficacy of human albumin versus hydroxyethyl starch for volume replacement in 61 children undergoing cardiac surgery. Although the trial was not powered to assess safety, outcomes measures were not different between the groups.

M Patient Selection for Day Case-eligible Surgery: Identifying Those at High Risk for **Major Complications**

Michael R. Mathis, Norah N. Naughton, Amy M. Shanks, Robert E. Freundlich, Christopher J. Pannucci, YiJia Chu, Jason Haus, Michelle Morris, and Sachin Kheterpal

In a review of nearly 250,000 cases in the National Surgery Quality Improvement Program, early perioperative morbidity and mortality occurred in approximately 1:1,000 patients. Predictors for morbidity and mortality were overweight or obesity, chronic obstructive pulmonary disease, history of transient ischemic attack/stroke, hypertension, prior cardiac surgical intervention, and prolonged operative time.

◇ M Predictors of Survival from Perioperative Cardiopulmonary Arrests: A Retrospective Analysis of 2,524 Events from the Get With The Guidelines-Resuscitation Registry

Satya Krishna Ramachandran, Jill Mhyre, Sachin Kheterpal, Robert E. Christensen, Kristen Tallman, Michelle Morris, and Paul S. Chan, for the American Heart Association's Get With The Guidelines-Resuscitation Investigators

One third of patients with perioperative cardiac arrests survive to hospital discharge. Two thirds of survivors have good neurological outcome. Shockable rhythms and asystolic arrests were associated with significant location-specific survival variance.

M Perioperative Metoprolol and Risk of Stroke after Noncardiac Surgery

George A. Mashour, Milad Sharifpour, Robert E. Freundlich, Kevin K. Tremper, Amy Shanks, Brahmajee K. Nallamothu, Phillip E. Vlisides, Adam Weightman, Lisa Matlen, Janna Merte, and Sachin Kheterpal

In this retrospective study, routine use of preoperative and intraoperative metoprolol—but not other β blockers—was associated with increased risk of stroke after noncardiac surgery. These data suggest that results of the POISE trial may

Find authenticated court documents without watermarks at docketalarm.com.

1340

1275

1310

1322

CONTENTS

M	Reconfiguration of Network Hub Structure after Propofol-induced Unconsciousness Heonsoo Lee, George A. Mashour, Gyu-Jeong Noh, Seunghwan Kim, and UnCheol Lee	1347
	Propofol reconfigures the brain's network hub structure. Reconfiguration of hub structure may explain the observed loss of frontal–parietal feedback connectivity.	
(11)	Incidence, Predictors, and Outcome of Difficult Mask Ventilation Combined with Difficult Laryngoscopy: A Report from the Multicenter Perioperative Outcomes Group Sachin Kheterpal, David Healy, Michael F. Aziz, Amy M. Shanks, Robert E. Freundlich, Fiona Linton, Lizabeth D. Martin, Jonathan Linton, Jerry L. Epps, Ana Fernandez-Bustamante, Leslie C. Jameson, Tyler Tremper, and Kevin K. Tremper, on behalf of the Multicenter Perioperative Outcomes Group (MPOG) Perioperative Clinical Research Committee	1360
	This study determined incidence of difficult mask ventilation combined with difficult laryngoscopy to be 0.4% of 176,679 adult cases, and succeeded in identifying 12 independent predictors for the critical situation.	
	CRITICAL CARE MEDICINE	
•	Propofol Limits Microglial Activation after Experimental Brain Trauma through Inhibition of Nicotinamide Adenine Dinucleotide Phosphate Oxidase Tao Luo, Junfang Wu, Shruti V. Kabadi, Boris Sabirzhanov, Kelsey Guanciale, Marie Hanscom, Juliane Faden, Katherine Cardiff, Charles Jeremy Bengson, and Alan I. Faden	1370
	Propofol reduces functional deficit after traumatic brain injury, in an animal model. These neuroprotective effects are mediated in part by inhibition of microglial nicotinamide adenine dinucleotide phosphate oxidase.	
	Critical Role of Interleukin-11 in Isoflurane-mediated Protection against Ischemic Acute Kidney Injury in Mice Ahrom Ham, Mihwa Kim, Joo Yun Kim, Kevin M. Brown, James Yeh, Vivette D. D'Agati, and H. Thomas Lee	1389
	Isoflurane increased interleukin-11 synthesis in human and mouse proximal tubular cells <i>via</i> TGF-β1 signaling to protect against ischemic acute kidney injury.	
۲	Positive End-expiratory Pressure Increments during Anesthesia in Normal Lung Result in Hysteresis and Greater Numbers of Smaller Aerated Airspaces Maurizio Cereda, Yi Xin, Kiarash Emami, Jessie Huang, Jennia Rajaei, Harrilla Profka, Biao Han, Puttisarn Mongkolwisetwara, Stephen Kadlecek, Nicholas N. Kuzma, Stephen Pickup, Brian P. Kavanagh, Clifford S. Deutschman, and Rahim R. Rizi	1402
	In healthy rats, larger lung volumes due to hysteresis were associated with smaller individual airspace dimensions, suggesting opening of previously nonaerated peripheral airspaces, rather than expansion of already opened airspaces. <i>SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT</i>	

PAIN MEDICINE

DOCKET

Δ

• Pain in 1,000 Women Treated for Breast Cancer: A Prospective Study of Pain Sensitivity and Postoperative Pain

1410

Mari A. Kaunisto, Ritva Jokela, Minna Tallgren, Oleg Kambur, Emmi Tikkanen, Tiina Tasmuth, Reetta Sipilä, Aarno Palotie, Ann-Mari Estlander, Marjut Leidenius, Samuli Ripatti, and Eija A. Kalso

In a study of 1,000 women undergoing breast surgery for cancer, a small portion of the variance in preoperative response to noxious heat and cold testing could be explained by anxiety, the presence of chronic pain, and the number of previous operations. There was a weak correlation between response to experimental pain testing and acute postoperative pain, with largely similar predictive factors across both. *SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT*

♦ ● Effect of Catechol-o-methyltransferase-gene (COMT) Variants on Experimental and Acute Postoperative Pain in 1,000 Women undergoing Surgery for Breast Cancer 1422 Oleg Kambur, Mari A. Kaunisto, Emmi Tikkanen, Suzanne M. Leal, Samuli Ripatti, and Eija A. Kalso In a more complete assessment of variants in the COMT gene in 1,000 women undergoing surgery for breast cancer, there were weak associations with sensitivity to heat or cold pain. The lack of association of COMT variants with acute postoperative opioid requirement raises questions regarding the clinical relevance of this gene in acute postoperative pain. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT 🌐 🚾 Survey Criteria for Fibromyalgia Independently Predict Increased Postoperative Opioid Consumption after Lower-extremity Joint Arthroplasty: A Prospective, Observational **Cohort Study** 1434 Chad M. Brummett, Allison M. Janda, Christa M. Schueller, Alex Tsodikov, Michelle Morris, David A. Williams, and Daniel J. Clauw Even though less than 9% of participants met "criteria" for fibromyalgia, increasing degrees of fibromyalgia-like symptoms were independently predictive of increased postoperative opioid requirements. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT Lidocaine Patch (5%) in Treatment of Persistent Inguinal Postherniorrhaphy Pain: A Randomized, Double-blind, Placebo-controlled, Crossover Trial 1444 Joakim M. Bischoff, Marian Petersen, Nurcan Üçeyler, Claudia Sommer, Henrik Kehlet, and Mads U. Werner Lidocaine patches (5%) caused pressure pain threshold increases when compared with placebo patch treatment. However, lidocaine patch treatment did not lead to decreases in summed pain intensity differences. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT 🔶 🚾 Pregabalin Rectifies Aberrant Brain Chemistry, Connectivity, and Functional Response in Chronic Pain Patients 1453 Richard E. Harris, Vitaly Napadow, John P. Huggins, Lynne Pauer, Jieun Kim, Johnson Hampson, Pia C. Sundgren, Bradley Foerster, Myria Petrou, Tobias Schmidt-Wilcke, and Daniel J. Clauw Using three complementary imaging techniques (proton magnetic resonance spectroscopy, functional magnetic resonance imaging, and functional connectivity) in chronic pain patients with fibromyalgia, it was shown that pregabalin treatment reduced brain insula glutamate levels and was associated with decreased connectivity of this structure to the default mode network. ■ CLASSIC PAPERS REVISITED Therapeutic Benefit of the Anesthesiologist-Patient Relationship 1465 Lawrence D. Egbert and Stephen H. Jackson This article is a revisiting of original material published as: Egbert LD, Battit GE, Turndorf H, Beecher HK: The value of the preoperative visit by an anesthetist: A study of doctor-patient rapport. JAMA 1963; 185:553-5. EDUCATION **IMAGES IN ANESTHESIOLOGY** Icteric Vocal Cords Recorded during Video Laryngoscopy 1469 Richard M. Cooper

Anomalous Origin of the Left Pulmonary Artery from the Right Pulmonary Artery (Pulmonary Artery Sling)

Ahmad Zabad and Mohanad Shukry

DOCKE

1470

DOCKET A L A R M



Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.