



ANESTHESIOLOGY

The Journal of the American Society of Anesthesiologists, Inc. • anesthesiology.org



ANESTHESIOLOGY





ON THE COVER:

We continue our end-of-the-year special focus on an institution which has developed and maintained research excellence in the specialty. This year a majority of the December issue comes from the department of anesthesiology at the University of Michigan (Ann Arbor, Michigan). (The heli-photo of the University of Michigan medical center was provided by Dale Fisher.)

\Diamond	THIS MONTH IN ANESTHESIOLOGY	1
	EDITORIAL VIEWS	
M	Translational Research and the Perioperative Phenotype: University of Michigan Kevin K. Tremper	123
M	Increased Risk of Awareness under Anesthesia: An Issue of Consciousness or of Memory? Kane O. Pryor and Hugh C. Hemmings, Jr.	123
M	Complications of Laryngeal Masks in Children: Big Data Comes to Pediatric Anesthesia Ronald S. Litman	123
	Neural and Immune Consequences of Traumatic Brain Injury: Does Propofol Reduce the Impact? Robert D. Sanders, Mark Coburn, and Pratik P. Pandharipande	124
	The Scientific Journey to Predicting and Preventing Postoperative Pain: Recalling Dr. Wall's Stories along the Way Samuel A. McLean	124
M	"Seeing" How Our Drugs Work Brings Translational Added Value Irene Tracey	124
	Equivalent Efficacy of Hydroxyethyl Starch 130/0.4 and Human Serum Albumin: If Nothing Is the Same, Is Everything Different? The Importance of Context in Clinical Trials and Statistics Richard B. Weiskopf	124
	SPECIAL ARTICLES	
	Altered States: Psychedelics and Anesthetics Eduardo F. Icaza and George A. Mashour	125

- Refers to This Month in Anesthesiology
- Refers to Editorial Views
- ▼ This article is from or about the University of Michigan

- See Supplemental Digital Content
- CME Article



CONTENTS

 \Diamond

	Thomas R. Vetter, Nataliya V. Ivankova, Lee A. Goeddel, Gerald McGwin, Jr., and Jean-Francois Pittet for the UAB Perioperative Surgical Home Group	1261
	PERIOPERATIVE MEDICINE	
♦ M	Increased Risk of Intraoperative Awareness in Patients with a History of Awareness Amrita Aranake, Stephen Gradwohl, Arbi Ben-Abdallah, Nan Lin, Amy Shanks, Daniel L. Helsten, David B. Glick, Eric Jacobsohn, Alex J. Villafranca, Alex S. Evers, Michael S. Avidan, and George A. Mashour	1275
	In a matched cohort analysis of patients drawn from three trials including over 25,000 patients, those with a history of intraoperative awareness had a five-fold increased incidence of awareness compared with propensity-matched controls who did not have a history of awareness. Anesthetic management did not differ between the cohorts; in view of the likely increased risk of awareness, clinicians should consider modifying anesthetic management in patients with a history of awareness.	
• M		
	A Study of Clinical Predictors and Outcomes Michael R. Mathis, Bishr Haydar, Emma L. Taylor, Michelle Morris, Shobha V. Malviya, Robert E. Christensen, Satya-Krishna Ramachandran, and Sachin Kheterpal	1284
	This study evidenced that laryngeal mask airway failures occur in 0.86% of 11,910 pediatric anesthesia cases with its planned use, and are independently associated with ear/nose/throat procedures, admission status, prolonged surgical duration, airway abnormalities, and patient transport.	
•	Six Percent Hydroxyethyl Starch 130/0.4 (Voluven®) versus 5% Human Serum Albumin for Volume Replacement Therapy during Elective Open-heart Surgery in Pediatric Patients Philippe Van der Linden, Andrée De Villé, Anna Hofer, Martina Heschl, and Hans Gombotz	1296
	This randomized clinical trial demonstrated equivalent efficacy of human albumin <i>versus</i> hydroxyethyl starch for volume replacement in 61 children undergoing cardiac surgery. Although the trial was not powered to assess safety, outcomes measures were not different between the groups.	
M	Patient Selection for Day Case-eligible Surgery: Identifying Those at High Risk for	1310
	Major Complications Michael R. Mathis, Norah N. Naughton, Amy M. Shanks, Robert E. Freundlich, Christopher J. Pannucci, YiJia Chu, Jason Haus, Michelle Morris, and Sachin Kheterpal	
	In a review of nearly 250,000 cases in the National Surgery Quality Improvement Program, early perioperative morbidity and mortality occurred in approximately 1:1,000 patients. Predictors for morbidity and mortality were overweight or obesity, chronic obstructive pulmonary disease, history of transient ischemic attack/stroke, hypertension, prior cardiac surgical intervention, and prolonged operative time.	
♦ M	Predictors of Survival from Perioperative Cardiopulmonary Arrests: A Retrospective Analysis	
	of 2,524 Events from the Get With The Guidelines-Resuscitation Registry Satya Krishna Ramachandran, Jill Mhyre, Sachin Kheterpal, Robert E. Christensen, Kristen Tallman, Michelle Morris, and Paul S. Chan, for the American Heart Association's Get With The Guidelines-Resuscitation Investigators	1322
	One third of patients with perioperative cardiac arrests survive to hospital discharge. Two thirds of survivors have good neurological outcome. Shockable rhythms and asystolic arrests were associated with significant location-specific survival variance.	
M	Perioperative Metoprolol and Risk of Stroke after Noncardiac Surgery George A. Mashour, Milad Sharifpour, Robert E. Freundlich, Kevin K. Tremper, Amy Shanks, Brahmajee K. Nallamothu, Phillip E. Vlisides, Adam Weightman, Lisa Matlen, Janna Merte, and Sachin Kheterpal	1340
	In this retrospective study, routine use of preoperative and intraoperative metoprolol—but not other β blockers—was associated with increased risk of stroke after noncardiac surgery. These data suggest that results of the POISE trial may	
· L	CT	

An Analysis of Methodologies That Can Be Used to Validate if a Perioperative Surgical

Home Improves the Patient-centeredness, Evidence-based Practice, Quality, Safety, and Value



CONTENTS

Reconfiguration of Network Hub Structure after Propofol-induced Unconsciousness Heonsoo Lee, George A. Mashour, Gyu-Jeong Noh, Seunghwan Kim, and UnCheol Lee	1347
Propofol reconfigures the brain's network hub structure. Reconfiguration of hub structure may explain the observed loss of frontal-parietal feedback connectivity.	
Incidence, Predictors, and Outcome of Difficult Mask Ventilation Combined with Difficult Laryngoscopy: A Report from the Multicenter Perioperative Outcomes Group Sachin Kheterpal, David Healy, Michael F. Aziz, Amy M. Shanks, Robert E. Freundlich, Fiona Linton, Lizabeth D. Martin, Jonathan Linton, Jerry L. Epps, Ana Fernandez-Bustamante, Leslie C. Jameson, Tyler Tremper, and Kevin K. Tremper, on behalf of the Multicenter Perioperative Outcomes Group (MPOG) Perioperative Clinical Research Committee	1360
This study determined incidence of difficult mask ventilation combined with difficult laryngoscopy to be 0.4% of 176,679 adult cases, and succeeded in identifying 12 independent predictors for the critical situation.	
CRITICAL CARE MEDICINE	
Propofol Limits Microglial Activation after Experimental Brain Trauma through Inhibition of Nicotinamide Adenine Dinucleotide Phosphate Oxidase Tao Luo, Junfang Wu, Shruti V. Kabadi, Boris Sabirzhanov, Kelsey Guanciale, Marie Hanscom, Juliane Faden, Katherine Cardiff, Charles Jeremy Bengson, and Alan I. Faden	1370
Propofol reduces functional deficit after traumatic brain injury, in an animal model. These neuroprotective effects are mediated in part by inhibition of microglial nicotinamide adenine dinucleotide phosphate oxidase.	
Critical Role of Interleukin-11 in Isoflurane-mediated Protection against Ischemic Acute Kidney Injury in Mice Ahrom Ham, Mihwa Kim, Joo Yun Kim, Kevin M. Brown, James Yeh, Vivette D. D'Agati, and H. Thomas Lee	1389
Isoflurane increased interleukin-11 synthesis in human and mouse proximal tubular cells <i>via</i> TGF-β1 signaling to protect against ischemic acute kidney injury.	
Positive End-expiratory Pressure Increments during Anesthesia in Normal Lung Result in Hysteresis and Greater Numbers of Smaller Aerated Airspaces Maurizio Cereda, Yi Xin, Kiarash Emami, Jessie Huang, Jennia Rajaei, Harrilla Profka, Biao Han, Puttisarn Mongkolwisetwara, Stephen Kadlecek, Nicholas N. Kuzma, Stephen Pickup, Brian P. Kavanagh, Clifford S. Deutschman, and Rahim R. Rizi	1402
In healthy rats, larger lung volumes due to hysteresis were associated with smaller individual airspace dimensions, suggesting opening of previously nonaerated peripheral airspaces, rather than expansion of already opened airspaces. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT	
PAIN MEDICINE	
Pain in 1,000 Women Treated for Breast Cancer: A Prospective Study of Pain Sensitivity and Postoperative Pain	1410
	Heonsoo Lee, George A. Mashour, Gyu-Jeong Noh, Seunghwan Kim, and UnCheol Lee Propofol reconfigures the brain's network hub structure. Reconfiguration of hub structure may explain the observed loss of frontal-parietal feedback connectivity. Incidence, Predictors, and Outcome of Difficult Mask Ventilation Combined with Difficult Laryngoscopy: A Report from the Multicenter Perioperative Outcomes Group Sachin Kheterpal, David Healy, Michael F. Aziz, Arny M. Shanks, Robert E. Freundlich, Fiona Linton, Lizabeth D. Martin, Jonathan Linton, Jerry L. Epps, Ana Fernandez-Bustamante, Leslie C. Jameson, Tyler Tremper, and Kevin K. Tremper, on behalf of the Multicenter Perioperative Outcomes Group (MPOG) Perioperative Clinical Research Committee This study determined incidence of difficult mask ventilation combined with difficult laryngoscopy to be 0.4% of 176.679 adult cases, and succeeded in identifying 12 independent predictors for the critical situation. CRITICAL CARE MEDICINE Propofol Limits Microglial Activation after Experimental Brain Trauma through Inhibition of Nicotinamide Adenine Dinucleotide Phosphate Oxidase Tao Luo, Junfang Wu, Shruti V. Kabadi, Boris Sabirzhanov, Kelsey Guanciale, Marie Hanscom, Juliane Faden, Katherine Cardiff, Charles Jeremy Bengson, and Alan I. Faden Propofol reduces functional deficit after traumatic brain injury, in an animal model. These neuroprotective effects are mediated in part by inhibition of microglial nicotinamide adenine dinucleotide phosphate oxidase. Critical Role of Interleukin-11 in Isoflurane-mediated Protection against Ischemic Acute Kidney Injury in Mice Ahrom Ham, Mihwa Kim, Joo Yun Kim, Kevin M. Brown, James Yeh, Vivette D. D'Agati, and H. Thomas Lee Isoflurane increased interleukin-11 synthesis in human and mouse proximal tubular cells via TGF-β1 signaling to protect against ischemic acute kidney injury. Positive End-expiratory Pressure Increments during Anesthesia in Normal Lung Result in Hysteresis and Greater Numbers of Smaller Aerated Airspaces Mauri

Mari A. Kaunisto, Ritva Jokela, Minna Tallgren, Oleg Kambur, Emmi Tikkanen, Tiina Tasmuth, Reetta Sipilä, Aarno Palotie, Ann-Mari Estlander, Marjut Leidenius, Samuli Ripatti, and Eija A. Kalso

In a study of 1,000 women undergoing breast surgery for cancer, a small portion of the variance in preoperative response to noxious heat and cold testing could be explained by anxiety, the presence of chronic pain, and the number of previous operations. There was a weak correlation between response to experimental pain testing and acute postoperative pain, with largely similar predictive factors across both. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT



CONTENTS

◇◆	Effect of Catechol-o-methyltransferase-gene (COMT) Variants on Experimental and Acute Postoperative Pain in 1,000 Women undergoing Surgery for Breast Cancer Oleg Kambur, Mari A. Kaunisto, Emmi Tikkanen, Suzanne M. Leal, Samuli Ripatti, and Eija A. Kalso	1422
	In a more complete assessment of variants in the <i>COMT</i> gene in 1,000 women undergoing surgery for breast cancer, there were weak associations with sensitivity to heat or cold pain. The lack of association of COMT variants with acute postoperative opioid requirement raises questions regarding the clinical relevance of this gene in acute postoperative pain. <i>SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT</i>	
M	Survey Criteria for Fibromyalgia Independently Predict Increased Postoperative Opioid Consumption after Lower-extremity Joint Arthroplasty: A Prospective, Observational Cohort Study Chad M. Brummett, Allison M. Janda, Christa M. Schueller, Alex Tsodikov, Michelle Morris, David A. Williams, and Daniel J. Clauw	1434
	Even though less than 9% of participants met "criteria" for fibromyalgia, increasing degrees of fibromyalgia-like symptoms were independently predictive of increased postoperative opioid requirements. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT	
(4)	Lidocaine Patch (5%) in Treatment of Persistent Inguinal Postherniorrhaphy Pain: A Randomized, Double-blind, Placebo-controlled, Crossover Trial Joakim M. Bischoff, Marian Petersen, Nurcan Üçeyler, Claudia Sommer, Henrik Kehlet, and Mads U. Werner	1444
	Lidocaine patches (5%) caused pressure pain threshold increases when compared with placebo patch treatment. However, lidocaine patch treatment did not lead to decreases in summed pain intensity differences. SUPPLEMENTAL DIGITAL CONTENT IS AVAILABLE IN THE TEXT	
♦ M	Pregabalin Rectifies Aberrant Brain Chemistry, Connectivity, and Functional Response in Chronic Pain Patients Richard E. Harris, Vitaly Napadow, John P. Huggins, Lynne Pauer, Jieun Kim, Johnson Hampson, Pia C. Sundgren, Bradley Foerster, Myria Petrou, Tobias Schmidt-Wilcke, and Daniel J. Clauw	1453
	Using three complementary imaging techniques (proton magnetic resonance spectroscopy, functional magnetic resonance imaging, and functional connectivity) in chronic pain patients with fibromyalgia, it was shown that pregabalin treatment reduced brain insula glutamate levels and was associated with decreased connectivity of this structure to the default mode network.	
	CLASSIC PAPERS REVISITED	
	Therapeutic Benefit of the Anesthesiologist–Patient Relationship Lawrence D. Egbert and Stephen H. Jackson	1465
	This article is a revisiting of original material published as: Egbert LD, Battit GE, Turndorf H, Beecher HK: The value of the preoperative visit by an anesthetist: A study of doctor–patient rapport. JAMA 1963; 185:553–5.	
	EDUCATION	
IMA	GES IN ANESTHESIOLOGY	
	Icteric Vocal Cords Recorded during Video Laryngoscopy Richard M. Cooper	1469
	Anomalous Origin of the Left Pulmonary Artery from the Right Pulmonary Artery (Pulmonary Artery Sling) Ahmad Zabad and Mohanad Shukry	1470



DOCKET

Explore Litigation Insights



Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

