



SIGMA-ALDRICH

3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

Product Information

L-Cysteine hydrochloride monohydrate Cell culture tested, meets USP testing specifications

Product Number **C6852**

Product Description

Molecular Formula: $C_3H_7NO_2S \cdot HCl \cdot H_2O$
Molecular Weight: 175.6
CAS Number: 7048-04-6

This product is tested for endotoxin levels and suitability for cell culture experiments.

Cysteine is a major biological source of sulfur and is one of the two common sulfur-containing amino acids. The biosynthesis of cysteine occurs through the initial condensation of homocysteine and serine via cystathionine synthase to form cystathionine, which in turn undergoes cleavage by cystathionase to give cysteine and α -ketobutyrate. Cysteine can readily dimerize to form cystine via the oxidation of the thiol side chain residues to give a disulfide covalent bond. The formation of such cystine links between cysteine residues in proteins is an important part of the stabilization of the three-dimensional structure of proteins.¹

An investigation into cysteine and cystine levels in normal and malignant cells with a relationship to γ -cystathionase levels and tumor sensitivity to L-cysteine and cystine depletion has been reported.² The toxicity of human neuronal cell lines to cysteine and its metabolites has been investigated.^{3,4} Cysteine can inhibit the activity of enzymes such as lecithin-cholesterol acyltransferase.⁵

A protocol for the use of cysteine in pulse-chase experiments to study protein expression from an inducible promoter has been reported.⁶

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

This product is soluble in water (100 mg/ml), yielding a clear, colorless solution.

Storage/Stability

Stock solutions of cysteine are relatively stable at acidic pH, especially in degassed solutions. Aqueous solutions of cysteine oxidize readily in air to give cystine at neutral or basic pH.⁷

References

1. Textbook of Biochemistry with Clinical Correlations, Devlin, T. M., ed., Wiley-Liss (New York, NY: 1992), p. 33.
2. Uren, J. R., and Lazarus, H., L-cyst(e)ine requirements of malignant cells and progress toward depletion therapy. *Cancer Treat. Rep.*, **63(6)**, 1073-1079 (1979).
3. Parsons, R. B., et al., Toxicity of cysteine and cysteine sulphinic acid to human neuronal cell-lines. *J. Neurol. Sci.*, **152** (Suppl 1), S62-66 (1997).
4. Parsons, R. B., et al., In vitro effect of the cysteine metabolites homocysteic acid, homocysteine and cysteic acid upon human neuronal cell lines. *Neurotoxicology*, **19(4-5)**, 599-603 (1998).
5. Albers, J. J., et al., Isolation, characterization, and assay of lecithin-cholesterol acyltransferase. *Methods Enzymol.*, **129**, 763-783 (1986).
6. *Molecular Cloning: A Laboratory Manual*, 3rd ed., Sambrook, J. and Russell, D. W., CSHL Press (Cold Spring Harbor, NY: 2001), pp. 15.18-15.19.
7. *Data for Biochemical Research*, 3rd ed., Dawson, R. M. C., et al., Oxford University Press (New York, NY: 1986), pp. 12-13.

GCY/NSB 5/06

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.