

L-CYSTEINE HYDROCHL x + v

https://web.archive.org/web/20170403170533/http://drugsdb.eu/drug.php?d=L-cystein

INTERNET ARCHIVE WayBack Machine 2 captures 24 Aug 2016 - 3 Apr 2017

www.drugsDB.eu With permission of U.S. National Library of Medicine

DRUGS DataBase

You are here: Home > Prescription(RX) Drugs > L > L-cysteine Hydrochloride (Sandoz Inc) Loading

Name:	L-cysteine Hydrochloride
Manufacturer:	Sandoz Inc
Category:	Prescription Marketed Drugs

Prescription Marketed Drugs Alphabetically
A| B| C| D| E| F| G| H| I| J| K| L| M| N| O| P| Q| R| S| T| U| V| W| X| Y| Z| 0-9

Categories:

Prescription(RX) Drugs
Over-the-counter (OTC) Drugs
Homeopathic Drugs
Animal Drugs
Feedback

L-CYSTEINE HYDROCHLORIDE - cysteine hydrochloride injection, solution
Sandoz Inc
Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

L-Cysteine Hydrochloride Injection, USP
0.5 g/10 mL (50 mg/mL)

PHARMACY BULK PACKAGE
NOT FOR DIRECT INFUSION

DESCRIPTION
L-Cysteine Hydrochloride Injection, USP, 50 mg/mL, is a sterile, nonpyrogenic solution. Each mL contains: 50 mg of L-Cysteine Hydrochloride Monohydrate USP; Water for Injection, USP q.s.; Air replaced with Nitrogen. pH 1.0-2.5
L-Cysteine is a sulfur-containing amino acid. In premixed solutions of crystalline amino acids, cysteine is relatively unstable over time, eventually converting to insoluble cystine. To avoid such precipitation, L-Cysteine Hydrochloride Injection USP is intended to be used as an additive with crystalline Amino Acid Injections immediately prior to administration to the patient.
The structural formula of Cysteine Hydrochloride Monohydrate USP is:

$$\begin{array}{c} \text{H} \\ | \\ \text{HSCH}_2 - \text{C} - \text{COOH} \cdot \text{HCl} \cdot \text{H}_2\text{O} \\ | \\ \text{NH}_2 \end{array}$$

Molecular Weight 175.63 **Molecular Formula** C₃H₇NO₂S·HCl·H₂O

L-CYSTEINE HYDROCHL X + v

https://web.archive.org/web/20170403170533/http://drugsdb.eu/drug.php?d=L-cystein

Wayback Machine 2 captures 24 Aug 2016 - 3 Apr 2017

CLINICAL PHARMACOLOGY

L-Cysteine is synthesized from methionine via the trans-sulfuration pathway in the adult, but newborn infants lack the enzyme necessary to effect this conversion. Therefore, L-Cysteine is generally considered to be an essential amino acid in infants.

INDICATIONS AND USAGE

L-Cysteine Hydrochloride Injection, USP is intended for use only after dilution as an additive to Crystalline Amino Acid Injections to meet the intravenous amino acid nutritional requirements of infants receiving total parenteral nutrition.

CONTRAINDICATIONS

This preparation should not be used in patients with hepatic coma or metabolic disorders involving impaired nitrogen utilization.

WARNINGS

Peripheral intravenous infusion of amino acids may induce a rise in blood urea nitrogen (BUN) especially in patients with impaired hepatic or renal function. Appropriate laboratory tests should be performed periodically and infusion discontinued if BUN levels exceed normal postprandial limits and continue to rise. It should be noted that a modest rise in BUN normally occurs as a result of increased protein intake.

Administration of amino acid solutions to a patient with hepatic insufficiency may result in serum amino acid imbalances, metabolic alkalosis, prerenal azotemia, hyperammonemia, stupor and coma.

Administration of amino acid solutions in the presence of impaired renal function may augment an increasing BUN, as does any protein dietary component.

Solutions containing sodium ion should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency, and in clinical states in which there exists edema with sodium retention.

Solutions which contain potassium ion should be used with great care, if at all, in patients with hyperkalemia, severe renal failure and in conditions in which potassium retention is present.

Solutions containing acetate ion should be used with great care in patients with metabolic or respiratory alkalosis. Acetate should be administered with great care in those conditions in which there is an increased level or an impaired utilization of this ion such as severe hepatic insufficiency.

Hyperammonemia is of special significance in infants, as it can result in mental retardation. Therefore it is essential that blood ammonia levels be measured frequently in infants.

Instances of asymptomatic hyperammonemia have been reported in patients without overt liver dysfunction. The mechanisms of this reaction are not clearly defined but may involve genetic defects and immature or subclinically impaired liver function.

Frequent Clinical Evaluation and Laboratory Determinations are Necessary for Proper Monitoring During Administration. Blood studies should include glucose, urea nitrogen, serum electrolytes, ammonia, cholesterol, acid-base balance, serum proteins, kidney and liver function tests, osmolarity and hemogram. White blood count and blood cultures are to be determined if indicated. Urinary osmolarity and glucose should be determined frequently.

Safe use during pregnancy has not been established, therefore, infusion of amino acids should be undertaken during pregnancy only when this is deemed essential to the patients' welfare, as judged by the physician.

WARNING: *This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.*

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration

L-CYSTEINE HYDROCHL

https://web.archive.org/web/20170403170533/http://drugsdb.eu/drug.php?d=L-cysteine

INTERNET ARCHIVE
Wayback Machine
2 captures
24 Aug 2016 - 3 Apr 2017

Go AUG APR MAY
2016 03 2017 2018

PRECAUTIONS
Special care must be taken when administering hypertonic glucose to provide calories in diabetic or prediabetic patients.
Because of its antianabolic activity, concurrent administration of tetracycline may reduce the nitrogen sparing effects of infused amino acids.
Do not withdraw venous blood for blood chemistries through the peripheral infusion site, as interference with estimations of nitrogen containing substances may occur.
Intravenous feeding regimens which include amino acids should be used with caution in patients with a history of renal disease, pulmonary disease, or with cardiac insufficiency so as to avoid excessive fluid accumulation.
The effect of infusion of amino acids, without dextrose, upon carbohydrate metabolism of children is not known at this time.
Nitrogen intake should be carefully monitored in patients with impaired renal function. For long-term total nutrition, or if a patient has inadequate fat stores, it is essential to provide adequate exogenous calories concurrently with the amino acids. Concentrated dextrose solutions are an effective source of such calories. Such strongly hypertonic nutrient solutions should be administered through an indwelling intravenous catheter with the tip located in the superior vena cava.

ADVERSE REACTIONS
Local reactions consisting of a warm sensation, erythema, phlebitis and thrombosis at the infusion site have occurred with peripheral intravenous infusion of amino acids, particularly if the other substances, such as antibiotics, are also administered through the same site. In such cases the infusion site should be changed promptly to another vein. Use of large peripheral veins, inline filters, and slowing the rate of infusion may reduce the incidence of local venous irritation. Electrolyte additives should be spread throughout the day. Irritating additive medications may need to be injected at another venous site. Generalized flushing, fever and nausea also have been reported during peripheral infusions of amino acid solutions.

Drug Abuse and Dependence
None known.

DOSAGE AND ADMINISTRATION
L-Cysteine Hydrochloride Injection USP is intended for use only after dilution in Crystalline Amino Acid Injection. Each 0.5 gram of L-Cysteine Hydrochloride Monohydrate should be combined aseptically with 12.5 grams of Crystalline Amino Acid Injection, such as that present in 250 mL of 5% Crystalline Amino Acid Injection. The admixture is then diluted with 250 mL of dextrose 50% or such lesser volume as indicated. Equal volumes of 5% Crystalline Amino Acid Injection and dextrose 50% produce a final solution which contains Crystalline Amino Acid Injection 2.5% in dextrose 25%, which is suitable for administration by central venous infusion. Administration of the final admixture should begin within one hour of mixing. Otherwise, the admixture should be refrigerated immediately and used within 24 hours of the time of mixing. For the recommended rate of administration, see the Crystalline Amino Acid Injection package insert.
Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

DIRECTIONS FOR PROPER USE OF PHARMACY BULK PACKAGE
The pharmacy bulk package is for use in a Pharmacy Admixture Service only.
Use of this product is restricted to a suitable work area, such as a laminar flow hood. Prior to entering the vial, remove the flip-off seal and cleanse the rubber closure with a suitable antiseptic agent.
The container closure may be penetrated only one time, utilizing a suitable sterile transfer device or dispensing set which allows measured distribution of the contents. The date and time the vial was initially opened should be recorded in the space provided on the label. Transfer individual doses(s) to appropriate intravenous infusion solutions. Use of a syringe with needle is not recommended. Multiple entries increase the potential of microbial and particulate contamination.

L-CYSTEINE HYDROCHL X + v

https://web.archive.org/web/20170403170533/http://drugsdb.eu/drug.php?d=L-cystein

Wayback Machine 2 captures he red 24 Aug 2016 - 3 Apr 2017

entries increase the potential of microbial and particulate contamination.
The withdrawal of container contents should be accomplished without delay using aseptic technique.
However, should this not be possible, a maximum time of 4 hours from initial closure entry is permitted to complete fluid transfer operations.

RECOMMENDED STORAGE CONDITIONS AFTER OPENING
Keep under laminar flow hood at room temperature. Any unused portion of the vial must be discarded within 4 hours after initial entry.

HOW SUPPLIED
L-Cysteine Hydrochloride Injection, USP (50 mg/mL) is supplied as follows:

PHARMACY BULK PACKAGE

NDC Number	Volume
66758-005-01	50 mL
66758-005-02	5 x 50 mL

Also available as:


SINGLE DOSE VIAL

NDC Number	Volume
66758-004-01	10 mL
66758-004-02	10 x 10 mL


Store at controlled room temperature 15°-30°C (59°-86°F) Do not freeze.
For Sandoz Inc. Customer Service, call [1-800-525-8747](tel:1-800-525-8747)

Rx only
Manufactured for:
SANDOZ
Princeton, NJ 08540
Revised: November 2009
L-028-00

Package Label - Principal Display Panel - 50 mL Vial
SANDOZ
Rx only NDC 66758-005-01
L-Cysteine
Hydrochloride Injection, USP
PHARMACY BULK PACKAGE
NOT FOR DIRECT INFUSION
50 mg/mL
For IV Use Only After Dilution
Do Not Dispense As A Unit
50 mL
L-018-00
PHARMACY BULK PACKAGE



Package Label - Principal Display Panel - 50 mL Vial Carton
SANDOZ
5 x 50 mL Vials **NDC 66758-005-02**
L-Cysteine Hydrochloride
Injection, USP
50 mg/mL
Pharmacy Bulk Package
Not For Direct Infusion
For IV Use Only After Dilution
Rx only



L-CYSTEINE HYDROCHLORIDE			
l-cysteine hydrochloride injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66758-005
Route of Administration	INTRAVENOUS	DEA Schedule	
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CYSTEINE HYDROCHLORIDE (CYSTEINE)	CYSTEINE HYDROCHLORIDE	50 mg in 1 mL	
Inactive Ingredients			
Ingredient Name	Strength		
WATER			
Product Characteristics			
Color		Score	
Shape		Size	
Flavor		Imprint Code	
Contains			
Packaging			
#Item Code	Package Description	Multilevel Packaging	
1 NDC:66758-005-02	5 VIAL, PHARMACY BULK PACKAGE (VIAL) in 1 CARTON	contains a VIAL, PHARMACY BULK PACKAGE (66758-005-01)	
1 NDC:66758-005-01	50 mL in 1 VIAL, PHARMACY BULK PACKAGE	This package is contained within the CARTON (66758-005-02)	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		12/01/2008	

Labeler - Sandoz Inc (005387188)
 Revised: 06/2010 Sandoz Inc

Source: <http://dailymed.nlm.nih.gov>
 Reproduced with permission of U.S. National Library of Medicine

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.