

Drug Facts

Active ingredients (in each tablet)	Purpose
Aspirin 500 mg.....	Analgesic
Citric acid 1000 mg.....	Antacid
Sodium bicarbonate (heat-treated) 1985 mg.....	Antacid

Uses for the relief of:

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if you have

- asthma
- ulcers
- bleeding problems
- stomach problems that last or come back frequently, such as heartburn, upset stomach, or pain

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- on a sodium-restricted diet
- taking a prescription drug for anticoagulation (blood thinning), diabetes, gout, or arthritis